

UNIVERSIDADE TÉCNICA DE LISBOA INSTITUTO SUPERIOR TÉCNICO

Thermodynamics as a Substantive and Formal Theory for the Analysis of Economic and Biological Systems

Tania Alexandra dos Santos Costa e Sousa (Mestre)

Dissertação para obtenção do Grau de Doutor em Engenharia do Ambiente

DOCUMENTO PROVISÓRIO

Abstract

In this dissertation, we use thermodynamic theory to analyse biological and economic systems, according to two methodological approaches. Substantive integration is a methodological approach that consists of making two different fields physically compatible either by reformulating current theories or by building up new theory. For example, thermodynamics must be compatible with the theories that explain and describe economic and biological systems because thermodynamic laws impose constraints on mass, energy and entropy flows. Formal analogy is a methodological approach that consists of the development of an isomorphism between the mathematical formalisms of different fields. For example, thermodynamics has a solid mathematical formal structure that describes equilibrium, non-equilibrium and self-organized systems that provides a possible common framework to describe thermodynamic, biological and economic systems. In this dissertation, we concentrate on the substantive integration between non-equilibrium thermodynamics and the Dynamic Energy Budget (DEB) theory that describes biological systems and on the formal analogy between equilibrium thermodynamics and neoclassical economic theory. In this dissertation, we also build a more systematic description of DEB theory as an intermediate step to a future substantive integration between thermodynamics, economics and biology.

Keywords: Thermodynamics, Formal Analogy, Substantive Integration, Dynamic Energy Budget Theory and Neoclassical Microeconomics

iv

Resumo

Nesta tese, a teoria termodinâmica é usada para analisar os sistemas biológicos e económicos através de duas metodologias. A integração substantiva é a metodologia que consiste em tornar dois campos do conhecimento compatíveis, através da reformulação das teorias existentes ou através da construção de nova teoria. Por exemplo, a termodinâmica tem que ser compatível com as teorias que explicam e descrevem os sistemas económicos e biológicos porque as leis termodinâmicas impõem restrições nos fluxos de massa, energia e entropia. A analogia formal é a metodologia que consiste no desenvolvimento de um isomorfismo entre os formalismos matemáticos de diferentes áreas. Por exemplo, a termodinâmica tem uma estrutura matemática formal que descreve os sistemas em equilíbrio, não-equilíbrio e auto-organizados que pode constituir uma base comum para a descrição dos sistemas termodinâmicos, biológicos e económicos. Nesta dissertação, é feita a integração substantiva entre a termodinâmica de não equilíbrio e a teoria 'Dynamic Energy Budget' (DEB) que descreve os sistemas biológicos e é construída uma analogia formal entre a termodinâmica de equilíbrio e a teoria económica neoclássica. Nesta dissertação, é também construída uma descrição mais sistemática da teoria DEB como um passo intermédio para uma futura integração substantiva entre a termodinâmica, a economia e a biologia.

Palavras Chave: Termodinâmica, Analogia Formal, Integração Substantiva, Teoria 'Dynamic Energy Budget' e Microeconomia Neoclássica vi

Samenvatting

In dit proefschrift gebruiken we thermodynamische theorie om biologische en economische systemen te analyseren, volgens twee methodologische benaderingen. De inhoudelijke integratie is een methodologische benadering die twee verschillende terreinen natuurkundig met elkaar vereenigbaar maakt door bestaande theoriën te herformuleren, of nieuwe op te zetten. De thermodynamika moet vereenigbaar zijn met theoriën die economische en biologische systemen beschrijven en verklaren, omdat haar wetmatigheden randvoorwaarden opleggen op massa, energie en entropie stromen. Een formele analogie is een methodologische benadering die bestaat uit de ontwikkeling van een gelijkvormig wiskundig formalisme voor verschillende terreinen. De thermodynamika heeft een solide formele struktuur die systemen beschrijft die wel of niet in evenwicht verkeren of zelf-organiserend zijn en kunnen worden toegepast in de thermodynamika, de biologie en de economie. In dit proefschrift concentreren we ons op een inhoudelijke integratie tussen niet-evenwichts thermodynamika en biologische systemen zoals die vastgelegd worden door de Dynamische Energie Budget (DEB) theorie en op de formele analogie tussen evenwichts-thermodynamika en economische systemen die beschreven worden door de nieuw-klassieke economische theorie. We zetten ook een meer systematische specificatie van de DEB theorie op als tussenstap voor een toekomstige inhoudelijke integratie tussen thermodynamika, economie en biologie.

Trefwoorden: Thermodynamika, Formele Analogie, Inhoudelijke Integratie, Dynamische Energie Budget Theorie en Nieuw-klassieke Microeconomie viii

Acknowledgements

First, I would like to thank Tiago Domingos and Bas Kooijman for the stimulating and fruitful discussions indoors and outdoors and for everything that I learnt with them during these years. I also would like to thank Paulo Ferrão - this thesis would have not been possible without him.

I thank Prof. Delgado Domingos and Prof. Jorge Calado for having contributed in a fundamental way to the building up of my knowledge in thermodynamics.

I thank the people that were part of my working environment in IST, Tiago Domingos, Ana Simões, João Rodrigues, Rui Mota, Ricardo Teixeira, Clara Fiuza, Gonçalo Marques and Cristina Marta for mantaining an inspiring working environment and for the nice weekly discussions and others. I also thank Stephan Baumgartner and Ralph Winkler for helpful comments and suggestions on chapter 2 of this dissertation.

Part of the thesis work was done while I was at the Theoretical Biological Department in the Vrije Universiteit. I want to thank Bas Kooijman and his group for their support and for the nice talks at the 9 o'clock coffee.

I thank Ana Simões, Inês Azevedo, Rui Mota and Ricardo Teixeira for their companionship in the economic courses that we attended.

I thank the financial support from PRODEP III and from FCT through grants POCTI/MGS/47731/2002 and POCI/AMB/55701/2004.

I thank Bas and Trus for their great hospitality in the Netherlands. I also thank Sparky (the dog) for those nice and noisy welcomes.

Finally, I want to thank my family and all my friends. Most important of all I want to thank Tiago, my husband, for everything.

х

Contents

1	Intro	oduction	1	
2	Is Ne	eoclassical Economics Formally Valid?	9	
	Published in <i>Ecological Economics</i> 58 (2006): 160-169			
	2.1	Introduction	10	
	2.2	Is the Formalism of Neoclassical Economics wrong?	12	
	2.3	A Unified Formalism for Neoclassical Economics and Equilibrium Ther-		
		modynamics	14	
	2.4	Discussing the Formalism	19	
	2.5	Conclusions	21	
3	Equi	librium Econophysics	29	
	Published in <i>Physica A</i> 371 (2006): 492-512			
	3.1	Introduction	30	
	3.2	Fundamental Equation and the Equilibrium State	31	
	3.3	Duality Formulation	33	
	3.4	Reversible, Irreversible and Impossible Processes	35	
	3.5	Many-step Processes	36	
	3.6	Legendre Transforms	38	
	3.7	Elasticities	40	
	3.8	Maxwell Relations	43	
	3.9	Stability Conditions and the Le Chatelier Principle	45	
	3.10	Equations of State and Integrability	47	
	3.11	First Order Phase Transitions	48	
	3.12	Conclusions	54	
	3.13	Appendices	57	

xii

4	The	The Thermodynamics of Organisms				
	Publ	Published in Physical Review E 74 (2006): 1-15				
	4.1	Introduction	66			
	4.2	Thermodynamic System	67			
	4.3	Thermodynamic Analysis	71			
	4.4	Empirical Application	82			
	4.5	Conclusions	90			
	4.6	Appendices	91			
5 A Formal Metabolic Theory of Life						
	Submitted					
	5.1	Introduction	99			
	5.2	Empirical Patterns	100			
	5.3	Theory on Metabolic Organization	102			
	5.4	Theory on Parameter Values	118			
	5.5	Conclusions	122			

Chapter 1

Introduction

In this dissertation, we use thermodynamic theory to analyse economic and biological systems, according to two methodological approaches. Substantive integration is a methodological approach that consists of making two different fields compatible, e.g., thermodynamics and biology or thermodynamics and economics, either by reformulating current theories or by building up new theory [6, 14]. For example, theories that explain and describe economic and biological systems must be compatible with thermodynamics because thermodynamic laws impose constraints on mass, energy and entropy flows. A formal analogy is a methodological approach of developing an isomorphism between the mathematical formalisms of different fields [6, 14]. For example, thermodynamics has a solid mathematical formal structure that describes equilibrium, non-equilibrium and self-organized systems that provides a possible common framework to describe thermodynamic, biological and economic systems.

In this dissertation, we concentrate 1) on the substantive integration between thermodynamics and biological systems, 2) on the formal analogy between thermodynamics and economic systems and 3) on the formalization of the Dynamic Energy Budget (DEB) Theory - a theory for biological systems as a prelude to a formal analogy between economic and biological systems.

In a formal analogy between thermodynamic, biological and economic systems, a mathematical framework is developed that simultaneously describes these systems. Formal analogies are important because 1) they reduce the number of logical reasonings used in the different fields of knowledge, 2) they provide a similar interpretation for different phenomena and 3) the knowledge that exists in one field is transposed to another contributing to new developments. However, formal analogies have to be done with great care identifying whether the conditions that have to be met in the original field of knowledge

are also met in the field where the analogy is taken. In this dissertation, an isomorphism is established between equilibrium macroscopic thermodynamics and neoclassical microeconomics using a mathematical framework based on the Gibbs-Tisza-Callen approach to equilibrium thermodynamics [2, 4, 19].

Equilibrium thermodynamics characterizes a macroscopic system that evolves in such a way that some thermodynamic potential is optimized. The existence of this extremum keeps the system stable because, when some fluctuation drives the system away from equilibrium, the optimization behaviour brings it back [12]. Neoclassical microeconomics describes the consumer maximizing his utility or the firm maximizing its profit. We focus on the neoclassical microeconomic description of the consumer's behavior because it is the standard theory used to describe the behavior of firms and consumers [15, 22].

In a substantive integration, thermodynamic laws are applied to economic and biological processes to obtain the constraints imposed on the relations between inputs, outputs and the system's state. Also, new theory compatible with thermodynamics can be developed to describe economic and biological processes. Substantive integration is important because it unifies the understanding of biological and economic systems as thermodynamic systems and therefore of the constraints imposed on them. In this dissertation, non-equilibrium thermodynamics [5, 8, 12, 16, 18], is used to make a substantive integration with biological systems.

Non-equilibrium thermodynamics should be used as a substantive theory because economic and biological systems are open thermodynamic systems kept out of thermodynamic equilibrium due to mass and energy flows. In this work, we take further the substantive integration between the most general framework of non-equilibrium thermodynamics and an organism described by the Dynamic Energy Budget (DEB) theory.

DEB theory describes the way organisms acquire and use matter and energy [13] linking all levels of organization, from membrane physiology to ecosystem dynamics. We use DEB theory to describe biological systems because 1) it has been successfully tested for many organisms including bacteria, yeasts, unicellular algae, lichens, fishes, nematodes, birds and mammals, 2) it is applicable to all species and 3) it is based on the description of mass and energy flows. In this dissertation, we build an axiomatic formalism to describe metabolism that is fully compatible with DEB theory. This formalization of DEB theory and the steps taken in the substantive integration between non-equilibrium thermodynamics and DEB theory facilitate the development of new economic theory compatible with thermodynamics by later establishing an analogy between economic and biological systems.

Introduction

A good example to ilustrate the simultaneous existence of a substantive integration between the firm and thermodynamics and a formal analogy between the organism and the firm is the field of industrial metabolism. In this field, there is a substantive integration because economic flows, i.e., the whole integrated collection of physical processes that convert raw materials and energy, plus labour, into finished products and wastes, are described as mass and energy flows [1] and there is a formal analogy because the firm is considered to be the economic analogue of a living organism [1].

Outline of the Dissertation

This dissertation comprises four chapters. Chapter 2 explores the formal analogy between microeconomics and equilibrium thermodynamics. Here, we show that neoclassical microeconomics is formally analogous to equilibrium thermodynamics proving that neoclassical economics has irreversibility embedded in it and we obtain a definition of irreversibility in economic behavior.

Chapter 3 is an elaboration on chapter 2. Here, the isomorphism between thermodynamics and economics is taken much further including the definition of Legendre transforms, Maxwell relations, Le Chatelier Principle and first order phase transitions in economic systems. The equivalence between variables used in microeconomics and thermodynamics is established, new results are obtained in both fields and the differences between the two optimization problems are systematized.

Chapter 4 addresses the substantive integration between thermodynamics and biology. Here, non-equilibrium thermodynamics is applied to an organism that follows the rules of DEB theory. Results include thermodynamic constraints on the behaviour and on the thermodynamic properties of aerobic and non-aerobic organisms. DEB's concept of reserve density is shown to be crucial in discussions concerning entropy production in organisms.

Chapter 5 formalizes into assumptions and propositions DEB theory and links it in a precise way with the empirical patterns that characterize metabolism. This formalization makes it easier to discuss the consequences of alternative metabolic theories. Results also include an alternative proof of a crucial concept in DEB theory and a novel way to present DEB's theory on parameter values.

In this dissertation, we have concluded that:

• The existence of a formal analogy does not imply a substantive integration and viceversa. This is evident in the relation between thermodynamics and microeconomics: a formal analogy between the two does not mean that microeconomic theory is compatible with thermodynamic laws.

- Formal analogies contribute to the solution of problems that exist in the theoretical structure of a scientific discipline. For example, the discussion on whether the microeconomic formalism has irreversibility embedded in it is clarified by the isomorphism established in chapter 2.
- Formal analogies contribute to obtain new results in a discipline. For example, the relationship between elasticities in economics and the distinction between mathematical integrability and optimization behaviour in thermodynamics.
- The identification of an isomorphism between two or more different disciplines simplifies the learning process. A clear example is presented in chapter 3 where the concept of economic integrability is shown to be equivalent to the Generalized Le Chatelier Principle.
- The substantive integration between thermodynamics and other disciplines contributes to new thermodynamic results about the systems studied. For example, in chapter 4 we obtain the result that anaerobic organisms have a higher flexibility in their thermodynamic behaviour.

Outlook

We used equilibrium thermodynamics to build up a formal analogy with one consumer in chapters 2 and 3. The formal analogy can be extended to a profit maximizing firm or to a social welfare maximizing central planner. Equilibrium thermodynamics can also be used to establish a formal analogy with the optimal foraging strategy in ecology [21].

The transition from maximizing utility to maximizing social welfare is not straightforward because the equilibrium states that result from multiple optimizing agents are different from the equilibrium state that results from an aggregate optimizing function. The set of conditions under which optimizing consumers and producers lead to the maximization of social welfare is a crucial issue in economics [10]. The same issue can be raised in ecological systems about the relation between optimizing organisms and the optimization of an ecosystem [11].

Statistical physics [9, 20] explains the macro thermodynamic behavior by imposing a probability distribution over the microscopic dynamical states. This paradigm would be useful as a formal tool to explore the relationships between different scales in economic

Introduction

and ecological systems. Also, the scaling up from microeconomic systems to macroeconomic systems that is achieved by using the equilibrium results at the micro scale to build models at the macro scale [17] can be applied to ecological systems.

In this dissertation, we do not use the formal structure that describes the evolution of thermodynamic systems from equilibrium, to non-equilibrium and self-organization. The dynamics of systems out of equilibrium, but close to equilibrium, is characterized by linear equations [2, 5, 12]. In these linear nonequilibrium thermodynamic systems (LNET) the production of irreversibilities is minimized. The dynamics of systems far away from equilibrium, studied in the framework of extended nonequilibrium thermodynamic systems (ENET), is characterized by nonlinear equations [12]. These far away from equilibrium systems are created and maintained by dissipative processes and evolve unpredictably with some of these possible final states being states that possess spatiotemporal organization [12]. This formal structure can be applied to economic and ecological systems to help characterize the evolution path for ecosystems and for economies [7].

The logical structure developed by Clausius-Kelvin-Carathéodory to describe equilibrium thermodynamics [19] can be used to establish a formal analogy with economics instead of the approach of Gibbs-Tisza-Callen used in this dissertation. A formal relationship between the two approaches to thermodynamics is already established [4, 19].

We use non-equilibrium thermodynamics to make a substantive analogy with the organism described by the Dynamic Energy Budget Theory in chapter 4. Although the main target of DEB theory is the organism, it also establishes links to the cellular and food web levels [13]. This means that DEB theory can be used together with non-equilibrium thermodynamics to extend the substantive integration to the cellular, organismic and ecological levels.

We formalize DEB theory into assumptions and propositions in chapter 5. This is a first step to use DEB theory to build up a non-optimizing theory in economics that would describe in more realistic terms the behavior of consumers and firms and at a more aggregate level the wealth of a country. This would also be an intermediate step for a substantive integration between thermodynamics and economics because DEB is compatible with non-equilibrium thermodynamics.

A description of both economic and ecological systems that is compatible with nonequilibrium thermodynamics is a crucial step to determine one of the most important issues in the interaction between ecologic and economic systems: the maximum scale of economic systems [3].

References

- [1] R. U. Ayres and U. E. Simonis. *Industrial Metabolism: Restructuring for Sustainable Development*. United Nations University Press, 1994.
- [2] H. Callen. *Thermodynamics and an Introduction to Thermostatistics*. John Wiley & Sons, New York, 1985.
- [3] R. Costanza, J. H. Cumberland, H. E. Daly, R. Goodland, and R. B. Norgaard. *An Introduction to Ecological Economics*. St. Lucie Press, Florida, 1997.
- [4] J. J. D. Domingos and T. Domingos. *Termodinâmica: Princípios e Conceitos Fun*damentais. IST Press, Lisboa, 2006.
- [5] J.J.D. Domingos. Transmissão do Calor por Condução e Convecção. Lisboa, 1964.
- [6] T. Domingos. *Theory and Modelling of Ecological Exploitation Systems*. PhD thesis, Instituto Superior Técnico, Lisbon, 2001.
- [7] C. Folke, S. Carpenter, B. Walker, M. Scheffer, T. Elmqvist, L. Gunderson, and Holling C. S. Regime shifts, resilience, and biodiversity in ecosystem management. *Annual Review of Ecology, Evolution and Systematics*, 35:557–581, 2004.
- [8] S. R. de Groot and P. Mazur. *Non-Equilibrium Thermodynamics*. North Holland, Amsterdam, 1962.
- [9] E. T. Jaynes. Information theory and statistical mechanics. *Physical Review*, 106:620–630, 1957.
- [10] G. Jehle. Advanced Microeconomic Theory. Prentice-Hall, Englewood Cliffs, NJ., 1991.
- [11] S. E. Jorgensen. *Integration of Ecosystem Theories: A Pattern*. Kluwer Academic Publishers, Dordrecht, 1997.
- [12] D. Kondepudi and I. Prigogine. Modern Thermodynamics: From Heat Engines to Dissipative Structures. Wiley, Chichester, 1998.
- [13] S. A. L. M. Kooijman. Dynamic Energy and Mass Budgets in Biological Systems. Cambridge University Press, Cambridge, 2000.

- [14] J. Martinez-Alier. Some issues in agrarian and ecological economics, In memory of Georgescu-Roegen. *Ecological Economics*, 22:225–238, 1997.
- [15] A. Mas-Collel and M. D. Whinston. *Microeconomic Theory*. Oxford University Press, New York, 1995.
- [16] M. Moran and H. Shapiro. Fundamentals of Engineering Thermodynamics. Wiley, New York, NY, 2000.
- [17] D. Romer. Advanced Macroeconomics. Mc Graw Hill, 1996.
- [18] E. Schrödinger. *What is Life? The Physical Aspects of a Living Cell*. Cambridge University Press, Cambridge, 1944.
- [19] L. Tisza. Generalized Thermodynamics. M.I.T. Press, Cambridge, MA, 1966.
- [20] M. Tribus. *Thermostatics and Thermodynamics*. D. Van Nostrand Company, Princeton, NJ, 1961.
- [21] J. Tschirhart. General equilibrium of an ecosystem. *Journal of Theoretical Biology*, 203:13–32, 2000.
- [22] H. Varian. Intermediate Microeconomics. A Modern Approach. Norton, New York, 1996.

Chapter 1

Chapter 2

Is neoclassical economics formally valid? An approach based on an analogy between Equilibrium Thermodynamics and Neoclassical Microeconomics

Abstract

The relation between Thermodynamics and Economics is a paramount issue in Ecological Economics. Two different levels can be distinguished when discussing it: formal and substantive. At the formal level, a mathematical framework is used to describe both thermodynamic and economic systems. At the substantive level, thermodynamic laws are applied to economic processes.

In Ecological Economics, there is a widespread claim that neoclassical economics has the same mathematical formulation as classical mechanics and is therefore fundamentally flawed because: 1) utility does not obey a conservation law as energy does; 2) an equilibrium theory cannot be used to study irreversible processes. Here, we show that neoclassical economics is based on a wrong formulation of classical mechanics, being in fact formally analogous to equilibrium thermodynamics. The similarity between both formalisms, namely that they are both cases of constrained optimisation, is easily perceived when thermodynamics is looked upon using the Tisza-Callen axiomatisation. In this paper, we take the formal analogy between equilibrium thermodynamics and economic systems far enough to answer the formal criticisms, proving that the formalism of neoclassical economics has irreversibility embedded in it.

However, the formal similarity between equilibrium thermodynamics and neoclassical

microeconomics does not mean that economic models are in accordance with mass, energy and entropy balance equations. In fact, neoclassical theory suffers from flaws in the substantive integration with thermodynamic laws as has already been fully demonstrated by valuable work done by ecological economists in this field.

Keywords: Thermodynamics, Entropy, Neoclassical Economics, Analogy and Irreversibility.

2.1 Introduction

The relation between Thermodynamics and Economics is a paramount issue in Ecological Economics. Two different levels can be distinguished when discussing it: formal and substantive.

At the formal level, a mathematical framework is used to describe both thermodynamic and economic systems. This allows for insights that were gained in one field of knowledge to be transposed to another. However, this has to be done with great care identifying whether the conditions that have to be met in the original field are also met in the field where the analogy is taken.

At the substantive level, thermodynamic laws are applied to economic processes. The integration between economics and thermodynamics at the substantive level is of crucial importance because economic processes obey thermodynamic laws and therefore a sound economic theory must be coherent with thermodynamics. This integration highlights the dependence between the economic system and the biophysical framework contributing to the analysis of the sustainability of economic systems.

This distinction between formal analogy and substantive integration is not a new issue, e.g., Martinez-Alier [33] says that 'the mathematical description of economic phenomena in the language of physics is different from applying the concepts of physics'. Also, Baumgärtner [9] discussing the different ways in which thermodynamics can be incorporated in economic analysis, considers the isomorphism of formal structure and the thermodynamic constraints on economic action among others, which are respectively, the formal analogy and the substantive integration discussed in this chapter.

The belief that neoclassical economics is based on a formal analogy to classical mechanics is common among ecological economists. For example, Amir [2] argues that 'most physical analogies in economic theory are borrowed from mechanics', Martinez-Alier [33] argues that 'economic science has used the mathematics of mechanics since the first neoclassical economists' and Costanza et al. [18] say that 'the market model has been formalized using the same mathematics as used by Newton for mechanical systems'. Outside ecological economics, this thesis has been most extensively argued by Mirowski [34], who considers that neoclassical economics is an attempt to emulate classical mechanics.

Based on this supposed analogy to classical mechanics, the main formal criticisms of neoclassical economics are: utility does not obey a conservation law as energy does; an equilibrium theory cannot be used to study irreversible processes.

Here, we argue that neoclassical economics is not formally identical to classical mechanics and that the correct identification of the formalism that underlies the construction of neoclassical economics is vital in the evaluation of its internal coherence. We show that economics is formally identical to thermodynamics because they are both problems of static constrained optimisation. The similarity between both formalisms has already been explored in the literature since the 40' [22, 30, 40, 41] and more recently [1, 12, 15, 16, 43, 44, 49, 50].

The formal analogies of Saslow [43], Berry et al. [12] and Amel'kin et al. [1] are driven from superficial similarities between the entities of economy and thermodynamics instead of being derived from fundamental principles. Therefore, these analogies are not helpful in answering the criticisms raised by ecological economists concerning the formal coherence of economic theory.

Candeal et al. [15, 16] prove that the mathematical representations of entropy and utility are analogous. Candeal et al. [15] and Cooper [17] investigate the mathematical foundations of the entropy representation where the entropy is built as an order preserving function that satisfies a continuity property. Candeal et al. [16] establish a formal relation between the entropy function and the utility function for the axioms that establish the existence of both ordering functions. Although, these authors do not develop their analysis any further, the formal analogy obtained at the function level is important and it lies behind the optimisation analogy pursued here.

The claim that neoclassical economics is formally identical to classical mechanics has also led many Ecological Economists to the substantive assertion that neoclassical economics is fundamentally flawed because it ignores thermodynamics. This argument lacks coherence because the existence of a formal analogy does not imply the existence of a substantive integration and vice-versa. This statement about the non-equivalence between the formal analogy and the substantive integration is easily argued, in this case, because the formal analogy uses the entities that are part of economic theory, i.e., utility, while the substantive integration uses the mass, energy and entropy flows in economic systems. Another argument that should help clarifying this issue is that different physical phenomena are described with different mathematical formalisms although they all have to obey thermodynamic laws. Whether neoclassical economics is formally identical to classical mechanics is not straightforwardly related to its substantive relation with thermodynamic laws.

For the same reason, although here we show that neoclassical economics is formally identical to thermodynamics, this does not imply that it is substantively compatible with thermodynamic laws. Whether neoclassical economics is in agreement with thermodynamic laws should be evaluated by looking at the 'entropic flow of energy and materials that runs through the economy' [33] instead of being based on the use of formal arguments. We would like to emphasize that ecological economics has already given many important contributions to this substantive integration between thermodynamics and economics with, among many others, the works of Georgescu-Roegen [25], Daly [19], Biancardi et al. [13], Ruth [38, 39], Stern [46], Ayres [4, 5, 6], Baumgärtner et al.[11], Ayres et al. [7], Tiezzi [47] and Frondel and Schmidt [24].

The roadmap of this chapter is as follows. In section 2.2, we motivate the reader for our formal analogy: (2.1) by explaining why there is the widespread idea that neoclassical economics is formally analogous to classical mechanics and (3.1) by reviewing some of the incorrect formal criticisms of neoclassical economics. In section 2.3, we present a unified formalism for thermodynamic and economic systems, based on the formalism of constrained optimisation. In section 2.4, some of the formalism's characteristics, namely its limits and scope and other related issues, are clarified. Section 3.12 concludes and argues that although there is a formal analogy between thermodynamics and neoclassical economics, these two fields are not substantively compatible.

2.2 Is the Formalism of Neoclassical Economics wrong?

It is generally claimed that neoclassical economics is based on classical mechanics because throughout the history of economics many economists used analogies from classical mechanics. Mirowski [34] gives some examples of the use of mechanical concepts and metaphors: (2.1) the lever rule by Nicolas Canard, (3.1) gravitation theory by Stanley Jevons, (3.5) force by Herman Gössen, (3.14) power by Frederick Soddy and (3.15) the Energy Minimum Principle by Francis Edgeworth.

This approach of establishing analogies between mechanics and economics was taken to its extreme by Irving Fisher who in 1892 established the most extensive relation between mechanics and economics (Table 2.2). According to Fisher [23], while economic equilibrium corresponds to maximum profit, mechanical equilibrium corresponds to minimum energy.

Mechanics	Economics
Particle	Individual
Space ^{*1}	Commodity*1
Force*1	MU^{*1}
Work = force \cdot space ^{*2}	Disutility = MU \cdot commodity ^{*2}
Energy = force \cdot space ^{*2}	Utility = MU \cdot commodity ^{*2}
Equilibrium: impelling and resisting	Equilibrium: MU and marginal
forces along each axis are equal	disutility along each axis are equal

Table 2.1: The analogy between mechanics and economics proposed by Fisher (1991, p. 85). *¹ (vector), *² (scalar) and MU(marginal Utility).

Given the history of economic analogies to mechanics, there is a widespread claim that neoclassical economics is fundamentally flawed because the assumptions on which classical mechanics is based do not apply to consumer theory. The most important aspects usually referred in the literature are: 1) utility does not obey a conservation law as energy does; 2) an equilibrium theory cannot be used to study irreversible processes. Some of the examples of this are described below.

Mirowski [34] considers that 'forgetting the conservation of energy while simultaneously appealing to the metaphor of energy ... is the Achilles heel of all neoclassical economics' because, according to Mirowski, although utility cannot be a conserved entity, the results obtained in consumer theory assume that it is.

Amir [2, 3] claims that the utility function is unlikely to be a conserved quantity, but that economic theory assumes that it is, which is supposedly patent in the use of the proportionality between marginal utilities and market prices in equilibrium.

Georgescu-Roegen [25] argues that Jevons and Walras, whose aim was to create an economic science similar to mechanics, built an economic theory that only describes reversible and qualityless motion. Georgescu-Roegen also argues that 'economics...is mechanistic in the same strong sense...[that] classical mechanics...because neither induces any qualitative change nor is affected by the qualitative change of the environment'.

Lozada [31] states that the entropy law is not reducible to mechanics by saying that 'the inconsistency between the logical structure of the entropy law and the logical structure of neoclassical economic analysis is that the former is evolutionary and the other is arithmomorphic and hence non-evolutionary'.

However it is important to emphasize that some authors do disagree with the supposed analogy between classical mechanics and neoclassical economics, e.g., Varian [51], in a review of Mirowski's book, argues that if the energy conservation principle implies that utility is not a coherent concept then this implies that utility is not energy', Marchionatti and Gambino [32] say that 'a critique, such as that of Mirowski, of the mechanical analogy in neoclassical economics, seems largely unhelpful and based on a misunderstanding' and Hands [27] argues that the standard Slutsky conditions, that are sufficient for the integrability of demand, do not seem to be sufficient to guarantee, as Mirowski argues, that prices form a conservative vector field.

We agree that if neoclassical economics were indeed formally identical to classical mechanics it would be internally incoherent. However, we argue that neoclassical economics is based on a wrong formulation of classical mechanics, being in fact formally identical to thermodynamics. Both neoclassical economics and thermodynamics are equilibrium theories and can be developed as formalisms of constrained optimisation as shown in the next section.

2.3 A Unified Formalism for Neoclassical Economics and Equilibrium Thermodynamics

In this section, a mathematical unified formalism based on the axiomatization of Tisza-Callen is first developed in the general case and then applied to thermodynamic and to economic systems.

2.3.1 General Formalism

The constrained optimisation problem describes the behaviour of the system that evolves in order to maximize some function y subject to a set of constraints¹. In equilibrium the values acquired by the variables, x_i , i = 1, ..., n, maximize the potential y, given the constraints.

This maximization is constrained because the variables in equilibrium have to obey a set of constraints, $g_z = 0$, z = 1, ..., m. These constraints are a function of the initial values

¹Minimization is an equivalent problem.

Is Neoclassical Economics Formally Valid?

of the variables, x_i^0 , and of some parameters k_j , $j=1,\ldots,l$:

$$\max_{x_1,...,x_n} \quad y(x_1,...,x_n) \quad s.t. \ g_z(x_1,...,x_n,x_1^0,...,x_n^0,k_1,...,k_l) = 0.$$
(2.1)

This problem is solved with the Method of Lagrangean Multipliers². The Lagrangean function, L, is defined as:

$$L(x_1, ..., x_n, k_1, ..., k_l, \lambda_1, ..., \lambda_m) =$$

$$y(x_1, ..., x_n) + \sum_{z=1}^m \lambda_z g_z(x_1, ..., x_n, x_1^0, ..., x_n^0, k_1, ..., k_l).$$
(2.2)

The values of the variables that maximize the objective function subject to the constraints are obtained solving the system of m + n equations:

$$\begin{aligned} \frac{\partial L}{\partial x_i} &= 0, \ i = 1, ..., n, \\ \frac{\partial L}{\partial \lambda_z} &= 0, \ \mathbf{z} = 1, ..., m \end{aligned}$$

which is equivalent to:

$$\frac{\partial y}{\partial x_i} = -\sum_{z=1}^n \lambda_z \frac{\partial g_z}{\partial x_i}, i = 1, ..., n$$

$$g_z = 0, \ z = 1, ..., m$$
(2.3)

If the function y and the constraints g_z are real valued and differentiable, if the number of constraints, m, is less than the number of variables, n, and if the gradient vectors of the constraint equations are linearly independent, the maximum exists. In this case, Lagrange's Method gives the first order equilibrium conditions [28], eq. (3.1), which define the state of the system. These equilibrium conditions give the optimal values of each x_i , x_i^* , as a function of the parameters k_j and of the initial conditions:

$$x_i^* = f\left(k_1, \dots, k_l, x_1^0, \dots, x_n^0\right).$$
(2.4)

With the set of equations (3.5) the optimal value of y, y^* , can also be written as:

$$y^* = f\left(k_1, \dots, k_l, x_1^0, \dots, x_n^0\right).$$
(2.5)

In the initial state, the system is characterized by the values x_i^0 ; which are called initial conditions. In the final state, the system is characterized by the optimal values assumed

 $^{^{2}}$ For a more detailed description of this method see, e.g., Jehle (1991).

by the variables, x_i^* . The evolution of the system from the initial to the final state is called a process. In a reversible process, the initial and final states have the same y^* . In an irreversible process, the final state has a higher value of y^* than the initial state. The amount of irreversibility (always positive) produced in an irreversible process can be measured by the increase in y. An impossible process is a process that will never occur because if it did the final state would have a lower value of y^* .

2.3.2 Thermodynamic Systems

The constrained optimisation problem in thermodynamics is the maximization of the entropy of a composite system with constant total internal energy, constant total volume and constant total mole numbers.

The entropy of a homogeneous thermodynamic system can be obtained from the fundamental equation [48, 14] ³:

$$S = S(U, V, N).$$

$$(2.6)$$

The variables are S, the entropy, U, the internal energy, V, the volume, and Nthe number of moles.

The problem of the maximization of entropy for a composite system⁴ comprising two simple systems (1 and 2) is formalized as⁵:

$$\max_{U_1, U_2, V_1, V_2, N_1, N_2} S = S_1 \left(U_1, V_1, N_1 \right) + S_2 \left(U_2, V_2, N_2 \right),$$

s. t.

$$U_{1} + U_{2} = U_{1}^{0} + U_{2}^{0}$$

$$V_{1} + V_{2} = V_{1}^{0} + V_{2}^{0}$$

$$N_{1} + N_{2} = N_{1}^{0} + N_{2}^{0}$$
(2.7)

where the constraints are that, over the two systems, the sums of the internal energies, the mole numbers and the volumes are kept constant. The initial conditions are the internal energy U_i^0 , the mole numbers N_i^0 and the volume V_i^0 for j=1,2.

³Where, for simplicity, we consider a system with a single chemical component.

⁴Caratheódory introduced the concept of composite systems because the entropy is axiomatically only defined for equilibrium systems.

⁵For simplicity, we consider only two subsystems.

Is Neoclassical Economics Formally Valid?

Defining temperature, T_j , pressure, P_j , and chemical potential, μ_j , for j = 1, 2, as:

$$\frac{\partial S_j}{\partial U_j} \equiv \frac{1}{T_j}$$

$$\frac{\partial S_j}{\partial V_j} \equiv \frac{P_j}{T_j}$$

$$\frac{\partial S_j}{\partial N_i} \equiv \frac{\mu_j}{T_i}$$
(2.8)

with $S_j = S_j(U_j, V_j, N_j)$, j = 1, 2, the equilibrium conditions, see eq. (3.1), for the thermodynamic system subject to the previous constraints are:

$$\frac{1}{T_1} = \frac{1}{T_2} \\ \frac{P_1}{T_1} = \frac{P_2}{T_2} \\ \frac{\mu_1}{T_1} = \frac{\mu_2}{T_2}$$

. That is, equilibrium is established when both systems have the same temperature, the same chemical potentials and the same pressure. Thus, an isolated composite system whose constraints are defined above increases its entropy towards the maximum value by turning into a system with a homogeneous temperature, homogeneous pressure and homogeneous chemical potential.

These equilibrium conditions are also explainable by reductio ad absurdum. Suppose that, in the equilibrium state, the temperatures of the two systems were not equal, e.g., $T_1 > T_2$. If energy would flow (in the form of heat) from system 1 to system 2, system 2 would increase its entropy and system 1 would decrease its entropy because the derivative of the entropy in relation with energy, eq. (2.8), is always positive given that the absolute temperature is always positive. The increase in entropy of system 2 would be higher than the decrease in entropy of the other system because its temperature is lower, again by eq. (2.8). Therefore, the global change in entropy would be positive and consequently the initial equilibrium state did not have the highest entropy possible, which means it was not an equilibrium state.

Each optimal variable, see eq. (3.5), can be written as:

$$U_{j}^{*} = f\left(V_{1}^{0}, V_{2}^{0}, U_{1}^{0}, U_{2}^{0}, N_{1}^{0}, N_{2}^{0}\right), j = 1, 2,$$

$$N_{j}^{*} = f\left(V_{1}^{0}, V_{2}^{0}, U_{1}^{0}, U_{2}^{0}, N_{1}^{0}, N_{2}^{0}\right), j = 1, 2$$

$$V_{j}^{*} = f\left(V_{1}^{0}, V_{2}^{0}, U_{1}^{0}, U_{2}^{0}, N_{1}^{0}, N_{2}^{0}\right), j = 1, 2,$$
(2.9)

These equations give the optimal value of the variables of each system as a function of the initial conditions. The maximum entropy, see eq. (3.14), as a function of the same arguments is:

$$S^* = f\left(V_1^0, V_2^0, U_1^0, U_2^0, N_1^0, N_2^0\right).$$
(2.10)

Reversible processes in a thermodynamic system are processes that occur in an isolated system at constant maximal entropy. Impossible processes are processes that do not occur because the isolated system would decrease its maximum entropy. Irreversible processes in a thermodynamic system are processes that occur in an isolated system at increasing maximal entropy. The amount of irreversibility produced is equal to the variation in entropy between the initial state and the equilibrium state.

2.3.3 Consumer System

The extremum principle, in consumer theory, is the maximization of utility of a consumer at constant market prices and constant endowment. When allowed to trade with the market, the consumer maximizes his utility, U.

The consumer's utility maximization problem is

$$\max_{x_1,\ldots,x_n} U = U\left(x_1,\ldots,x_n\right)$$

s.t.

$$p_1 x_1 + \dots + p_n x_n = p_1 x_1^0 + \dots + p_n x_n^0.$$
(2.11)

where the constraint is that the total cost of goods that will be bought (left hand side of the constraint) is equal to his initial endowment $p_1x_1^0 + \ldots + p_nx_n^0$. The variables are the different amounts of goods, x_i , the parameters are the market price p_i of good i, $i = 1, \ldots, n$. The initial conditions are the amounts of each good i in the initial state, x_i^0 .

Defining marginal utilities as:

$$\frac{\partial U}{\partial x_i} \equiv MU_i, \quad i = 1, \dots, n,$$
(2.12)

with $U = U(x_1, ..., x_n)$, the equilibrium conditions, see eq. (3.1), for the consumer are:

$$\frac{MU_1}{p_1} = \dots = \frac{MU_n}{p_n}.$$

A consumer system whose constraints are that the endowment and the prices remain constant increases his utility towards the maximum value by exchanging with the market until the ratio between marginal utility and price is the same for all products. Varian [52] proves this optimum condition by reductio ad absurdum using the following argument. Suppose that the consumer's utility is maximum but that the ratio between marginal utilities of product i and product j is two and the ratio of prices is one. Then, the consumer will exchange one unit of product j for one unit of product i obeying the endowment constraint (they are worth the same) and increase his utility. This means that the initial state was not at the maximum utility.

Each optimal value, see eq. (3.5), can be written as:

$$x_{j}^{*} = f(p_{1}, p_{2}, ..., p_{n}, x_{1}^{0}, ..., x_{n}^{0}) \quad j = 1, ..., n$$
 (2.13)

These equations give the optimal amount of product j as a function of the prices and initial amounts of each product. The maximum utility as a function of the same arguments is the equilibrium condition analogous to eq. (3.14),

$$U^* = f\left(p_1, \dots, p_n, x_1^0, \dots, x_n^0\right).$$
(2.14)

Reversible processes for the same consumer are processes in which he trades with the market but does not increase his maximum utility. Impossible processes are trades that do not occur because the consumer would decrease his maximum utility. Irreversible processes for the consumer are processes characterized by an increase in utility: if a consumer engages in a trade that increases his utility, he will not reverse the trade. The amount of irreversibility produced is equal to the amount of utility increase between the initial state and the equilibrium state.⁶

2.4 Discussing the Formalism

In this section we clarify some of the points of the formalism presented.

2.4.1 On the correct thermodynamic framework for substantive integration

We would like to emphasize that the substantive integration between thermodynamics and economic systems should not be based on the thermodynamic theory of isolated systems that we present in this chapter. Economic systems are open thermodynamic systems far from equilibrium and therefore a thermodynamic analysis of economic systems should be based on the thermodynamics of nonequilibrium open systems [29, 37, 35, 36, 26].

⁶This statement does not mean that we are considering that utility is a cardinal concept. This is further discussed in section 2.4.3.

2.4.2 On the concept of equilibrium

In the formalism of constrained optimization, equilibrium is the extremum point subject to the constraints considered. It is equilibrium because the system will not move from that point unless the constraints are changed. For example, for a certain amount of total internal energy, total volume and total number of moles, a thermodynamic system is in equilibrium if its entropy is at the maximum level (S^*). This entropy equilibrium value (S^*) can only be changed if the constraints are changed, e.g., the system receives a heat flow from the exterior hence increasing its total energy. The same is applicable to the consumer: for a certain total cost of goods, a consumer system is in equilibrium if its utility is at the maximum level (U^*). This utility level (U^*) can only be changed if the constraints are changed, e.g., the consumer receives an additional endowment.

The formal equilibrium considered for the consumer problem is not the thermodynamic equilibrium of the consumer. The thermodynamic equilibrium of the consumer would be a dead consumer.

2.4.3 On the ordinal concept of utility

Utility is usually considered in an ordinal framework, i.e, utility functions order the consumption bundles according to the preferences of the consumer but the absolute values of utility have no meaning. This means that if utility function $U(x_1, \ldots, x_n)$, represents the preferences of the consumer, then so does any other utility function obtained through a positive monotonic transformation $U^{\theta} = f(U(x_1, \ldots, x_n))$, (Jehle, 1991), i.e.,

$$\frac{\partial U^{\theta}}{\partial U} = \frac{\partial f(U)}{\partial U} > 0.$$
(2.15)

Our proposed formalism does not entail a cardinal concept of utility and is fully compatible with the ordinal framework because (2.1) the equilibrium conditions obtained and (3.1) the ranking of the amount of irreversibilities produced are both independent of the specific utility function chosen within the family of monotonic transformations. (2.1) is proved by Samuelson [42] and (3.1) is obtained below.

The amount of irreversibility produced between an initial state characterized by x_1, \ldots, x_n and a final equilibrium state x_1^*, \ldots, x_n^* can be measured using the utility function U or any other positive monotonic transformation $U^{\theta} = f(U)$:

$$\sigma = U^* (x_1^*, \dots, x_n^*) - U (x_1, \dots, x_n)$$

$$\sigma^{\theta} = U^{\theta *} (x_1^*, \dots, x_n^*) - U^{\theta} (x_1, \dots, x_n)$$
(2.16)

We prove that σ^{θ} is a positive monotonic transform of σ , using eq. (2.15),

$$\frac{\partial \sigma^{\theta}}{\partial \sigma} = \frac{\partial \left(U^{\theta*} \left(x_1^*, \dots, x_n^* \right) - U^{\theta} \left(x_1, \dots, x_n \right) \right)}{\partial \left(U^* \left(x_1^*, \dots, x_n^* \right) - U \left(x_1, \dots, x_n \right) \right)} = \frac{\partial U^{\theta*} \left(x_1^*, \dots, x_n^* \right)}{\partial U^* \left(x_1^*, \dots, x_n^* \right)} + \frac{\partial U^{\theta} \left(x_1, \dots, x_n \right)}{\partial U \left(x_1, \dots, x_n \right)} > 0.$$
(2.17)

Summarizing, the amount of irreversibilities produced between any initial state and the equilibrium state has an ordinal meaning, i.e., if the amount of irreversibilities produced going from state A to equilibrium is higher than the amount of irreversibilities produced going from state B to equilibrium, then the consumer will behave such as to move from state A to state B but not the reverse.

2.5 Conclusions

In the ecological economics literature, the validity of neoclassical economics has been widely discussed. This discussion has not been as fruitful as expected because the distinction between formal and substantive criticisms is blurred in many of the arguments presented.

In this chapter, we present and explore the validity of the formal criticisms of neoclassical economic theory. We concluded that formal criticisms are wrong because they are either based on mixing up the substantive and formal levels [20, 45] or they are based on the wrong assumption that the microeconomic formalism is analogous to the classical mechanics formalism [2, 3, 25, 31, 34].

We have shown that the argument that the neoclassical formalism is wrong because it was built from mechanics, is not valid; in fact, the neoclassical formalism is identical to the Tisza-Callen axiomatization of thermodynamics. For both thermodynamic and economic systems, we have identified the constrained optimisation problems along with the equilibrium conditions. We have proved that the formalism of neoclassical economics does not consider utility to be a conserved entity because it predicts the existence of irreversible processes characterized by increasing utility at constant market prices.

Smith and Foley [44] also present a formal analogy between neoclassical economics and thermodynamics based on constrained optimisation. These authors consider that the main difference between the two fields is that 'there is no counterpart in physics for the way neoclassical economists attach importance to transformations respecting initial endowments ... and there seems to be no counterpart in economics to the importance thermodynamics attaches to reversible transformations.' The first remark is not correct because initial endowments in consumer theory (initial amounts of goods) are similar to the initial extensive parameters characterizing an isolated composite system (volume, internal energy and number of moles). The second remark is not correct because in consumer theory many results are also obtained for reversible transformations, e.g., the compensating and the equivalent variations. In both fields, reversible transformations are just limiting cases of actual transformations but their study is important because they set limits on real transformations.

Finally, it is of fundamental importance that the fact that neoclassical economics is formally identical to thermodynamics does not mean that it is compatible with thermodynamic laws. Examples of flaws in the integration between economic theory and thermo-dynamic laws already identified are: economic theory considers a circular flow between households and firms without considering the one-way flow that begins with resources and ends with waste [25]; energy and capital are generally not substitutes, as assumed by production functions, but complements [21, 38, 46]; the Inada conditions are inconsistent with mass conservation [8]; and production theory does not fully possess thermodynamic irreversibility [10].

However these flaws do not result from the similarities between the formal structure of economics and mechanics. This misconception, which results from the use of the formal to argue the substantive, is present in many works like Daly [20], Sollner [45] and Geogescu-Roegen [25].

Daly [20] argues that standard economics explains circular flows because they are mechanistic in nature (reversible and qualityless) and that it does not explain the one-way flow of resources into waste because it is irreversible and mechanistic models cannot deal with irreversibility. We have proved that formally standard economics has irreversibility embedded in it; therefore, the fact that standard economics does not explain the physical directionality of economic flows cannot be accounted for by its formal structure.

Sollner [45] considers that the definition of value in economic theory is incompatible with the entropy law because it was taken from neoclassical mechanics. However, to exemplify this formal flaw, Sollner uses substantive arguments analogous to the ones provided by Georgescu-Roegen.

We argue that the flaws in the economic theory, referred by Sollner [45], Daly [20] and Georgescu-Roegen [25] do exist, but the causality relation used to explain these flaws is not correct. These flaws are due to a lack of substantive integration between thermodynamics and economics instead of being rooted in the supposed 'mechanistic' nature of neoclassical economics.

Concluding, we proved that the analogies to classical mechanics used by economists to build up neoclassical economic theory were wrongly formulated and that instead, neoclassical economic theory was built formally analogous to thermodynamics. Additionally, by clarifying the distinction between formal analogy and substantive integration we made clear that although neoclassical economics and thermodynamics are both formalisms of constrained optimisation they are not substantively compatible.

References

- S. Amel'kin, K. Martinás, and M. Tsirlin. Optimal control for irreversible processes in thermodynamics and microeconomics. *Automation and Remote Control*, 63:519– 539, 2002.
- [2] S. Amir. Welfare maximization in economic theory: Another viewpoint. *Structural Change and Economic Dynamics*, 6:359–376, 1995.
- [3] S. Amir. The role of thermodynamics for ecological economics. *Ecological Economics*, 27:213–214, 1998.
- [4] R. U. Ayres. Eco-thermodynamics: economics and the second law. *Ecological Economics*, 26:189–209, 1998.
- [5] R. U. Ayres. The second law, the fourth law, recycling and limits to growth. *Ecological Economics*, 29:473–483, 1999.
- [6] R. U. Ayres. The minimum complexity of endogeneous growth models: the role of physical resource flows. *Energy*, 26:817–838, 2001.
- [7] R. U. Ayres, L. Ayres, and B. Warr. Exergy, Power and Work in the US economy, 1900-1998. *Energy*, 28:219–273, 2003.
- [8] S. Baumgärtner. The Inada Conditions for material resource inputs reconsidered. *Environmental & Resource Economics*, 29:307–322, 2004.
- [9] S. Baumgärtner. Thermodynamic models. In P. Proops, J. & Safonov, editor, *Modeling in Ecological Economics.*, pages 102–129. Edward Elgar, Cheltenham, 2004.

- [10] S. Baumgärtner. Temporal and thermodynamic irreversibility in production theory. *Economic Theory*, 26:725–728, 2005.
- [11] S. Baumgärtner, H. Dyckhoff, M. Faber, J. Proops, and J. Schiller. The concept of joint production and ecological economics. *Ecological Economics*, 36:365–372, 2001.
- [12] R. Berry, V. Kasakov, S. Sieniutycz, Z. Szwast, and A. Tsirlin. *Thermodynamic Optimization of Finite-Time Processes*. Wiley, West Sussex, UK, 2000.
- [13] C. Biancardi, E. Tiezzi, and U. Ulgiati. Complete recycling in the frameworks of physics, biology and ecological economics. *Ecological Economics*, 8:1–5, 1993.
- [14] H. Callen. *Thermodynamics and an Introduction to Thermostatistics*. John Wiley & Sons., New York, NY, 2 edition, 1985.
- [15] J. Candeal, J. Miguel, E. Indurain, and G. Mehta. On a theorem of Cooper. *Journal of Mathematical Analysis and Applications*, 258:701–710, 2001.
- [16] J. Candeal, J. Miguel, E. Indurain, and G. Mehta. Utility and entropy. *Economic Theory*, 17:233–238, 2001.
- [17] J. Cooper. The foundations of thermodynamics. *Journal of Mathematical Analysis and Applications*, 17:172–193, 1967.
- [18] R. Costanza, J. Cumberland, H. Daly, R. Goodland, and R. Norgaard. An Introduction to Ecological Economics. St. Lucie Press: ISEE, Boca Raton, FL., 1997.
- [19] H. Daly. Steady-State Economics. Island Press, Washington, DC, 1991.
- [20] H. Daly. On Nicholas Georgescu-Roegen's contributions to economics: An obituary essay. *Ecological Economics*, 13:149–154, 1995.
- [21] H. Daly. Georgescu-Roegen versus Solow/Stiglitz. *Ecological Economics*, 22:261–266, 1997.
- [22] H. Davis. The Theory of Econometrics. Principia Press, Bloomington, IN., 1941.
- [23] I. Fisher. Mathematical Investigations in the Theory of Value and Prices. Reprint, Augustus M. Kelley, Fairfield, NJ., 1991.
- [24] M. Frondel and C. Schmidt. Facing the truth about separability: nothing works without energy. *Ecological Economics*, 51:217–223, 2004.
- [25] N. Georgescu-Roegen. The Entropy Law and the Economic Process. Harvard University Press, Cambridge, MA., 1971.
- [26] S. Groot and P. Mazur. *Non-Equilibrrium Thermodynamics*. North Holland, Amsterdam, 1962.
- [27] W. Hands. More light on integrability, symmetry and utility as potential energy in Mirowski's critical history. *History of Political Economy*, 25:118–130, 1993.
- [28] G. Jehle. Advanced Microeconomic Theory. Prentice-Hall, Englewood Cliffs, NJ., 1991.
- [29] D. Kondepudi and I. Prigogine. *Modern Thermodynamics: From Heat Engines to Dissipative Structures*. Wiley, Chichester, 1998.
- [30] J. Lisman. Econometrics and Thermodynamics: A Remark on Davis' Theory of Budgets. *Econometrica*, 17:59–62, 1949.
- [31] G. Lozada. Georgescu-Roegen's defense of classical thermodynamics revisited. *Ecological Economics*, 14:31–44, 1995.
- [32] R. Marchionatti and E. Gambino. Pareto and political economy as a science: Methodological revolution and analytical advances in economic theory in the 1890's. *Journal of Political Economy*, 105:1322–1348, 1997.
- [33] J. Martinez-Alier. Some issues in agrarian and ecological economics, in memory of Georgescu-Roegen. *Ecological Economics*, 22:225–238, 1997.
- [34] P. Mirowski. Economics as Social Physics, Physics as Nature's Economics. Cambridge University Press,, New York, 1999.
- [35] L. Onsager. Reciprocal Relations in Irreversible Processes, I. *Physical Review*, 37:405–426, 1931.
- [36] L. Onsager. Reciprocal Relations in Irreversible Processes, II. *Physical Review*, 38:2265–2279, 1931.

- [37] I. Prigogine. *Thermodynamics of irreversible processes*. Wiley, New York, NY, 1961.
- [38] M. Ruth. Integrating Economy, Ecology and Thermodynamics. Kluwer Academic, Dordech, 1993.
- [39] M. Ruth. Thermodynamic constraints on optimal depletion of copper and aluminium in the united states: a dynamic model of substitution and technical change. *Ecological Economics*, 15:197–213, 1995.
- [40] P. A. Samuelson. Essays in Economics and Econometrics: A Volume in Honor of Harold Hotteling, chapter Structure of a Minimum Equilibrium System. University of North Carolina Press, Chapel Hill, NC., 1960.
- [41] P. A. Samuelson. An extension of the Le Chatelier principle. *Econometrica*, 28:368–379, 1960.
- [42] P. A. Samuelson. Foundations of Economic Analysis. Harvard University Press, Cambridge, MA., 1983.
- [43] W. Saslow. An economic analogy to thermodynamics. *Americal Journal of Physics*, 67:1239–1247, 1999.
- [44] E. Smith and D. Foley. Classical thermodynamics and economic general equilibrium theory. *New School University, Center for Economic Policy Analysis Working Papers*, 2004.
- [45] F. Sollner. A reexamination of the role of thermodynamics for environmental economics. *Ecological Economics*, 22:175–201, 1997.
- [46] D. Stern. Limits to substitution and irreversibility in production and consumption: A neoclassical interpretation of ecological economics. *Ecological Economics*, 21:197– 215, 1997.
- [47] E. Tiezzi. *The Essence of Time*. WIT Press, Southampton, 2002.
- [48] L. Tisza. Generalized Thermodynamics. M.I.T. Press, Cambridge, MA., 1966.
- [49] M. Tsirlin and S. Amel'kin. Dissipation and conditions of equilibrium for an open microeconomic system. *Open Systems and Information Dynamics*, 8:157–168, 2001.

- [50] M. Tsirlin, V. Kasakov, and N. Kolinko. Irreversibility and limiting possibilities of macrocontrolled systems: II. Microeconomics. *Open Systems and Information Dynamics*, 8:329–347, 2001.
- [51] H. Varian. Review of more heat than light: Economics as social physics: Physics as nature's economics, by Philip Mirowski. *Journal of Economic Literature*, XXIX:595–596, 1991.
- [52] H. Varian. *Intermediate Microeconomics. A Modern Approach*. Norton, New York, 1996.

Chapter 2

28

Chapter 3

Equilibrium Econophysics: A unified formalism for neoclassical economics and equilibrium thermodynamics

Abstract

We develop a unified conceptual and mathematical structure for Equilibrium Econophysics, i.e., the use of concepts and tools of equilibrium thermodynamics in neoclassical microeconomics and vice-versa. Within this conceptual structure the results obtained in microeconomic theory are: (1) the definition of irreversibility in economic behavior, (2) the clarification that the Engel curve and the offer curve are not descriptions of real processes dictated by the maximization of utility at constant endowment, (3) the derivation of a relation between elasticities proving that economic elasticities are not all independent, (4) the proof that Giffen goods do not exist in a stable equilibrium, (5) the derivation that 'economic integrability' is equivalent to the Generalized Le Chatelier Principle and (6) the definition of a first order phase transition, i.e., a transition between separate points in the utility function. In thermodynamics the results obtained are: (1) a relation between the non-dimensional isothermal and adiabatic compressibilities and the increase or decrease in the thermodynamic potentials, (2) the distinction between mathematical integrability and optimization behavior and (3) the generalization of the Clapeyron Equation.

Keywords: Econophysics, Entropy, Utility, Optimization and Unification of Knowledge.

3.1 Introduction

A general definition of econophysics is the use of concepts and tools in Economics that were developed in Physics. The state of the art in this field is the use of statistical physics to get new insights into the behavior of financial markets [8, 10, 11, 15, 19, 21, 29, 32, 36, 39]. Here, we focus instead on the similarity between Neoclassical Microeconomics and Equilibrium Thermodynamics. Some similarities between both formalisms have been explored in the literature since the 40's and 60's [9, 16, 24, 23, 25, 27, 26] and more recently [1, 2, 5, 6, 28, 30, 31, 34, 35]. In this chapter, we develop a unified formalism, for both fields, using the Tisza-Callen axiomatization of thermodynamics [4, 33], and take it further, making contributions both to Economics and to Thermodynamics. Also, by formulating both theories with the same mathematical formalism, neoclassical economics becomes accessible to physicists and thermodynamics becomes accessible to economists. This makes this chapter also an interesting pedagogical tool, e.g., in B.Sc. courses in econophysics like the one proposed by Grech [12].

The road-map of this chapter is as follows. All sections start by the description of the general formalism followed by its application to the equilibrium thermodynamic system and then to the consumer. In section 3.2, the fundamental equation, the constraints and the extremum principle that describe the behavior of these systems are explained and the equilibrium conditions are obtained. An alternative formulation to describe the systems' behavior, the duality equation, is presented in section 3.3. The reversible, irreversible and impossible processes undertaken by these systems are clarified in sections 3.4 and 3.5. In section 3.6, adequate descriptions of systems with a special type of constraints are obtained with the Legendre transforms. In section 3.7, the properties that characterize these systems are identified and in section 3.8 the relation between them is established. The stability conditions, i.e., the constraints imposed on the values of the systems' properties by the maximization principle, are presented in section 3.9 and the process that occurs as a result of instability, i.e., the first order phase transition, is explained in section 3.11. The issue of whether the system is described by the constrained optimization formalism developed in the chapter is addressed in section 3.10. Section 3.12 concludes.

The microeconomic formalism of the consumer can contemplate two cases: (1) a consumer who, in each of many periods, chooses between alternative bundles constrained by the same budget and (2) a consumer who, over a short period of time faces a series of opportunities to buy or sell goods at constant endowment; in the first case the bundle is consumed within that period and in the second case the consumption is postponed [20].

In this chapter, we will focus on the second case because: (1) it is more adequate for financial markets, currently the major focus of econophysics and (2) the similarities with the equilibrium thermodynamics systems are more straightforward.

3.2 Fundamental Equation and the Equilibrium State

A system that evolves in order to maximize some function y subject to a set of constraints ¹ is described by a constrained optimization problem: in equilibrium the values acquired by the state variables, x_i , i = 1, ..., n, maximize the function y, given the constraints. The function $y(x_1, ..., x_n)$ is the **fundamental equation** of the system. The state variables are **extensive variables**, i.e., additive variables.

This maximization is constrained because the variables in equilibrium have to obey a set of constraints, which are a function of the initial values of the variables, x_i^0 , and of some parameters k_j , j=1,...,l:

$$\max_{x_1,...,x_n} y(x_1,...,x_n) \quad s.t.$$

$$g^z(x_1,...,x_n,x_1^0,...,x_n^0,k_1,...,k_l) = 0,$$

$$z = 1,...,m.$$
(3.1)

This problem is solved with the Method of Lagrangean Multipliers². The Lagrangean function, \mathcal{L} , is defined as:

$$\mathcal{L}(x_{1},...,x_{n},x_{1}^{0},...,x_{n}^{0},k_{1},...,k_{l},\lambda_{1},...,\lambda_{m}) = y(x_{1},...,x_{n}) + \sum_{z=1}^{m} \lambda_{z}g^{z}(x_{1},...,x_{n},x_{1}^{0},...,x_{n}^{0},k_{1},...,k_{l}).$$
(3.2)

where λ_z are the Lagrangean multipliers.

The values of the variables that maximize the objective function subject to the constraints are obtained solving the system of m + n equations:

$$\frac{\partial \mathcal{L}}{\partial x_i} = 0, \quad i = 1, \dots, n, \tag{3.3}$$

$$\frac{\partial \mathcal{L}}{\partial \lambda_z} = 0, \quad z = 1, \dots, m,$$
(3.4)

¹Minimization is an equivalent problem.

²For a more detailed description of this method see, e.g., Jehle (1991).

which is equivalent to

$$\pi_i = -\sum_{z=1}^n \lambda_z \frac{\partial g^z}{\partial x_i}, \quad i = 1, \dots, n,$$
(3.5)

$$g^z = 0, \quad z = 1, \dots, m,$$
 (3.6)

where $\pi_i \equiv \frac{\partial y}{\partial x_i}$, i = 1, ..., n. The variable π_i is an **intensive variable**, i.e., a non-additive variable. If the function y and the constraints g^z are real valued and differentiable, if the number of constraints, m, is less than the number of variables, n, and if the gradient vectors of the constraint equations are linearly independent, the maximum exists. In this case, Lagrange's Method gives the first order **equilibrium conditions** [13], Eq. 3.5 and Eq. 3.6, which define the state of the system. The **differential form** of the fundamental equation is

$$dy = \sum_{i=1}^{n} \pi_i dx_i. \tag{3.7}$$

The constrained optimization problem in thermodynamics is the maximization of entropy, $S(U^1, U^2, V^1, V^2, N_1^1, N_1^2, \ldots, N_r^1, N_r^2)$, of a composite system, with subsystems 1 and 2 (see Appendix 3.13.1 for further details), with constant total internal energy, U, constant total volume, V, and constant total mole numbers for each chemical component, N_i , $i = 1, \ldots, r$, where N_i is the number of moles of the chemical compound i^3 . Thermodynamic intensive variables are:

$$1/T^{j} \equiv \partial S/\partial U^{j}, \qquad (3.8)$$
$$P^{j}/T^{j} \equiv \partial S/\partial V^{j},$$
$$-\mu_{i}^{j}/T^{j} \equiv \partial S/\partial N_{i}^{j},$$

with j = 1, 2 and i = 1, ..., r, where T^j is the temperature, P^j is the pressure and μ_i^j is the chemical potential of the *i* component of subsystem *j*. Temperature is defined positive by Nernst's postulate [4]. For the thermodynamic system, the equilibrium conditions (Eq. 3.5) are: $1/T^1 = 1/T^2 \equiv 1/T$, $P^1/T^1 = P^2/T^2 \equiv P/T$ and $\mu_i^1/T^1 = \mu_i^2/T^2 \equiv \mu_i/T$ with i = 1, ..., r. Thus, an isolated composite system with the above constraints increases its entropy towards the maximum value by turning into a system with a homogeneous temperature, homogeneous pressure and homogeneous chemical potentials for all compounds. The differential form of the fundamental equation is

$$dS = \sum_{j=1}^{2} \frac{1}{T^{j}} dU^{j} + \sum_{j=1}^{2} \frac{P^{j}}{T^{j}} dV^{j} - \sum_{j=1}^{2} \sum_{i=1}^{r} \frac{\mu_{i}^{j}}{T^{j}} dN_{i}^{j}.$$
(3.9)

³It is considered that no chemical reactions occur.

The extremum principle, in consumer theory, is the maximization of utility,

 $U(x_1, \ldots, x_n)$, of a consumer at constant market prices, p_i , and constant endowment given by $w = p_1 x_1^0 + \ldots + p_n x_n^0$ where x_i , $i = 1, \ldots, n$, are the available amounts of each good. The utility function, U, is the consumer's fundamental equation because it describes its behavior: when allowed to trade with the market, the consumer behaves in order to maximize his utility. The consumer intensive variables are the marginal utility of good i,

$$MU_i \equiv \partial U / \partial x_i. \tag{3.10}$$

Marginal utilities are defined positive by the assumption of monotonicity usually made on the utility function (Axiom 5, [13]): if a consumption bundle involves more of at least one good and no less of any other good, then it provides a higher utility than the original consumption bundle. This is equivalent to assuming that there are no 'bads', only goods ⁴. The equilibrium conditions (Eq. 3.5) for the consumer are: $MU_1/p_1 =$ $\dots = MU_n/p_n$. Thus, a consumer whose constraints are that the endowment and the prices remain constant increases his utility towards the maximum value by exchanging with the market until the ratio of marginal utility to price is the same for all products. The differential form of the fundamental equation is

$$dU = MU_1 dx_1 + \ldots + MU_n dx_n. \tag{3.11}$$

3.3 Duality Formulation

A duality formulation of the fundamental equation is the inverse function $x_1 = x_1 (y, x_2, ..., x_n)$. Its intensive variables are given by:

$$\frac{\partial x_1}{\partial y} = \frac{1}{\pi_1}$$
$$\frac{\partial x_1}{\partial x_j} = -\frac{\pi_j}{\pi_1}, \quad j = 2, \dots, n.$$
(3.12)

The duality formulation also has an extremum principle if y is a monotonically increasing function of x_1 , i.e., $\pi_1 > 0$:

$$\min_{\substack{y, x_2, \dots, x_n \\ y, x_2, \dots, x_n, y^0, x_2^0, \dots, x_n^0, k_1, \dots, k_l \\} s.t.$$

$$g^z \left(y, x_2, \dots, x_n, y^0, x_2^0, \dots, x_n^0, k_1, \dots, k_l \right) = 0,$$

$$z = 1, \dots, m.$$
(3.13)

⁴This is a simplification because 'bads' do exist, e.g., air pollution.

The proof by reduction to absurd is a generalization of theorem 4.3.2 given by Jehle [13]. If $y^* = \max y$ but $x_1^* \neq \min x_1$ then there exists at least $y^{*,a}(x_1^{*,a}, x_2, \ldots, x_n) = y^*(x_1^*, x_2, \ldots, x_n)$ such that $x_1^{*,a} < x_1^*$. Since x_1 is increasing in y then $y^* > y^{*,a}$ which contradicts the initial assumption.

In the thermodynamic system, the duality is between entropy and energy. Entropy is a concave function and an increasing function of the internal energy, i.e., $\partial^2 S/\partial U^2 < 0$ and $\partial S/\partial U > 0$ which means that the internal energy is a convex function. Thus, the fundamental equation can also be written in the energy representation, $U(S^1, S^2, V^1, V^2, N_1^1, N_1^2, \dots, N_r^1, N_r^2)$ (see Appendix 3.13.1 for further details); in this formulation the relevant extremum principle is the *Energy Minimum Principle*. In the energy representation, the intensive variables are straightforwardly the temperature, the (minus) pressure and the chemical potentials.

An intuitive explanation for the Energy Minimum Principle is the following: if the thermodynamic system is in equilibrium then its entropy is maximum, but if the energy is not in its minimum value then this quantity 'in excess' can be withdrawn from the system keeping the entropy constant. This quantity can then be returned to the system increasing the entropy because temperatures are positive. Thus, the initial state was not an equilibrium state because it was not at the maximum entropy [4].

In the consumer system, the duality is between utility and product demand. Utility is a concave function and an increasing function of the amount of product, i.e., $\partial^2 U/\partial x_i^2 < 0$ and $\partial U/\partial x_i > 0$ which means that the product demand function is a convex function. Thus, the fundamental equation can also be written in the product demand representation, $x_i(U, x_1, \ldots, x_{j \neq i}, \ldots, x_n)$. This equation represents the demand for product *i* as a function of utility and the amounts of the other products. In the product demand representation the extremum principle is the *Product Minimum Principle*.

An intuitive explanation for the Product Minimum Principle is the following: if the consumer system is in equilibrium then its utility is maximum, but if the consumption of product i is not in its minimum value then this quantity 'in excess' can be withdrawn from the system keeping the utility constant. This quantity could be later returned to the system increasing the utility because marginal utilities are positive. Thus, the initial state was not an equilibrium state because the consumer was not at the maximum utility.

The economic variables x_1, \ldots, x_n , in contrast to the thermodynamic variables, have a similar meaning, in the sense that, using prices, they can all be added. Therefore, the Product Minimum Principle can be applied for each product, implying that the feasible bundle is the one with the lowest amount of each product and the lowest cost. The explanation is that if the consumer did not choose the bundle with the lowest cost, he could trade it for a bundle with the same utility but a lower cost and use the remaining budget to buy goods and increase his utility. In consumer theory, this is known as (1) the duality between maximum utility and minimum expenditure, $E(x_1, \ldots, x_n) = \sum_{i=1}^n p_i x_i$ or (2) the equivalence between maximum utility and the cheapest bundle that provides that utility (axiom 5, [13]). While the maximum utility problem computes the maximum level of utility given a certain endowment, w, the minimum expenditure problem computes the maximum level of utility given a certain endowment, w, the minimum expenditure problem computes the minimum endowment required to reach a certain level of utility [18].

3.4 Reversible, Irreversible and Impossible Processes

The evolution of the system from the initial to the final state as described by Eq. 3.1, is triggered by a change in parameters; this is called a one step process. In a **reversible process**, the initial and final states have equal y. In an **irreversible process**, the final state has a higher value of y than the initial state. The amount of irreversibility (always positive) produced in an irreversible process can be measured by the increase in y. An **impossible process** is a process that will never occur because if it did the final state would have a lower value of y.

Reversible processes in a thermodynamic system are processes that occur in an isolated system⁵ at constant maximal entropy. An impossible process is a process that will never occur because if it did the maximum entropy of an isolated system would decrease. Irreversible processes in a thermodynamic system are processes that occur in an isolated system at increasing maximal entropy. The amount of irreversibility produced is equal to the variation in entropy between the initial state and the equilibrium state.

Reversible processes for the consumer are processes in which he trades at constant endowment with the market but does not increase his maximum utility. Impossible processes are trades that do not occur because the consumer with a constant endowment would decrease his maximum utility. Irreversible processes for the consumer are processes characterized by an increase in utility: if a consumer with constant endowment engages in a trade that increases his utility, he will not reverse the trade. The amount of irreversibility produced is equal to the amount of utility increase between the initial state and the equilibrium state.

⁵An isolated thermodynamic system is a system with the set of constraints defined in Sec. 3.2.

3.5 Many-step Processes

The evaluation of the change in the equilibrium value of the potential function, y^* , in response to the change in its parameters is known as *comparative statics* analysis.

Given a change in parameters dk_i, dk_j, \ldots , the optimal values x_i^* , as a function of the parameters and of the initial conditions,

$$x_i^* = x_i^*(k_1, \dots, k_l, x_1^0, \dots, x_n^0),$$
(3.14)

are obtained from the equilibrium conditions, Eq. 3.5 and 3.6, and the optimal value function y^* ,

$$y^* = y^*(k_1, \dots, k_l, x_1^0, \dots, x_n^0),$$
(3.15)

is obtained from the set of Eq. 3.14. Thus, Eq. 3.14 and 3.15 can be used to study a **many-step process** composed of many maximization steps where: (1) each maximization step is given by Eq. 3.1 and triggered by a change in one or more parameters and (2) the optimal value of x_i obtained at the end of each step is the initial value for the following step.

Not all trajectories described by Eq. 3.14 and 3.15 are processes because not all parameters are independent of each other. Suppose that $k_1(k_2, k_3)$; in this case, it is not possible to impose independently dk_1 , dk_2 and dk_3 .

Quasi-static Processes are processes characterized by infinitesimally close equilibrium points y^* ; this requires infinitesimal changes in the parameters. Quasi-static processes are reversible, irreversible or impossible according to the criteria defined.

The effects of changes dk_i in the value function y^* are addressed by the **envelope the**orem summarized next (for more details consult [18]). Assuming that y^* is differentiable then at the equilibrium x_1^*, \ldots, x_n^* :

$$\frac{\partial y^*}{\partial k_i} = \frac{\partial y}{\partial k_i} - \sum_{z=1}^m \lambda_z \frac{\partial g^z}{\partial k_i}.$$
(3.16)

For the thermodynamic system, Eq. 3.14 is, e.g., $U_1^*(U, V, N_1, \ldots, N_r)$ while Eq. 3.15 is $S^*(U, V, N_1, \ldots, N_r)$.

The application of the envelope theorem to the maximization of entropy gives:

$$\frac{\partial S^*}{\partial U} = -\lambda_1 = \frac{1}{T},$$

$$\frac{\partial S^*}{\partial V} = -\lambda_2 = \frac{P}{T},$$

$$\frac{\partial S^*}{\partial N_i} = -\lambda_j = -\frac{\mu_i}{T}, \quad j = 3, \dots, r+2.$$
(3.17)

The application of the envelope theorem to the minimization of energy gives:

$$\frac{\partial U^*}{\partial S} = -\lambda_1 = T,$$

$$\frac{\partial U^*}{\partial V} = -\lambda_2 = -P,$$

$$\frac{\partial U^*}{\partial N_i} = -\lambda_j = \mu_i, \quad j = 3, \dots, r+2.$$
(3.18)

For the consumer, Eq. 3.14 is the Walrasian demand function given by $x_i^*(w, p_1, \ldots, p_n)$ while Eq. 3.15 is the *indirect utility function*, $U^*(w, p_1, \ldots, p_n)$. For a given a set of prices and endowment, the Walrasian demand function specifies the amount of product x_i that maximizes utility while the indirect utility function specifies the maximum utility achievable.

The *Engel curve* is the Walrasian demand function with constant prices [18]. The Engel curve does not depict a process because endowment is a function of prices, $w(p_1, \ldots, p_n)$, i.e., it is not possible to change endowment while keeping all prices constant. It would only depict a process if at the end of each process the consumer either (1) had completely consumed all its goods and received a new amount of money equal to w + dw or (2) exchanged all its goods for money and then received the extra endowment dw. These are not the conditions assumed for a many-step process.

The offer curve depicts the relation between price p_i and the optimal quantity of good i with all other prices and endowment constant, [18]. Like the Engel curve, the offer curve does not depict a process because $p_i(w, p_1, \ldots, p_{j,\forall j \neq i}, \ldots, p_n)$, i.e., it is not possible to change the price of good i while keeping all other prices and the endowment constant.

For the minimum expenditure representation, Eq. 3.14 is the *Hicksian demand func*tion given by $x_i^{h,*}(U, p_1, \ldots, p_n)$ while Eq. 3.15 is the expenditure function given by $w^*(U, p_1, \ldots, p_n)$. Maintaining a specified level of utility given a set of prices requires the amounts of products x_i , $i = 1, \ldots, n$, given by the Hicksian demand function and the endowment specified by the expenditure function.

The application of the envelope theorem for the maximization of utility gives

$$\frac{\partial U^*}{\partial p_i} = -\lambda x_i^*,$$

$$\frac{\partial U^*}{\partial w} = \lambda.$$
(3.19)

Combining these results yields Roy's Identity [18]:

$$x_i^* = -\frac{\frac{\partial U^*}{\partial p_i}}{\frac{\partial U^*}{\partial w}}.$$
(3.20)

Roy's identity gives a method to obtain the Walrasian demand functions knowing the indirect utility functions. The intuition is the following: the higher the amount of x_i that a consumer needs to maximize his utility, the higher is the dependence of the maximum utility on p_i relative to its dependence on endowment. The application of the envelope theorem for the minimization of expenditure yields *Shepard's Lemma* [18]:

$$x_i^{h,*} = \frac{\partial w^*}{\partial p_i}.$$
(3.21)

Shephard's lemma gives a relationship between the Hicksian demand and the expenditure function. The intuition is the following: the higher the amount of x_i that a consumer needs to maintain his utility, the higher must be the expenditure increase with an increase in p_i to maintain the same utility.

3.6 Legendre Transforms

The **Legendre transform** is a mathematical technique by which one or more of the partial derivatives π_i replace one or more of the extensive variables x_i as independent variables. If the substitution of x_i by π_i is made for one variable, the Legendre transform is

$$\varphi_i(x_1,\ldots,\pi_i,\ldots,x_n) = y(x_1,\ldots,x_n) - \pi_i x_i, \qquad (3.22)$$

if it is is made for all variables, we have the complete Legendre transform

$$\varphi(\pi_1, \dots, \pi_i, \dots, \pi_n) = y(x_1, \dots, x_n) - \sum_{i=1}^n \pi_i x_i.$$
 (3.23)

The meaning of the Legendre transform is easily grasped for two simplified cases: (1) the fundamental equation is a first order homogeneous function, i.e., the *Euler Equation*, $\pi_1 x_1 + \ldots + \pi_n x_n = y$, applies (see Appendix 3.13.4 for further details), implying that:

$$\varphi_i\left(x_1,\ldots,\pi_i,\ldots,x_n\right) = \sum_{j \neq i} \pi_j x_j, \qquad (3.24)$$

(2) π_i is constant implying that the differential of the Legendre transform simplifies to: $d\varphi_i(x_1, \ldots, \pi_i, \ldots, x_n) = dy - \pi_i dx_i = \sum_{j \neq i} \pi_j dx_j.$

Suppose that an extremum principle applies to the initial function and to its Legendre transform. The Langrangean of the Legendre transform is

$$L(x_1,\ldots,x_n,\lambda_1,\ldots,\lambda_n) =$$
(3.25)

$$\underbrace{\begin{array}{c} \varphi(x_1,\dots,\pi_w,\dots,x_n) \\ y(x_1,\dots,x_w,\dots,x_n) - \pi_w x_w \\ + \sum_{z=1,m} \lambda'_z g'^z(x_1,\dots,\pi_w,\dots,x_n), \end{array}}_{\substack{\varphi(x_1,\dots,\pi_w,\dots,x_n)}}$$

and the Lagrangean of function y is given by Eq. 3.2. The first order conditions to the optimization problems are respectively:

$$-\sum_{i=1,n}\lambda'_{z}\frac{\partial g'^{z}(x_{1},\ldots,x_{n})}{\partial x_{i\neq w}} = \frac{\partial\varphi(x_{1},\ldots,\pi_{w},\ldots,x_{n})}{\partial x_{i\neq w}} = \pi'_{i},$$
(3.26)

and Eq. 3.5. If π_w is constant, then $\pi_i = \pi'_i$ for all $i \neq w$ and Eq. 3.26 is equal to Eq. 3.5. Thus, whenever an extremum principle is imposed on the original function and π_w remains constant, an extremum principle is also imposed on the Legendre transform. Theorem 2.3.3 [13] determines the sufficient conditions for the constrained optimal point to be either a maximum or a minimum (see a more detailed discussion in sec. 3.9). According to this theorem the extremum is a constrained maximum if the principal minors of the Hessian matrix of the Lagrangean evaluated at the extremum alternate in sign beginning with positive (Appendix 3.13.2). The extremum is a constrained minimum if the principal minors of the Hessian matrix of the Lagrangean evaluated at the extremum are all negative. Whenever π_w is constant then (1) if the initial function is maximized, the same occurs with the Legendre transform and (2) if the initial function is minimized, the same occurs with the Legendre transform. This is because the Hessian matrix of the Lagrangean of the Hessian matrix of the Legendre transform.

In thermodynamics, intensive variables are constant when the system is no longer isolated but in contact with a reservoir that maintains, e.g., a constant temperature. The most common Legendre transforms in thermodynamics are (1) the *Helmholtz potential*, $F \equiv U - TS$; (2) the *Enthalpy*, $H \equiv U + PV$ and (3) the *Gibbs potential* $G \equiv U - TS + PV$ (for further details see Appendix 3.13.1). The extremum principles are: (1) the system reaches the minimum Helmholtz Energy subject to the reservoir temperature; (2) the system reaches the minimum Enthalpy subject to the reservoir pressure and (3) the system reaches the minimum Gibbs Energy subject to the reservoir pressure and temperature. The differential form of the Legendre transforms with constant intensive properties are:(1) dF = dU - TdS, (2) dH = dU + PdV, and (3) dG = dU - TdS + PdV.

For the consumer the Legendre transforms have an extremum principle when one or more of the marginal utilities are constant, i.e., the consumption of these goods is such that their marginal utilities are constant. For example: (1) Public Goods or non-excludable goods that have a null price and therefore null marginal utilities, i.e., these goods are consumed until the additional utility derived from their consumption is null and (2) rationed goods. These Legendre transforms have the following general form: $\Psi \equiv U - \sum_{i=1,l} x_i M U_i$. The extremum principle is that the system reaches the maximum Ψ subject to constant marginal utilities for the products x_i , $i = 1, \ldots, l$. In this case, the differential form of the Legendre transforms is: $d\Psi = dU - \sum_{i=1,l} M U_i dx_i = \sum_{i=l+1,n} M U_i dx_i$. The economic intuition is that the consumer maximizes the utility obtained with the consumption of goods $i = l + 1, \ldots, n$, because the amount of other goods is constrained in order to provide constant marginal utilities.

The complete Legendre transform $\Psi \equiv U - \sum_{i=1,n} x_i M U_i$ can be written as the difference in the utility gain from consumption and the cost of consumption, $\Psi = U - \frac{M U_q}{p_q} \sum_{i=1,n} x_i p_i$ where $\frac{M U_q}{p_q}$, is equal for all goods, in equilibrium. If marginal utilities are null then the complete Legendre transform is equal to the utility. Thus, the complete Legendre transform is a measure of the net benefits of consumption, i.e., *consumer's surplus* [38].

3.7 Elasticities

While the first derivatives of the fundamental equation are intensive variables, the second derivatives, i.e., elasticities, stand for important properties of the system. The nondimensional **elasticities** have the general form

$$q_{\Pi,X}^{ij} = \frac{\pi_j}{x_i} \left(\frac{\partial x_i}{\partial \pi_j}\right)_{\Pi,X},\tag{3.27}$$

where Π is the set of π_k variables kept constant (with $k \neq j$) and X is the set of x_z variables kept constant (with $z \neq i$) and the total number of variables kept constant is n-1. The absolute value of these non-dimensional elasticities has a straightforward interpretation: it is the ratio of the proportional change in x_i to the proportional change in π_j . These characteristics can be obtained empirically. Additionally,

$$\frac{1}{x_i}\frac{\partial(x_i\pi_i)}{\partial\pi_i} = 1 + \frac{\pi_i}{x_i}\frac{\partial x_i}{\partial\pi_i} = 1 + q_{\Pi,X}^{ii}, \qquad (3.28)$$

which means that for $x_i > 0$ if (1) q > -1 then $\pi_i x_i$ increases while if (2) q < -1 then $\pi_i x_i$ decreases. If the fundamental equation is first order homogeneous then $\pi_i x_i =$

 $\varphi(\pi_1, \ldots, x_i, \ldots, \pi_n)$ and the value of the non-dimensional elasticity determines the sign of the change in the potential given by the Legendre transform.

In thermodynamic systems, elasticities are descriptive of material properties. The most used elasticities are respectively the thermal expansion, the isothermal compressibility, the heat capacity at constant pressure, the adiabatic compressibility and the heat capacity at constant volume:

$$\alpha \equiv \frac{1}{V} \left(\frac{\partial V}{\partial T} \right)_{P,N}, \qquad (3.29)$$

$$\kappa_T \equiv -\frac{1}{V} \left(\frac{\partial V}{\partial P} \right)_{T,N}, \qquad (3.29)$$

$$c_P \equiv T \left(\frac{\partial S}{\partial T} \right)_{P,N}, \qquad (3.29)$$

$$\kappa_s \equiv -\frac{1}{V} \left(\frac{\partial V}{\partial P} \right)_{S,N}, \qquad (3.29)$$

$$\kappa_s \equiv T \left(\frac{\partial S}{\partial T} \right)_{V,N}, \qquad (3.29)$$

(for further details see Appendix 3.13.1).

In thermodynamics non-dimensional elasticities are not used although they would be useful given Eq. 3.28. For example, consider an open system with a movable and diathermic⁶ wall in contact with a temperature reservoir. If the pressure of the reservoir is increased then: (1) the system is compressed by the reservoir until the pressures are equal (2) a heat flow is established between the system and the reservoir to maintain the system at constant temperature and (3) a mass flow is established between the system and the reservoir to maintain the system at constant chemical potential. If the non-dimensional isothermal compressibility of the system $\kappa'_T = \kappa_T P > 1$ then the change in internal energy associated with the change in pressure will be positive, i.e., the change in the Legendre transform $d\phi(T, V, \mu) = d(-PV)$, while if $\kappa'_T < 1$, the change in internal energy associated with the change in pressure will be negative. This is summarized for all thermodynamic non-dimensional elasticities in Tab. 3.1.

In economic systems, elasticities are descriptive of consumer characteristics. The non-dimensional elasticities used in consumer theory are called elasticities:

$$\varepsilon_{\{p_k,\forall k\neq i\}} \equiv \frac{p_i}{x_i} \left(\frac{\partial x_i}{\partial p_i}\right)_{\{p_k,\forall k\neq i\}},\tag{3.30}$$

⁶A diathermic wall allows for energy transfer in the form of heat between the system and the environment.

Elasticities	Legendre transform behavior
κ_T'	$\phi(T,V,\mu)$ increases with P at constant T if $\kappa_T'>1,$ remains constant if
	$\kappa_T' = 1$ and decreases otherwise.
κ_s'	$\phi(T,V,\mu)$ increases with P at constant S if $\kappa_s'>1,$ remains constant if
	$\kappa_s' = 1$ and decreases otherwise.
c'_P	$\phi(S, -P, \mu)$ increases with T at constant P if $c'_P > -1^7$.
c'_v	$\phi(S, -P, \mu)$ increases with T at constant V if $c'_v > -1^8$.

Table 3.1: The relation between Adimensional Thermodynamic Elasticities and the change in the potentials given by the Legendre transforms of the Energy Function.

$$\begin{split} \varepsilon_{\{x_k,\forall k\neq j\}} &\equiv \frac{p_j}{x_j} \left(\frac{\partial x_j}{\partial p_j}\right)_{\{x_k,\forall k\neq j\}} \\ e_{\{p_k,\forall k\neq i\}} &\equiv \frac{p_i}{x_j} \left(\frac{\partial x_j}{\partial p_i}\right)_{\{p_k,\forall k\neq i\}}, \\ e_{\{x_k,\forall k\neq j\}} &\equiv \frac{p_i}{x_j} \left(\frac{\partial x_j}{\partial p_i}\right)_{\{x_k,\forall k\neq j\}}, \\ \mathfrak{e}_{\{p_k,\forall k\neq i\}} &\equiv \frac{w}{x_i} \left(\frac{\partial x_i}{\partial w}\right)_{\{p_k,\forall k\neq i\}}. \end{split}$$

The $\varepsilon_{\{p_k,\forall k\neq i\}}$ is the adimensionalization of the inverse demand curve. It is called the own price elasticity of demand. It measures how responsive demand is to changes in price [38]. Products are classified as *Giffen Goods* if $\varepsilon_{\{p_k,\forall k\neq i\}} > 0$. A product has an elastic demand if the absolute value of the own-price elasticity is higher than 1 and an inelastic demand otherwise. Eq. 3.28 says that if the demand of x_i is elastic then an increase in price will decrease the amount demanded so much that the revenue will decrease; if the demand is inelastic then an increase in price will not change demand very much and the overall revenue will increase. The $e_{\{p_k, \forall k \neq i\}}$ is called the cross-price elasticity of demand [13]. The $\varepsilon_{\{x_k,\forall k\neq j\}}$ and $e_{\{x_k,\forall k\neq j\}}$ are also prices elasticities but with the amounts of other products kept constant. The $e_{\{p_k,\forall k\neq i\}}$ is the adimensionalization of the inverse income curve. It is called the income elasticity of demand. It measures how responsive demand is to some change in income [38]. Products are classified according to their income elasticity: normal goods ($\mathfrak{e}_{\{p_k,\forall k\neq i\}} > 0$) and inferior goods ($\mathfrak{e}_{\{p_k,\forall k\neq i\}} < 0$). Normal goods can be either necessary goods ($\mathfrak{e}_{\{p_k,\forall k\neq i\}} < 1$) or luxury goods ($\mathfrak{e}_{\{p_k,\forall k\neq i\}} > 1$). In the consumer elasticities, prices are used instead of marginal utilities, because in equilibrium marginal utilities are proportional to prices.

y	$\left(\frac{\partial \pi_i}{\partial x_j}\right)_{\{x_k,\forall k\neq j\}}$	=	$\left(\frac{\partial \pi_j}{\partial x_i}\right)_{\{x_z,\forall z\neq i\}}$
φ_i	$\left(\frac{\partial \pi_l}{\partial x_j}\right)_{\pi_i, \{x_k, \forall k \neq i, j\}}$	=	$\left(\frac{\partial \pi_j}{\partial x_l}\right)_{\pi_i, \{x_z, \forall z \neq i, l\}}$
$\varphi_{i,g}$	$\left(\frac{\partial \pi_l}{\partial x_j}\right)_{\pi_i,\pi_g,\{x_k,\forall k\neq i,j,g\}}$	=	$\left(\frac{\partial \pi_j}{\partial x_l}\right)_{\pi_i,\pi_g,\{x_z,\forall z\neq i,l,g\}}$
φ_i	$\left(\frac{\partial \pi_k}{\partial \pi_i}\right)_{\{x_j, \forall j \neq i\}}$	=	$-\left(rac{\partial x_i}{\partial x_k} ight)_{\pi_i,\{x_j,orall j eq i,k\}}$
$\varphi_{i,g}$	$\left(\frac{\partial \pi_k}{\partial \pi_i}\right)_{\pi_g, \{x_j, \forall j \neq i, g\}}$	=	$-\left(\frac{\partial x_i}{\partial x_k}\right)_{\pi_i,\pi_g,\{x_j,\forall j\neq i,k,g\}}$
$\varphi_{i,g}$	$\left(\frac{\partial x_i}{\partial \pi_g}\right)_{\pi_i, \{x_j, \forall j \neq i, g\}}$	=	$\left(\frac{\partial x_g}{\partial \pi_i}\right)_{\pi_g, \{x_j, \forall j \neq i, g\}}$
φ	$\left(\frac{\partial x_i}{\partial \pi_g}\right)_{\{\pi_j,\forall j\neq g\}}$	=	$\left(\frac{\partial x_g}{\partial \pi_i}\right)_{\{\pi_j,\forall j\neq i\}}$

Table 3.2: Maxwell Relations obtained from $y = y(x_1, ..., x_n)$ and its Legendre transforms $\varphi_i = \varphi_i(x_1, ..., \pi_i, ..., x_n)$, $\varphi_{i,g} = \varphi_{i,g}(x_1, ..., \pi_i, \pi_g, ..., x_n)$ and $\varphi = \varphi(\pi_1, ..., \pi_n)$.

3.8 Maxwell Relations

The **Maxwell Relations** result from the equality of mixed partial derivatives valid for scalar functions whose second derivatives exist and are continuous (Young's Theorem). The equality between the mixed partial derivatives of the fundamental equation $y = y(x_1, \ldots, x_n)$ implies that:

$$\left(\frac{\partial \pi_i}{\partial x_j}\right)_{x_{k\neq j}} = \left(\frac{\partial \pi_j}{\partial x_i}\right)_{x_{k\neq i}}.$$
(3.31)

The equality between the mixed partial derivatives is also applicable to the other formulations of the fundamental equation, i.e., the Duality Formulation and the Legendre transforms. The Maxwell Relations for the Legendre transforms are given in Tab. 3.2.

Maxwell relations are important because they make explicit that not all second mixed derivatives are independent. In fact, all second derivatives can be written as a function of a small set of independent variables. The relations between the elasticities,

. . .

$$q_{\{\pi_i,\forall i\neq k\}}^{kk} = q_{x_j,\{\pi_i,\forall i\neq k,j\}}^{kk} + \frac{\left(q_{\{\pi_i,\forall i\neq k\}}^{jk}\right)^2}{q_{\{\pi_i,\forall i\neq j\}}^{jj}} \frac{\pi_j x_j}{\pi_k x_k},$$
(3.32)

$$q_{x_{j},\{\pi_{i},\forall i\neq k,j\}}^{kk} = q_{x_{j},x_{z},\{\pi_{i},\forall i\neq k,j,z\}}^{kk} + \frac{\left(q_{x_{j},\{\pi_{i},\forall i\neq k,j\}}^{zk}\right)^{2}}{q_{x_{j}\{\pi_{i},\forall i\neq j,z\}}^{zz}} \frac{\pi_{z}x_{z}}{\pi_{k}x_{k}},$$
(3.33)

U(S, V, N)	$\left(\frac{\partial T}{\partial V}\right)_{S,N} = -\left(\frac{\partial P}{\partial S}\right)_{V,N}$	$\left(\frac{\partial T}{\partial N}\right)_{S,V} = \left(\frac{\partial \mu}{\partial S}\right)_{V,N}$	$\left(\frac{\partial P}{\partial N}\right)_{S,V} = -\left(\frac{\partial \mu}{\partial V}\right)_{S,N}$
F(T, V, N)	$\left(\frac{\partial S}{\partial V}\right)_{T,N} = \left(\frac{\partial P}{\partial T}\right)_{V,N}$	$\left(\frac{\partial S}{\partial N}\right)_{T,V} = -\left(\frac{\partial \mu}{\partial T}\right)_{V,N}$	$\left(\frac{\partial P}{\partial N}\right)_{T,V} = -\left(\frac{\partial \mu}{\partial V}\right)_{T,N}$
H(S, P, N)	$\left(\frac{\partial T}{\partial P}\right)_{S,N} = \left(\frac{\partial V}{\partial S}\right)_{P,N}$	$\left(\frac{\partial T}{\partial N}\right)_{S,P} = \left(\frac{\partial \mu}{\partial S}\right)_{P,N}$	$\left(\frac{\partial V}{\partial N}\right)_{S,P} = \left(\frac{\partial \mu}{\partial P}\right)_{S,N}$
G(T, P, N)	$\left(\frac{\partial S}{\partial P}\right)_{T,N} = -\left(\frac{\partial V}{\partial T}\right)_{P,N}$	$\left(\frac{\partial S}{\partial N}\right)_{T,P} = -\left(\frac{\partial \mu}{\partial T}\right)_{P,N}$	$\left(\frac{\partial V}{\partial N}\right)_{T,P} = \left(\frac{\partial \mu}{\partial P}\right)_{T,N}$

Table 3.3: Thermodynamic Maxwell Relations obtained from U, F, H and G.

$$q_{\pi_{l},\{x_{i},\forall i\neq l,k\}}^{kk} = q_{\{x_{i},\forall i\neq k\}}^{kk} + \frac{\left(q_{\pi_{l},\{x_{i},\forall i\neq l,k\}}^{lk}\right)^{2}}{q_{\pi_{k},\{x_{i},\forall i\neq l,k\}}^{ll}} \frac{\pi_{l}x_{l}}{\pi_{k}x_{k}},$$
(3.34)

are obtained using the Maxwell relations (see Appendix 3.13.3 for further details). The relation between $q_{\pi_{i\neq k}}^{kk}$ and $q_{x_{i\neq k}}^{kk}$ (see also Appendix 3.13.3 for further details) is:

$$q_{\{\pi_{i},\forall i \neq k\}}^{kk} = q_{\{x_{i},\forall i \neq k\}}^{kk} + \sum_{\substack{j \neq k}} \frac{x_{j}\pi_{j}}{x_{k}\pi_{k}} \frac{q_{\{\pi_{i},\forall i \neq k\}}^{jk} q_{\pi_{j},\{x_{i},\forall i \neq j,k\}}^{jk}}{q_{\pi_{k},\{x_{i},\forall i \neq j,k\}}^{jj}}.$$
(3.35)

The Maxwell relations that result from the Energy representation of the fundamental equation and its Legendre transforms are given in Tab. 3.3 and the relations between the thermodynamic elasticities for constant mole numbers are presented next.

The relation between the specific heat at constant pressure and the specific heat at constant volume,

$$c_P = c_v + \frac{T v \alpha^2}{\kappa_T},\tag{3.36}$$

is obtained using Eq. 3.33 and the relation between the adiabatic compressibility and compressibility at constant temperature,

$$\kappa_T = \kappa_s + \frac{T v \alpha^2}{c_P},\tag{3.37}$$

is obtained using Eq. 3.33 and the Maxwell Relation $\left(\frac{\partial S}{\partial P}\right)_{T,N} = -\left(\frac{\partial V}{\partial T}\right)_{P,N}$. Eq. 3.36 and 3.37 imply that only three of five coefficients defined by Eq. 3.38 are independent. The relation between specific heats and compressibilities is

$$\frac{c_P}{c_v} = \frac{\kappa_T}{\kappa_s}.$$
(3.38)

The Maxwell relations that result from the Utility function and its Legendre transforms are given in Tab. 3.4 and the relations between the economic elasticities for two products are summarized next.

$U(x_1,\ldots,x_n)$	$\left(\frac{\partial p_i}{\partial x_j}\right)_{\{x_k,\forall k\neq j\}}$	=	$\left(\frac{\partial p_j}{\partial x_i}\right)_{\{x_k,\forall k\neq i\}}$
$\varphi_i(x_1,\ldots,p_i,\ldots,x_n)$	$-\left(\frac{\partial p_j}{\partial p_i}\right)_{\{x_k,\forall k\neq i\}}$	=	$\left(\frac{\partial x_i}{\partial x_j}\right)_{p_i,\{x_k,\forall k\neq i,j\}}$
$\varphi_i(x_1,\ldots,p_i,\ldots,x_n)$	$\left(\frac{\partial p_j}{\partial x_k}\right)_{p_i,\{x_l,\forall l\neq i,k\}}$	=	$\left(\frac{\partial p_k}{\partial x_j}\right)_{p_i,\{x_l,\forall l\neq i,j\}}$
$\varphi(p_1,\ldots,p_i,\ldots,p_m)$	$\left(\frac{\partial p_j}{\partial x_i}\right)_{\{p_k,\forall k\neq j\}}$	=	$\left(\frac{\partial p_i}{\partial x_j}\right)_{\{p_k,\forall k\neq i\}}$

Table 3.4: Economic Maxwell Relations obtained from U, φ_i and φ .

The relation between the own price elasticity of demand at constant prices and the own price elasticity of demand at constant amounts for product 2,

$$\varepsilon_{p_2} = \varepsilon_{x_2} + e_{p_2}^2 \frac{1}{\varepsilon_{p_1}} \frac{p_2 x_2}{p_1 x_1},$$
(3.39)

is obtained using Eq. 3.33 and for product 1,

$$\varepsilon_{p_1} = \varepsilon_{x_1} + e_{p_2}^2 \frac{1}{\varepsilon_{p_2}} \frac{p_2 x_2}{p_1 x_1},$$
(3.40)

is obtained using Eq. 3.33 and the Maxwell relation $\left(\frac{\partial x_1}{\partial p_2}\right)_{p_1,N} = \left(\frac{\partial x_2}{\partial p_1}\right)_{p_2}$. Eq. 3.39 and 3.40 imply that only three of the elasticities defined by Eq. 3.30 are independent. The relation between the own price elasticities at constant prices and at constant amounts is

$$\frac{\varepsilon_{p_1}}{\varepsilon_{p_2}} = \frac{\varepsilon_{x_1}}{\varepsilon_{x_2}}.$$
(3.41)

3.9 Stability Conditions and the Le Chatelier Principle

The **stability of the equilibrium state** of a function y depends on the sign of d^2y . If the equilibrium is a maximum, the function is stable if it is strictly concave, i.e., $d^2y < 0$; if the equilibrium is a minimum, the function is stable if it is strictly convex, i.e., $d^2y > 0$. The sufficient condition for the twice continuously differentiable real-valued function y on a set C to be (1) strictly convex is that its Hessian matrix is positive definite, i.e., all sub matrices of the Hessian matrix defined by the first k rows and k columns have positive determinants and to be (2) strictly concave is that its Hessian matrix is negative definite, i.e., the determinants of the Hessian matrix are negative if k is odd and positive if k is even [22]. These necessary and sufficient conditions impose certain constraints on the signs of the elasticities (second derivatives).

Function $y(x_1, \ldots, x_n)$ is strictly concave if $(\partial \pi_i)/(\partial x_i)_{\{x_k, \forall k \neq i\}} < 0$, i.e., $q_{\{x_k, \forall k \neq i\}}^{kk} < 0$ if $\pi_i > 0$ and $x_i > 0$. The Legendre transforms of the concave function y are convex

with respect to its intensive variables (a proof is presented in sec. 3.6), i.e., its second derivatives are positive

$$\frac{\partial^2 \varphi(x_1, \dots, x_l, \pi_{l+1}, \dots, \pi_n)}{\partial \pi_{l+1}^2} = -\frac{\partial x_{l+1}(x_1, \dots, x_l, \pi_{l+1}, \dots, \pi_n)}{\partial \pi_{l+1}} = -\frac{x_{l+1}}{\pi_{l+1}} q_{\Pi, X}^{l+1, l+1} > 0,$$
(3.42)

with $\Pi = \{\pi_j, j = l+1, \ldots, n\}$ and $X = \{x_i, i = 1, \ldots, l\}$. Thus, the elasticity $q_{\Pi,X}^{l+1,l+1}$ in Eq. 3.42 is negative if $\pi_i > 0$ and $x_i > 0$. Additionally, Eq. 3.32 to Eq. 3.34 imply the Generalized Le Chatelier Principle: $0 > q_{\{x_i, \forall i \neq k\}}^{kk} > \ldots > q_{x_j, \{\pi_i, \forall i \neq k, j\}}^{kk} > q_{\{\pi_i, \forall i \neq k\}}^{kk}$.

The Generalized Le Chatelier Principle relations describe the behavior of a stable system displaced from equilibrium by a change in its extensive variable dx_k : (1) its conjugate intensive variable will change by $d\pi_k$ and (2) the absolute value of $d\pi_k$ increases with the number of extensive variables kept constant.

The entropy is strictly concave if $(\partial^2 S/\partial U^2) < 0$, i.e., $c_v > 0$. The Gibbs Energy is strictly concave on its intensive independent variables T and P, if $(\partial^2 G/\partial T^2) < 0$, i.e., $c_P > 0$, and that $(\partial^2 G/\partial P^2) < 0$, i.e., $\kappa_T > 0$. These constraints on c_v , c_P and κ_T together with Eq. 3.38 imply that $\kappa_s > 0$. Additionally, Eq. 3.36 and Eq. 3.37 imply that $c_P > c_v$ and $\kappa_T > \kappa_s$.

The conditions $(\partial^2 S/\partial U^2) < 0$ and $(\partial^2 S/\partial V^2) < 0$ imply that: (1) the amount of entropy increase that results from an unit increase in internal energy at constant volume (due to a heat flow) decreases with the internal energy of the system and (2) the amount of entropy increase that results from an unit increase of volume at constant internal energy (due to both heat and work flows) decreases with the volume of the system.

Conditions $c_P > c_v > 0$ and $\kappa_T > \kappa_s > 0$ embody the Generalized Le Chatelier Principle for thermodynamic systems: the absolute change in temperature triggered by a change in entropy is higher for a system at constant volume than for a system at constant pressure and the absolute change in pressure triggered by a change in volume is higher for a system at constant entropy than for a system at constant pressure. As an example, we can think of a diathermic cylinder fitted with a piston that is in contact with a temperature reservoir. Suppose that the piston is pulled and an increase in volume occurs. The system will decrease its pressure ($\kappa_T > 0$), but this decrease will be smaller ($\kappa_T > \kappa_s > 0$) than at constant entropy, because the system receives a flow of heat from the temperature

Thermodynamic Constraints	Economic Constraints
$c_P > c_v > 0$	$0 > \varepsilon_{\{x_k, \forall k \neq i\}} > \varepsilon_{\{p_k, \forall k \neq i\}}$

Table 3.5: Constraints on the thermodynamic and economic elasticities imposed by stability.

reservoir.

The utility is strictly concave if $(\partial^2 U/\partial x_i^2) < 0$, i.e., $\varepsilon_{\{x_k,\forall k\neq i\}} < 0$. The Economic Potential Ψ is strictly convex on its intensive variables MU_i if $\varepsilon_{\{p_k,\forall k\neq i\}} < 0$ and, Eq. 3.39 and Eq. 3.40 imply that $\varepsilon_{\{x_k,\forall k\neq i\}} > \varepsilon_{\{p_k,\forall k\neq i\}}$ (assuming that prices and amounts are positive). These stability results imply that Giffen goods do not exist if the equilibrium state is stable⁹.

The condition $(\partial^2 U/\partial x_i^2) < 0$ corresponds to the 'Law' of Diminishing Marginal Utility: the utility gained from an extra increase of one unit of product x_i decreases with the amount of x_i already owned.

Conditions $0 > \varepsilon_{\{x_k, \forall k \neq i\}} > \varepsilon_{\{p_k, \forall k \neq i\}}$ embody the Generalized Le Chatelier Principle for the consumer: the absolute change in demand triggered by a change in price is higher for constant product amounts than for constant prices. As an example, we can think of a consumer in contact with a market that has only two goods. If the price of product p_1 decreases the consumer buys an additional amount of good x_1 ($\varepsilon_{p_2} < 0$). The amount of x_1 traded is smaller if the consumer can also trade product 2 with the market ($0 > \varepsilon_{x_2} > \varepsilon_{p_2}$).

Tab. 3.5 summarizes the constraints imposed by stability on the thermodynamic and economic elasticities.

3.10 Equations of State and Integrability

Equations of state are equations that establish a relation in equilibrium between one intensive variable and the independent extensive variables. They have the following functional form

$$\pi_i = \pi_i \left(x_1, x_2, \dots, x_n \right).$$

The knowledge of all equations of state is equivalent to the knowledge of the fundamental equation under certain conditions. This is known as the **integrability problem** and it can be summarized as follows: 'When is it possible to build a fundamental function using the equations of state?'. First, mathematical integrability implies that the fundamental

⁹Note that this would not occur if only quasi-concavity of the utility function was required (remark pointed out by one of the reviewers).

equation must satisfy the Maxwell relations. Second, the optimization behavior embedded in the fundamental equation implies that the equations of state do not contradict that the behavior is indeed optimized, i.e., the Generalized Le Chatelier Principle is verified. These two conditions are equivalent to imposing that the Hessian matrix of the fundamental function is (1) symmetric and (2) negative semidefinite. If the fundamental equation represents an order relation¹⁰ then these two conditions are met (for further details see Candeal et al. [6]).

In the thermodynamic system, the equations of state, derived from the fundamental equation written in the energy representation, are: $T = T(S, V, N_1, ..., N_r)$, $P = P(S, V, N_1, ..., N_r)$ and $\mu_j = \mu_j(S, V, N_1, ..., N_r)$ with j = 1, ..., r (see Appendix 3.13.1 for further details). These equations can be computed based on observed behavior. The energy fundamental equation can be built knowing all the equations of state by using Eq. 3.18. We can identify the first condition of integrability in thermodynamics as the Maxwell relations and the second as the constraints imposed on the thermodynamic coefficients by the Generalized Le Chatelier Principle. In thermodynamics, the entropy represents a preference relation imposed by the second law (for further details see Cooper [7]); thus, it is always possible to obtain an entropy function from the equations of state.

In the consumer, the equations of state, derived from the fundamental equation written in the expenditure representation, are: $p_j = p_j(x_1, ..., x_n)$. These equations can be computed based on observed behavior. The expenditure function can be built knowing all the Walrasian demand functions (obtained from the equations of state) by using Eq. 3.20. The first condition of integrability is known in economics as *Antonelli's integrability condition* and the second condition, i.e., the constraints imposed on economic elasticities, is known as *'economic integrability'* [17]. In the consumer the utility represents a preference relation if and only if the consumer is rational; thus, it might not be possible to obtain a utility function from the equations of state.

3.11 First Order Phase Transitions

Phase transitions occur where the fundamental equation is unstable (at least one of the stability criteria is violated). In Fig.3.1, the fundamental equation is unstable where $q_{\{\pi_i, \forall i \neq i\}}^{ii} > 0$, assuming that intensive and extensive variables are positive. In this case,

¹⁰A function f represents a preference relation \leq if: $x \leq y \Leftrightarrow f(x) \leq f(y)$.

Equilibrium Econophysics



Figure 3.1: First Order Phase Transition.

the fundamental equation is no longer valid, i.e., the system is not characterized by the equilibrium conditions. However, the system can be decomposed into *phases*, i.e, subsystems in equilibrium.

The two states¹¹ between which a first order transition occurs are distinct and are characterized (1) by being in equilibrium with each other, i.e., by having equal intensive variables [14],

$$\pi_i^1 = \pi_i^2, \qquad j = 1, \dots, n,$$
(3.43)

where π_j^k is the intensive variable π_j in phase k, k = 1, 2 and (2) by having at least one extensive variable that is different $x_j^1 \neq x_j^2$ where x_j^k is the extensive variable x_j in phase k. Conditions (1) and (2) imply that (3) $q_{\{\pi_i,\forall i\neq j\}}^{jj} = \infty$.

The total number of independent variables needed to characterize the phase transition is n because Eq. 3.43 applies. If the function y is homogeneous first order then the **Gibbs-Duhem Equation**,

$$0 = x_1^k d\pi_1 + \ldots + x_n^k d\pi_n, (3.44)$$

applies to each phase k (see Appendix 3.13.4 for further details) and the number of independent intensive variables or degrees of freedom decreases to f = n - 2 because (1) the intensive variable π_j can be written as a function of the other intensive variables (eq. 3.44) and (2) Eq. 3.43 applies. This is the **Generalized Gibbs phase rule**. Thus, the minimum number of extensive variables that are needed to characterize the phase transition is n - f.

The relation between the intensive variables $\pi_p \equiv \pi_p^1 = \pi_p^2$ and $\pi_g \equiv \pi_g^1 = \pi_g^2$ in

¹¹More than two states are possible but will not be considered in this chapter.

equilibrium can be computed with the Generalized Clapeyron Equation:

$$\left(\frac{\partial \pi_p}{\partial \pi_g}\right)_{\{\pi_i^1 - \pi_i^2, \forall i \neq p, g\}} = -\frac{\left(\frac{\partial (\pi_k^1 - \pi_k^2)}{\partial \pi_g}\right)_{\{\pi_i^1 - \pi_i^2, \forall i \neq k, g\}}}{\left(\frac{\partial (\pi_k^1 - \pi_k^2)}{\partial \pi_p}\right)_{\{\pi_i^1 - \pi_i^2, \forall i \neq k, p\}}} = -\frac{\left(\frac{\partial \pi_k^1}{\partial \pi_g}\right)_{\{\pi_i^1 - \pi_i^2, \forall i \neq k, g\}}}{\left(\frac{\partial \pi_k^1}{\partial \pi_p}\right)_{\{\pi_i^1 - \pi_i^2, \forall i \neq k, p\}}} - \left(\frac{\partial \pi_k^2}{\partial \pi_p}\right)_{\{\pi_i^1 - \pi_i^2, \forall i \neq k, p\}}}.$$
(3.45)

where $\pi_i^1 - \pi_i^2$ is constant and equal to zero (Eq. 3.43). If the potential y is a first order homogeneous function then Eq. 3.45 simplifies to

$$\left(\frac{\partial \pi_p}{\partial \pi_g}\right)_{\{\pi_i^1 - \pi_i^2, \forall i \neq p, g\}} = -\frac{\frac{x_g^1}{x_k^1} - \frac{x_g^2}{x_k^2}}{\frac{x_p^1}{x_k^1} - \frac{x_p^2}{x_k^2}},$$
(3.46)

by computing the partial derivatives using the Gibbs-Duhem Equation (Eq. 3.44) for each phase.

The system goes through a heterogeneous state where both phases coexist, with a fraction of the system in one phase and the remainder in the other. The intensive variables π_i that characterize the line that links the two phases (line A in Fig. 3.1) are such that: $\sum_{i \neq j} \int_{\pi_i^1}^{\pi_i^2} (x_i^1 - x_i^2) d\pi_i = 0$; this is the **Maxwell construction** proved next. Each point in a vertical line in Fig. 3.1 has a different π_i for $i \neq j$; therefore, vertical lines have a variable value of the Legendre transform $\varphi(\pi_1, \ldots, \pi_n)$. The horizontal line or π_i that correspond to the phase transition are such that the Legendre transform is at its maximum:

$$d\varphi = \sum_{i \neq j} x_i d\pi_i = 0. \tag{3.47}$$

By integrating Eq. 3.47 we obtain the Maxwell construction. Its geometrical meaning is best understood for the case where all intensive variables are kept constant with the exception of π_k : $\int_{\pi_k^1}^{\pi_k^2} x_k d\pi_k = 0$, i.e., the area below the dashed line must be equal to the area above. However, if the function y is homogeneous first order then the Legendre transform $\varphi(\pi_1, \ldots, \pi_n)$ is always null because Eq. 3.44 applies. In this case, the adequate Legendre transform has one extensive variable, $\varphi(x_1, \pi_2, \ldots, \pi_n)$. Also, the horizontal line or the π_i that corresponds to the phase transition is such that the Legendre transform is at its maximum. With x_1 constant:

$$d\varphi = \sum_{i \neq 1,j} x_i d\pi_i = 0. \tag{3.48}$$

Also for systems where x_1 is constant and all intensive variables with the exception of π_k are also kept constant: $\int_{\pi_k^1}^{\pi_k^2} x_k d\pi_k = 0.$

The system moves from one phase to the other by exchanging extensive variables with the exterior, i.e., **flows**, subject to the constraints imposed:

$$g^z = 0, \quad z = 1, \dots, m.$$
 (3.49)

The amount of each extensive variable is computed using the specific extensive variables, $\mathcal{X}_i = \frac{x_i}{X}$, where X is

$$X = X^1 + X^2 \equiv \alpha X + (1 - \alpha)X$$

 X^1 is the amount of X in phase 1 and α is the fraction of X in phase 1. Using the specific extensive variables the amount of x_z in each phase is

$$\begin{aligned} x_z^1 &= X \alpha \mathcal{X}_z^1, \\ x_z^2 &= X(1-\alpha) \mathcal{X}_z^2. \end{aligned}$$

Thus the amount of x_z is

$$x_{z} = x_{z}^{1} + x_{z}^{2} = X\alpha \mathcal{X}_{z}^{1} + X(1-\alpha)\mathcal{X}_{z}^{2}.$$
(3.50)

This is the **Generalized Lever Rule**, i.e., the set of variables x_z , z = 1, ..., n is a linear combination of the sets of specific variables \mathcal{X}_z^k , k = 1, 2. With the set of Eq. 3.50 and the constraints imposed by Eq. 3.49 it is possible to compute x_z , z = 1, ..., n. Whenever y is first order homogeneous function, the specific variables \mathcal{X}_z^k are independent of the size of the system and therefore the set of independent intensive variables completely specify the set of specific variables \mathcal{X}_z^k .

In a thermodynamic system, a phase transition occurs where the entropy fundamental equation is unstable, i.e., it is no longer concave. Fig. 3.2 shows where the stability criteria are violated: where the compressibility at constant temperature, κ_T , becomes negative. In this case, the system is no longer homogeneous but two different phases coexist. The two phases are characterized by: (1) equal temperature, pressure and chemical potential for



Figure 3.2: First Order Phase Transition in Thermodynamics.

each chemical species and (2) different volumes, number of moles and internal energy. These conditions imply that $c_P = \infty$, $\alpha = \infty$, $\kappa_T = \infty$.

The number of independent intensive variables, given by the Gibbs phase rule, for thermodynamics, is r = (r + 2) - 2 where r is the number of different chemical species and r+2 is the number of extensive variables. For systems with constant number of moles, constant temperature and constant chemical potentials for all species, the horizontal line of the phase transition is such that: $\int_{P^1}^{P^2} V dP = 0$ (Fig. 3.2).

In the thermodynamic phase transition, the change in pressure with temperature given by the Clapeyron Equation (Eq. 3.46) is

$$\left(\frac{\partial P}{\partial T}\right)_{\{\mu^{1}=\mu^{2}\}} = \frac{\frac{S^{1}}{N^{1}} - \frac{S^{2}}{N^{2}}}{\frac{V^{1}}{N^{1}} - \frac{V^{2}}{N^{2}}} = \frac{\Delta s}{\Delta v},$$
(3.51)

obtained for a system with one chemical component using Eq. 3.46 with $\pi_p = -P$ and $\pi_g = T$ and where Δs and Δv are the changes in specific entropy, $s \equiv S/N$, and specific volume, $v \equiv V/N$, between the two phases. The other relations that can be obtained are:

$$\left(\frac{\partial\mu}{\partial T}\right)_{\{P^1=P^2\}} = -\frac{\frac{S^1}{V^1} - \frac{S^2}{V^2}}{\frac{N^1}{V^1} - \frac{N^2}{V^2}} = -\frac{s^1v^2 - s^2v^1}{\Delta v},\tag{3.52}$$

and

$$\left(\frac{\partial\mu}{\partial P}\right)_{\{T^1=T^2\}} = \frac{\frac{V^1}{S^1} - \frac{V^2}{S^2}}{\frac{N^1}{S^1} - \frac{N^2}{S^2}} = \frac{v^1 s^2 - v^2 s^1}{\Delta s}.$$
(3.53)

In a thermodynamic system, the phase transition occurs as the system exchanges energy and/or volume, in the form of heat and/or work with the environment, i.e., U and/or V change during the phase transition, at constant total number of moles

$$N \equiv N^{1} + N^{2} = \alpha N + (1 - \alpha)N, \qquad (3.54)$$

where α is the fraction of N in phase 1. The aggregate amounts of volume and internal energy are given by the *Lever Rule*: $V = V^1 + V^2 = \alpha Nv^1 + (1 - \alpha)Nv^2$ and $U = U^1 + U^2 = \alpha Nu^1 + (1 - \alpha)Nu^2$, where $u \equiv U/N$ is the specific energy.

Phase transitions in consumer theory occur where the utility function is unstable (no longer concave) or the expenditure function is unstable (no longer convex). Fig. 3.3 shows where the stability criteria are violated: the own price elasticity of demand, ε_{p_k} , becomes positive. In this case, the consumer has two different stable consumption bundles. The two consumption bundles are characterized by: (1) equal MU_i and (2) different values of the variables x_i . These conditions imply that $\varepsilon_{\{p_k,\forall k\neq i\}} = \infty$, $e_{\{p_k,\forall k\neq i\}} = \infty$, $e_{\{p_k,\forall k\neq i\}} = \infty$.

The number of independent intensive variables, given by the Gibbs phase rule, for the consumer, is n and decreases to n-2 is the utility is a first order homogeneous function. For consumers in a market with constant prices for all products except product k, the horizontal line of the phase transition is such that: $\int_{1}^{p_k^2} x_k dp = 0$.

In the consumer phase transition, the change in price of good i with price of good j, is

$$\left(\frac{\partial p_{i}}{\partial p_{j}}\right)_{\{p_{z}^{1}=p_{z}^{2},\forall z\neq i,j\}} = -\frac{\left(\frac{\partial p_{k}^{1}}{\partial p_{j}}\right)_{\{p_{z}^{1}=p_{z}^{2},\forall z\neq k,j\}} - \left(\frac{\partial p_{k}^{2}}{\partial p_{j}}\right)_{\{p_{z}^{1}=p_{z}^{2},\forall z\neq k,j\}}}{\left(\frac{\partial p_{k}^{1}}{\partial p_{i}}\right)_{\{p_{z}^{1}=p_{z}^{2},\forall i\neq k,i\}} - \left(\frac{\partial p_{k}^{2}}{\partial p_{i}}\right)_{\{p_{z}^{1}=p_{z}^{2},\forall i\neq k,i\}}}.$$
(3.55)

obtained using Eq. 3.45 with $\pi_p = p_i$ and $\pi_g = p_j$. If the utility is a first order homogeneous function then Eq. 3.55 simplifies to:

$$\frac{\partial p_i}{\partial p_j}_{\{p_z^1 = p_z^2, \forall z \neq i, j\}} = -\frac{\frac{x_j^1}{x_z^1} - \frac{x_j^2}{x_z^2}}{\frac{x_1^1}{x_z^1} - \frac{x_i^2}{x_z^2}}.$$
(3.56)

In the consumer system, the phase transition occurs as the consumer exchanges goods with the market at constant prices. Thus, a phase transition in consumer theory, can be thought of as a transition between two separate points in the utility function, motivated by an increasing (or decreasing) endowment:

$$w \equiv w^{1} + w^{2} = \alpha w + (1 - \alpha)w,$$
 (3.57)

where α is the fraction of w spent in phase 1. The aggregate amounts of each product are given by the *Consumer Lever Rule*: $p_i x_i = p_i x_i^1 + p_i x_i^2 = \alpha w \mathcal{X}_i^1 + (1 - \alpha) w \mathcal{X}_i^2$, where $\mathcal{X}_i^1, \mathcal{X}_i^2$ are the equilibrium amounts of endowment spent on x_i^k per unit of endowment spent in phase k.



Figure 3.3: First Order Phase Transition in the Consumer.

3.12 Conclusions

We have contributed to the unification of knowledge between Equilibrium Thermodynamics and Neoclassical Microeconomics by building a formalism common to both fields (Tab. 3.6 and 3.7). This general formalism includes (1) a derivation of a general relation between non-dimensional elasticities (Eq. 3.32 to Eq. 3.34) and (2) a derivation of the general formalism of first order phase transitions including a Generalized Clapeyron Equation (Eq. 3.45).

We obtained the following results in thermodynamics: (1) a relation between the non-dimensional thermodynamic coefficients and the change in Legendre transforms which highlights the importance of the non-dimensional coefficients in thermodynamics (Sec. 3.7), (2) the distinction between mathematical integrability and the optimization (Sec. 3.10) and (3) the generalization of the Clapeyron Equation (Eq. 3.52 and Eq. 3.53).

We obtained the following results in economics: (1) the definition of irreversibility in economic behavior (Sec. 3.4), (2) the clarification that the Engel curve and the offer curve are not descriptions of real processes dictated by the maximization of utility at constant endowment (Sec. 3.5), (3) a relation between elasticities (Eq. 3.39 to 3.41), (4) the proof that *Giffen Goods* do not exist in a stable equilibrium (Sec. 3.9), (5) the establishment that 'economic integrability' is equivalent to the Generalized Le Chatelier Principle (Sec. 3.10) and (6) the definition of a phase transition of first order characterized by an infinite own price elasticity of demand (Sec. 3.11). The importance of phenomena similar to phase transitions in economics has been recently highlighted for financial markets [21, 37] and for an Ising economy [3]. With the general formalism provided in this chapter for the first order phase transition, this concept can be applied more rigorously and extended to other fields of economics.

Differences between the formalism of thermodynamics and the microeconomic for-

	General Formalism	Thermodynamic Formalism
Fundamental Function	$y(x_1,\ldots,x_n)$	$S(U^1, U^2, V^1, V^2, N_1^1, \dots, N_r^2)$
Extensive Variables	x_i	U^j,V^j,N^j_i
Intensive Variables	π_i	$rac{1}{T^j}, rac{P^j}{T^j}, -rac{\mu_i^j}{T^j}$
Duality Function	$x_1(y, x_2, \ldots, x_n)$	$U(S^1, S^2, V^1, V^2, N_1^1, \dots, N_r^2)$
Irreversibility	Increase in y at constant g^z	Increase in S at constant U, V, N_i
Value Function	$y^*(k_1,\ldots,k_l)$	$S^*(U, V, N_1, \ldots, N_r)$
Legendre Transforms	$\varphi(x_1,\ldots,\pi_i,\ldots,x_n)$	F = U - TS
		H = U + PV
		G = U - TS + PV
Elasticities	$q_{\Pi,X}^{ij}$	$\alpha, c_v, c_P, \kappa_s, \kappa_T$
Maxwell Relations		$\frac{c_P}{c_v} = \frac{\kappa_T}{\kappa_s}$
Le Chatelier Principle	$0 > q_{\{x_i, \forall i \neq k\}}^{kk} > \dots$	$c_P > c_v > 0$ and $\kappa_T > \kappa_s > 0$
	$> q_{x_j,\{\pi_i,\forall i \neq k,j\}}^{kk} > q_{\{\pi_i,\forall i \neq k\}}^{kk}$	
Equations of State	$\pi_i(x_1,\ldots,x_n)$	$T(S, V, N_1, \ldots, N_r)$
		$P(S, V, N_1, \ldots, N_r)$
		$\mu_i(S, V, N_1, \ldots, N_r)$
Integrability	Hessian matrix is: symmetric	Maxwell Relations
	and negative semidefinite	Constraints on elasticities
Phase transition	$q^{jj}_{\{\pi_i,\forall i\neq j\}} = \infty$	$c_P = \infty, \alpha = \infty, \kappa_T = \infty$
Maxwell construction	$\pi_k \colon \int_{\pi_k^1}^{\pi_k^2} x_k d\pi_k = 0$	$\int_{P^1}^{P^2} V dP = 0$

Table 3.6: Table of Correspondences I.

malism are: (1) the fundamental equation in thermodynamics must be a homogeneous first order equation because entropy is additive over the subsystems while in microeconomics the utility is not ¹², (2) the integrability problem is solved in microeconomics by imposing that the consumer is always rational (which is not always empirically true) while in thermodynamics it is solved by imposing the second law (which has always been empirically true), (3) there is no equivalent to the Expenditure Duality in thermodynamics because thermodynamic intensive variables do not all have the same meaning and (4) the composite vs. simple systems issue only exists in thermodynamics because the entropy is only defined in equilibrium while utility is defined out of equilibrium. The measurement

¹²In the field of Nonextensive Thermodynamics entropy is no longer a homogeneous first order function.

	General Formalism	Economic Formalism
Fundamental Function	$y(x_1,\ldots,x_n)$	$U(x_1,\ldots,x_n)$
Extensive Variables	x_i	x_i
Intensive Variables	π_i	MU_i
Duality Function	$x_1(y, x_2, \ldots, x_n)$	$x_1(U, x_2, \ldots, x_n)$
		$w(x_1,\ldots,x_n)$
Irreversibility	Increase in y at constant g^z	Increase in U at constant w
Value Function	$y^*(k_1,\ldots,k_l)$	$U^*(w, p_1, \ldots, p_n)$
Legendre Transforms	$\varphi(x_1,\ldots,\pi_i,\ldots,x_n)$	$\Psi = w - \sum_{i=1,l} x_i p_i$
Elasticities	$q_{\Pi,X}^{ij}$	$\varepsilon_{\{p_k,\forall k\neq i\}},\varepsilon_{\{x_k,\forall k\neq j\}}$
		$e_{\{p_k,\forall k\neq i\}}, e_{\{x_k,\forall k\neq j\}}, \mathfrak{e}_{\{p_k,\forall k\neq i\}}$
Maxwell Relations	$\frac{\varepsilon_{p_1}}{\varepsilon_{p_2}} = \frac{\varepsilon_{x_1}}{\varepsilon_{x_2}}$	
Le Chatelier Principle	$0 > q_{\{x_i, \forall i \neq k\}}^{kk} > \dots$	$0 > \varepsilon_{x_k} > \varepsilon_{p_k}$
Equations of State	$\pi_i(x_1,\ldots,x_n)$	$p_j(x_1,\ldots,x_n)$
Integrability	Hessian matrix is: symmetric	Antonelli's integrability conditions
	and negative semidefinite	Constraints on elasticities or
		economic integrability
Phase transition	$q^{jj}_{\{\pi_i,\forall i\neq j\}} = \infty$	$\varepsilon_{\{p_k,\forall k\neq i\}}=\infty, e_{\{p_k,\forall k\neq i\}}=\infty$
	2	$\mathbf{e}_{\{p_k,\forall k\neq i\}}=\infty$
Maxwell construction	$\pi_k : \int_{\pi_k^1}^{\pi_k^2} x_k d\pi_k = 0$	$\int_{p_k^k}^{p_k^*} x_k dp = 0$

Table 3.7: Table of Correspondences II.

of the entropy in a thermodynamic system out of equilibrium implies that: (1) the system be divided into subsystems where the entropy is defined, i.e., subsystems in equilibrium and (2) the total entropy of the composite system be obtained by summing up the entropies of the subsystems. In contrast, the measurement of the utility out of equilibrium is straightforward because utility is defined for all consumption baskets.

Among the works cited in the introduction on the analogy between equilibrium thermodynamics and neoclassical microeconomics we would like to emphasize the contributions made by Rozonoer [23, 24, 25], Samuelson [27, 26] and Smith and Foley [30]. In this chapter we follow the same approach that is followed in these works considering that the entropy is formally analogous to the utility. Samuelson applies Le Chatelier Principle to economic systems. Rozonoer establishes an analogy between mechanisms of resource exchange and allocation in thermodynamics and economics. This author also obtains a version of the Le Chatelier Principle, although it is less general then the one obtained here. Smith and Foley depart from the three axioms for the mathematical representation of thermodynamic and economic systems (the three laws of thermodynamics). These authors obtain as the second law for the consumer theory a definition of irreversibility in economic behavior similar to the one proposed in this chapter, although our definition of irreversibility is valid only for a constant endowment (constant energy for a thermodynamic system). We believe that this constraint is necessary in the definition of irreversibility because if the consumer is forced to decrease his endowment, e.g., due to a new tax, then he might find optimal to reverse a trade that has increased his utility in the past.

Future work on the topic of equilibrium econophysics should encompass: (1) the derivation of first order phase transitions for the consumer theory with other constraints and between different utility functions, (2) the derivation of second-order phase transitions, (3) the establisment of a relation between consumer theory and social welfare analogous to the relation between statistical physics and macrothermodynamics and (4) the improvement of consumer theory to include thermodynamic irreversibility (building on the work of Sousa and Domingos [31]).

3.13 Appendices

3.13.1 Appendix I

The composite system can have an arbitrary number of subsystems; in this chapter, we consider only the simplified case of two subsystems. Each subsystem is a homogeneous system, i.e., it has a uniform temperature, pressure and chemical potentials for each chemical species. Each subsystem is a simple system.

According to Callen's postulate III [4], the entropy of a composite system is additive over the constituent subsystems, i.e., $S = \sum_{j=1}^{2} S^{i}$, where the entropy of each subsystem is a function of its extensive variables $S^{j}(U^{j}, V^{j}, N_{1}^{j}, \dots, N_{r}^{j})$. Thus, the entropy of the system is $S(U^{1}, U^{2}, V^{1}, V^{2}, N_{1}^{1}, N_{1}^{2}, \dots, N_{r}^{1}, N_{r}^{2})$. Applying the same reasoning to the internal energy, it can be concluded that $U(S^{1}, S^{2}, V^{1}, V^{2}, N_{1}^{1}, N_{1}^{2}, \dots, N_{r}^{1}, N_{r}^{2})$.

The enthalpy of subsystem j is $H^j = U^j + P^j V^j$ and the enthalpy of the composite system is $H = \sum_{j=1}^{2} H^j = \sum_{j=1}^{2} U^j + P^j V^j$. If the intensive variable is the same for both

subsystems, i.e., $P^1 = P^2$ then $H = \sum_{j=1}^2 U^j + PV^j = U + PV$. The same reasoning can be applied to the other Legendre transforms to obtain the expressions given in sec. 3.6.

Elasticities defined in sec. 3.7 can always be used to characterize simple systems. In the case of composite systems: (1) α , c_P and κ_T are adequate for composite systems with homogeneous temperature and pressure, (2) κ_s for composite systems with homogeneous pressure and (3) c_v for composite systems with homogeneous temperature.

Equations of state can describe: (1) composite systems in equilibrium or (2) each subsystem separately.

3.13.2 Appendix II

The Hessian of the Lagrangean function is given by:

$$\begin{bmatrix} \mathcal{L}_{11} & \cdots & \mathcal{L}_{1n} & g_1^1 & \cdots & g_1^m \\ \vdots & \ddots & \vdots & \vdots & \ddots & \vdots \\ \mathcal{L}_{n1} & \cdots & \mathcal{L}_{nn} & g_n^1 & \cdots & g_n^m \\ g_1^1 & \cdots & g_1^m & 0 & \cdots & 0 \\ \vdots & \ddots & \vdots & \vdots & \ddots & \vdots \\ g_n^1 & \cdots & g_n^m & 0 & \cdots & 0 \end{bmatrix}$$

where $\mathcal{L}_{ij} = \frac{\partial^2 \mathcal{L}}{\partial x_j \partial x_i}$ and $g_j^i = \frac{\partial g^i}{\partial x_j}$ [13].

3.13.3 Appendix III

A more detailed deduction of the relation between elasticities given by Eq. 3.32 is presented below. Relations given by Eq. 3.33 to Eq. 3.34 are obtained in a similar manner. Starting from

$$\begin{pmatrix} \frac{\partial x_k}{\partial \pi_k} \end{pmatrix}_{\{\pi_i, \forall i \neq k\}} = \begin{pmatrix} \frac{\partial x_k}{\partial \pi_k} \end{pmatrix}_{x_j, \{\pi_i, \forall i \neq k, j\}} + \\ \begin{pmatrix} \frac{\partial x_k}{\partial x_j} \end{pmatrix}_{\{\pi_i, \forall i \neq j\}} \begin{pmatrix} \frac{\partial x_j}{\partial \pi_k} \end{pmatrix}_{\{\pi_i, \forall i \neq k\}}$$
(3.58)

and (1) multiplying by $\left(\frac{\partial x_j}{\partial \pi_j}\right)_{\{\pi_i,\forall i\neq j\}}$ and (2) using the Maxwell relation $\left(\frac{\partial x_k}{\partial \pi_j}\right)_{\{\pi_i,\forall i\neq j\}} = \left(\frac{\partial x_j}{\partial \pi_k}\right)_{\{\pi_i,\forall i\neq k\}}$ we obtain,

$$\left(\frac{\partial x_j}{\partial \pi_j}\right)_{\{\pi_i,\forall i\neq j\}} \left(\frac{\partial x_k}{\partial \pi_k}\right)_{\{\pi_i,\forall i\neq k\}} =$$

Equilibrium Econophysics

$$= \left(\frac{\partial x_j}{\partial \pi_j}\right)_{\{\pi_i,\forall i\neq j\}} \left(\frac{\partial x_k}{\partial \pi_k}\right)_{x_j,\{\pi_i,\forall i\neq k,j\}} + \left(\frac{\partial x_j}{\partial \pi_k}\right)_{\{\pi_i,\forall i\neq k\}} \left(\frac{\partial x_j}{\partial \pi_k}\right)_{\{\pi_i,\forall i\neq k\}}.$$
(3.59)

Eq. 3.32 is obtained by multiplying Eq. 3.59 by $\frac{\pi_k}{x_k}$ and rearranging.

The deduction of Eq. 3.35 is presented next. The differential of the extensive variable x_k can be written as:

$$dx_k = \frac{\partial x_k}{\partial \pi_1} d\pi_1 + \ldots + \frac{\partial x_k}{\partial \pi_n} d\pi_n.$$
(3.60)

Starting with Eq.3.60 the following steps are taken: (1) the Maxwell relation, $\left(\frac{\partial x_i}{\partial \pi_k}\right)_{\{\pi_j, \forall j \neq k\}} = \left(\frac{\partial x_k}{\partial \pi_i}\right)_{\{\pi_j, \forall j \neq i\}}$ (Tab 3.2) is used to replace the partial derivatives, (2) the expression is differentiated in relation to π_k with x_i constant and (3) the expression is multiplied by $\frac{\pi_k}{x_k}$ and the following equation is obtained:

$$\frac{\pi_k}{x_k} \left(\frac{\partial x_k}{\partial \pi_k}\right)_{\{x_i,\forall i\neq k\}} = \frac{\pi_k}{x_k} \left(\frac{\partial x_1}{\partial \pi_k}\right)_{\{\pi_i,\forall i\neq k\}} \left(\frac{\partial \pi_1}{\partial \pi_k}\right)_{\{x_i,\forall i\neq k\}} + \dots + \frac{\pi_k}{x_k} \left(\frac{\partial x_n}{\partial \pi_k}\right)_{\{\pi_i,\forall i\neq k\}} \left(\frac{\partial \pi_n}{\partial \pi_k}\right)_{\{x_i,\forall i\neq k\}}.$$
(3.61)

Using the elasticity given by Eq. 3.27, Eq. 3.61 can be written:

$$q_{\{x_i,\forall i\neq k\}}^{kk} = \frac{x_1}{x_k} q_{\{\pi_i,\forall i\neq k\}}^{1k} \left(\frac{\partial \pi_1}{\partial \pi_k}\right)_{\{x_i,\forall i\neq k\}} + \dots + \frac{x_n}{x_k} q_{\{\pi_i,\forall i\neq k\}}^{nk} \left(\frac{\partial \pi_n}{\partial \pi_k}\right)_{\{x_i,\forall i\neq k\}}.$$
(3.62)

With the partial derivatives replaced by

$$\left(\frac{\partial \pi_j}{\partial \pi_k}\right)_{x_i \neq k} = -\frac{\pi_j q_{\pi_j, \{x_i, \forall i \neq k, j\}}^{jk}}{\pi_k q_{\pi_k, \{x_i, \forall i \neq k, j\}}^{jj}},$$
(3.63)

obtained using Eq. 3.27, Eq. 3.62 simplifies to Eq. 3.35.

3.13.4 Appendix IV

A simple and direct demonstration of the Euler Equation is presented by Callen [4, pp. 59], where Equation $y(\lambda x_1, \ldots, \lambda x_n) = \lambda y(x_1, \ldots, x_n)$ is differentiated with respect to λ and then λ is set to 1.

The Gibbs-Duhem Equation, $0 = x_1 d\pi_1 + \ldots + x_n d\pi_m$, is obtained by combining the differential of the Euler Equation, $dy = \pi_1 dx_1 + \ldots + \pi_m dx_n + x_1 d\pi_1 + \ldots + x_n d\pi_m$, with the fundamental equation $dy = \pi_1 dx_1 + \ldots + \pi_m dx_n$.

References

- S. A. Amel'kin, K. Martinás, and A. M. Tsirlin. Optimal control for irreversible processes in thermodynamics and microeconomics. *Automation and Remote Control*, 63:519–539, 2002.
- [2] R. S. Berry, V. A. Kasakov, S. Sieniutycz, Z. Szwast, and A. M. Tsirlin. *Thermodynamic Optimization of Finite-Time Processes*. Wiley, Chichester, 2000.
- [3] S. Bornholdt and Wagner F. Stability of money: phase transitions in an Ising economy. *Physica A*, 316:453–468, 2002.
- [4] H. Callen. *Thermodynamics and an Introduction to Thermostatistics*. John Wiley & Sons, New York, NY, 1985.
- [5] J. C. Candeal, J. R. De Miguel, E. Indurain, and G. B. Mehta. On a theorem of Cooper. *Journal of Mathematical Analysis and Applications*, 258:701–710, 2001.
- [6] J. C. Candeal, J. R. De Miguel, E. Indurain, and G. B. Mehta. Utility and entropy. *Economic Theory*, 17:233–238, 2001.
- [7] J. L. B. Cooper. The foundations of thermodynamics. *Journal of Mathematical Analysis and Applications*, 17:172–193, 1967.
- [8] M. G. Daniels, J. D. Farmer, L. Gillemot, G. Iori, and Smith E. Quantitaive model of price diffusion and market friction based on trading as a mechanistic random process. *Physical Review Letters*, 90:108102, 2003.
- [9] H. T. Davis. The Theory of Econometrics. Principia Press, Bloomington, IN, 1941.
- [10] A. Dragulescu and V. M. Yakovenko. Exponential and power-law probability distributions of wealth and income in the United Kingdom and the United States. *Physica* A, 299:213–221, 2001.
- [11] J. D. Farmer, P. Patelli, and I. I. Zovko. The predictive power of zero intelligence in financial markets. *Proceedings of the National Academy of Sciences*, 102:2254– 2259, 2005.
- [12] D. Grech. Econophysics teaching at the University of Wroclaw. *Physica A*, 344:335–339, 2004.
- [13] G. Jehle. Advanced Microeconomic Theory. Prentice-Hall, Englewood Cliffs, NJ, 1991.
- [14] L. D. Landau and E. M. Lifshitz. *Statistical Physics*. Pergamon Press, London, 1959.
- [15] F. Lillo, J. D. Farmer, and R. N. Mantegna. Master curve for price-impact function. *Nature*, 421:129–130, 2003.
- [16] J. H. C. Lisman. Econometrics and thermodynamics: A remark on Davis' theory of budgets. *Econometrica*, 17:59–62, 1949.
- [17] A. Martina. Antonelli's analytical techniques: their exploitation to derive results in duality theory. *European Journal History of Economic Thought*, 7:363–376, 2000.
- [18] A. Mas-Collel, M. Whinston, and J. R. Green. *Microeconomic Theory*. Oxford University Press, New York, NY, 1995.
- [19] J. L. McCauley. Thermodynamic analogies in economics and finance: instability of markets. *Physica A*, 329:199–212, 2003.
- [20] A. Munro and R. Sugden. On the theory of reference-dependent preferences. *Journal of Economic Behavior and Organization*, 50:407–428, 2003.
- [21] V. Plerou, P. Gopikrishnan, and H. E. Stanley. Two-phase behaviour of financial markets. *Nature*, 421:130–130, 2003.
- [22] R. T. Rockafeller. Convex Analysis. Princeton University Press, Princeton, NJ, 1970.
- [23] L. I. Rozonoer. A generalized thermodynamic approach to resource exchange and allocation II. Automatation and Remote Control, 34:915–927, 1973.
- [24] L. I. Rozonoer. Large scale systems and operations management. a generalized thermodynamic approach to resource exchange and allocation I. *Automatation and Remote Control*, 34:781–795, 1973.

- [25] L. I. Rozonoer. A generalized thermodynamic approach to resource exchange and allocation III. Automatation and Remote Control, 34:1272–1289, 1977.
- [26] P. A. Samuelson. An extension of the Le Chatelier principle. *Econometrica*, 28:368– 379, 1960.
- [27] P. A. Samuelson. Structure of a Minimum Equilibrium System. In: Ralph W. Pfouts, ed., Essays in economics and econometrics: a volume in honor of Harold Hotelling. University of North Carolina Press, 1960.
- [28] W. M. Saslow. An economic analogy to thermodynamics. American Journal of Physics, 67:1239–1247, 1999.
- [29] E. Smith, J. D. Farmer, L. Gillemot, and S. Krishnamurthy. Statistical theory of the continuous double auction. *Quantitative Finance*, 3:481–514, 2003.
- [30] E. Smith and D. Foley. Classical thermodynamics and economic general equilibrium theory. *New School University, Center for Economic Policy Analysis Working Papers*, 2004.
- [31] T. Sousa and T. Domingos. Is neoclassical microeconomics formally valid? An approach based on an analogy with equilibrium thermodynamics. *Ecological Economics*, 58:160–169, 2006.
- [32] H. E. Stanley, L.A.N. Amaral, D. Canning, P. Gopikrishnan, Y. Lee, and Y. Liu. Econophysics: Can physicists contribute to the science of economics? *Physica A*, 269:156–169, 1999.
- [33] L. Tisza. Generalized Thermodynamics. M.I.T Press, Cambridge, MA, 1966.
- [34] A. M. Tsirlin and S. A. Amel'kin. Dissipation and conditions of equilibrium for an open microeconomic system. *Open Systems and Information Dynamics*, 8:157–168, 2001.
- [35] A. M. Tsirlin, V. Kasakov, and N. A. Kolinko. Irreversibility and limiting possibilities of macrocontrolled systems: II. Microeconomics. *Open Systems and Information Dynamics*, 8:329–347, 2001.
- [36] N. Vandewalle and M. Ausloos. Coherent and random sequences in financial fluctuations. *Physica A*, 246:454–459, 1997.

- [37] N. Vandewalle, Ph. Boveroux, A. Minguet, and M. Ausloos. The crash of October 1987 seen as a phase transition: amplitude and universality. *Physica A*, 255:201–210, 1998.
- [38] H. Varian. Intermediate Microeconomics. A Modern Approach. Norton, New York, NY, 1996.
- [39] K. Yamasaki, L. Muchnik, S. Havlin, A. Bunde, and H. E. Stanley. Scaling and memory in volatility return intervals in financial markets. *Proceedings of the National Academy of Sciences*, 102:9424–9428, 2005.

Chapter 3

64

Chapter 4

The thermodynamics of organisms in the context of DEB theory

Abstract

We carry out a thermodynamic analysis to an organism. It is applicable to any type of organism because 1) it is based on a thermodynamic formalism applicable to all open thermodynamic systems and 2) uses a general model to describe the internal structure of the organism - the Dynamic Energy Budget Model. Our results on the thermodynamics of DEB organisms are the following. (1) Thermodynamic constraints for the following types of organisms: (a) aerobic and exothermic, (b) anaerobic and exothermic and (c) anaerobic and endothermic; showing that anaerobic organisms have a higher thermodynamic flexibility. (2) A way to compute the changes in the enthalpy and in the entropy of living biomass that accompany changes in growth rate solving the problem of evaluating the thermodynamic properties of biomass as a function of the amount of reserves. (3) Two expressions for Thornton's coefficient that explain its experimental variability and theoretically underpin its use in metabolic studies. (4) A new mechanism that organisms in non steady-state use to rid themselves of internal entropy production: 'dilution of entropy production by growth'. To demonstrate the practical applicability of DEB theory to quantify thermodynamic changes in organisms we use published data on Klebsiella aerogenes growing aerobically in a continuous culture. We obtain different values for molar entropies of the reserve and the structure of Klebsiella aerogenes proving that the reserve density concept of DEB theory is essential in discussions concerning (a) the relationship between organization and entropy and (b) the mechanism of storing entropy in new biomass. Additionally, our results suggest that the entropy of dead biomass is significantly

different from the entropy of living biomass.

Keywords: Organism, Thermodynamics, Dynamic Energy Budget Theory, Entropy, Enthalpy, Thornton's coefficient.

4.1 Introduction

Many thermodynamic analyses of living organisms and cells have been made. Demirel et al.[10], for example, used thermodynamics to describe the coupled phenomena of transport and chemical reactions that take place inside living systems; Battley analyzed the entropy change accompanying the growth of *E. coli* [2] and the growth of *Saccharomyces cerevisiae* [5]; Stockar et al. [31, 32] analyzed the internal entropy production in some micro-organisms; Esener et al. [12] studied the energetics of *Klebsiella aerogenes*; Duboc et al. [11] evaluated the thermodynamic efficiency of *Saccharomyces cerevisae*; Qian and Beard [29] introduced a thermodynamic formalism to study metabolic biochemical reaction networks, etc.

The thermodynamic analyses mentioned use equations that describe chemical reactions that take place inside an organism as the model describing the organism's behavior; examples of these aggregated chemical reactions are given in Battley[5, Table 2] for some microorganisms. Because these models need a considerable amount of data they are not useful in thermodynamic analyses made of more complex organisms. Even for microorganisms these models present problems because they do not give a mechanistic explanation of many of the energetic aspects of an organism's growth process; this is patent in the ad-hoc explanations given for the empirical results.

The thermodynamic analyses that have been made would benefit from a general theory underlying the description of the energetic fluxes in order to build up solid knowledge about organism's metabolism. The Dynamics Energy Budget (DEB) theory is the most general non-species-specific theory of this kind [22, 23, 28]. It consists of a set of simple, mechanistically inspired rules that fully specify the uptake and use of mass and energy by an organism. The frequently applied classical models by Monod and Marr-Pirt on bacterial growth, and the well-known model by Droop for nutrient limited growth of unicellular algae are all special cases of DEB theory. DEB theory also considers phenomena of a complexity well beyond these simple models, including simultaneous nutrient limitation, adaptation, co-metabolism, flocculated growth, product formation, aging and syntrophy.

Here, we will carry out a thermodynamic analysis of an organism using 1) the most

general framework of non-equilibrium thermodynamics applicable to all open thermodynamic systems (de Groot and Mazur [19], Bejan [7] and Moran et al. [26]) and 2) the knowledge of its internal dynamics given by DEB theory.

This chapter is organized as follows. In section 4.2 we begin by defining the DEB thermodynamic system, i.e., the mass and energy flows in the organism and the dynamics of its state variables. This is followed by a thermodynamic analysis of the organism in section 4.3. This analysis uses the mass, energy and entropy balances together with DEB to obtain the thermodynamic constraints imposed on different types of organisms and to discuss the use of Thornton's coefficient. In Section 4.4 a thermodynamic characterization of *Klebsiella aerogenes* is done using published data from chemostat experiments. Section 4.5 concludes.

4.2 Thermodynamic System

To simplify our analysis, we focus on a single-reserve, single-structure heterotrophic that lives on a single substrate. It is a V1 morph, i.e., an organism whose surface area is proportional to volume. Extensions to include more substrates, reserves and/or structures have been formulated in the DEB theory to include autotrophs and nutritional details [22]. However, such extensions do not pose new thermodynamic problems. Appendix 4.6.1 presents an explanation of the notation used throughout the chapter.

Fig. 5.1 presents a scheme, according to DEB theory, for an asexually propagating heterotrophic organism ¹. The simplest DEB model delineates one reserve, E, with M_E C-moles, and one structure, V, with M_V C-moles; the chemical composition of each is assumed to be constant (the strong homeostasis assumption). This simplifying assumption combines nicely with the idea that the enzymes that catalyse chemical transformations in the organism do so in a constant chemical environment, which is essential for full control. Reserve has the dual role of providing both energy and essential compounds for growth, maintenance, development and reproduction.

The incoming and outgoing fluxes through the outer surface of the system can be divided into organic (food, X and products, P), and mineral (CO_2 , H_2O , O_2 , and nitrogenous waste, N_{waste}) compounds. The energy flow of the food uptake per C-mol of structure \dot{p}_X , enters and is transformed into an assimilation energy flux per C-mol of structure, \dot{p}_A , that goes into the reserves and into products, P, that are excreted. The catabolic

¹The simplifications chosen are suitable for our empirical application.

energy flux, \dot{p}_C , i.e., the rate per C-mol of structure, at which the organism mobilizes its reserve for metabolic purposes ², is partitioned between growth, \dot{p}_G , and maintenance, \dot{p}_M , i.e.,

$$\dot{p}_C = \dot{p}_M + \dot{p}_G. \tag{4.1}$$

Maintenance includes a variety of requirements, such as the turnover of chemical compounds of structure (e.g. proteins), the maintenance of concentration gradients across membranes, the maintenance of defense systems (e.g. the immune system), activity (including behavior), the heating of the body to a near constant temperature (endotherms only) and osmotic work (especially freshwater organisms) [22]. Growth is defined as the increase of structure; body weight has contributions from both reserve and structure.

Ellipses in Fig. 5.1 are idealized reactors where anabolic and catabolic processes take place. These transformations occur all over the organism, rather than at specific sites. Consequently, the idealized reactors are the transformations themselves. These processes are characterized by constant conversion efficiencies between mass flows, y_{*1*2} , i.e., the number of moles of *1 needed to produce one mol of *2. In the assimilation reactor food is converted into reserve, y_{XE} , and in the growth reactor reserve is converted into structure, y_{EV} .

The ratio $m_E \equiv M_E/M_V$ is the reserve density. For any constant food level, $X = X^* > 0$, there is a reserve density, m_E^* , that remains constant along the growth process. Furthermore $\lim_{X^*\to\infty} m_E^* \equiv m_{Em}$ where m_{Em} is the maximum reserve density (the weak homeostasis assumption).

We now quantify the dynamics of the state variables structure and reserve. The amount of structure and reserve and the flows in the organism are measured in mass and Gibbs energy and the conversion between them is done using chemical potentials: the chemical potential of food, μ_X , converts the flow of food per C-mol of structure, j_X , to \dot{p}_X ; the chemical potential of reserve, μ_E , converts the flow of reserve that exits the assimilation reactor to \dot{p}_A and the flow of reserve that exits the reserve compartment to $\dot{p}_C = \dot{p}_M + \dot{p}_G$.

The change in C-moles of reserve per C-mol of structure is:

$$j_E \equiv \frac{1}{M_V} \frac{dM_E}{dt} = \frac{\dot{p}_A - \dot{p}_C}{\mu_E},$$
(4.2)

because the reserve is continuously used (catabolic power) and replenished (assimilation power). With Eq. 4.2, the reserve density dynamics is

$$\frac{dm_E}{dt} \equiv \frac{d(M_E/M_V)}{dt} = \frac{\dot{p}_A - \dot{p}_C}{\mu_E} - m_E \frac{1}{M_V} \frac{dM_V}{dt},$$
(4.3)

²Catabolism has a somewhat different meaning in the biochemistry literature.

where the last term on the right hand side is the dilution by growth.

To further evaluate the reserve dynamics we need to quantify feeding and assimilation. The food uptake per C-mol of structure is given by $j_X = f j_{Xm}$ where f is the scaled functional response and j_{Xm} is the maximum food uptake per C-mol of structure, i.e., the food uptake that would occur at abundant food availability. The assimilation energy flow per C-mol of structure is

$$\dot{p}_A = \frac{j_X}{y_{XE}} \mu_E,\tag{4.4}$$

where $\frac{j_X}{y_{XE}}$ is the reserve flow that exits the assimilation reactor per C-mol of structure.

The reserve dynamics can be further deduced considering 1) that the catabolic power is independent of the food availability, 2) that the mobilization of reserves does not depend on how they are partitioned among aggregates with different chemical compositions and 3) the weak homeostasis assumption (for details on the derivation see Appendix 4.6.2). The reserve dynamics is then

$$\frac{dm_E}{dt} = \dot{k}_E \left(fm_{Em} - m_E \right), \tag{4.5}$$

where \dot{k}_E is the maximum reserve turnover rate:

$$\dot{k}_E = \frac{j_{Xm}/y_{XE}}{m_{Em}}.$$
 (4.6)

The parameter k_E is a turnover rate because it is the ratio between the maximum flow of C-moles of reserve into the 'reserve compartment', $j_{Xm}/y_{XE}M_V$, and the maximum amount of C-moles of reserve kept in the 'reserve compartment', $m_{Em}M_V$.

By combining Eq. 4.3 with Eq. 4.5 the catabolic power simplifies to:

$$\dot{p}_C = \mu_E m_E \left(\dot{k}_E - \frac{1}{M_V} \frac{dM_V}{dt} \right). \tag{4.7}$$

To evaluate the dynamics of the structure we need to quantify maintenance and growth. The maintenance energy flow per C-mol of structure is

$$\dot{p}_M = \dot{k}_M y_{EV} \mu_E, \tag{4.8}$$

where $\dot{k}_M y_{EV}$ is the reserve flow that exits the reserve for maintenance purposes per Cmol of structure and \dot{k}_M is the maintenance rate coefficient. The growth energy flow per C-mol of structure is

$$\dot{p}_G = j_V y_{EV} \mu_E, \tag{4.9}$$



Figure 4.1: Schematization of an asexually propagating heterotrophic organism following DEB theory. Boxes represent boundaries. Ovals represent chemical reactors. Lines represent mass and energy flows: \dot{p}_X has the chemical composition of food; \dot{p}_A , \dot{p}_C , \dot{p}_G and \dot{p}_M have the chemical composition of reserve and the flow that goes from the growth reactor to the structure compartment has the chemical composition of structure.

	Compound
X	Food
P	Products
E	Reserve
V	Structure
W	Biomass
\mathcal{O}	class of organic compounds (X, P, E, V)
\mathcal{M}	class of mineral compounds $(CO_2, H_2O, O_2, N_{waste})$

Table 4.1: List of compounds.

where $j_V y_{EV}$ is the flow that exits the reserve for growth purposes per C-mol of structure and j_V is the molar flow of structure.

The dynamics of structure

$$j_V \equiv \frac{1}{M_V} \frac{dM_V}{dt} = \frac{\dot{p}_G}{y_{EV}\mu_E} = \frac{(\dot{k}_E \frac{m_E}{m_{Em}} - \dot{k}_M g)}{(\frac{m_E}{m_{Em}} + g)},$$
(4.10)

is obtained by combining Eq. 4.1, Eq. 4.8, Eq. 4.9, and Eq. 4.7, where the investment ratio, *g*, is a dimensionless quantity given by

$$g \equiv y_{EV}/m_{Em},\tag{4.11}$$

that stands for the ratio between the number of moles of reserve allocated to growth per mole of structure and the maximum number of moles of reserve allocated to maintenance plus growth per mole of structure. At constant food the dynamics of structure simplifies to:

$$j_V = (\dot{k}_E f - \dot{k}_M g) / (f + g),$$
(4.12)

because the weak homeostasis assumption implies that $\frac{dm_E}{dt} = 0$.

4.3 Thermodynamic Analysis

We now formalise the thermodynamic analysis. See Table 4.1 for a list of compounds.

If the mole numbers of each compound, the total energy and the total entropy of the organism are constant, the organism is in steady state. In this state there is still a continuous flow of matter, energy and entropy through the system from and to external reservoirs. This is implied by maintenance requirements. Here, we will consider that the organism is not in steady state because our life cycle perspective of an individual comes with the necessity to consider changes in amounts of reserve and structure, both in mass and energy aspects.

The strong homeostasis assumption justifies one of the main simplifications used in the thermodynamic analysis: the molar chemical compositions, internal energies, enthalpies and entropies of reserve and structure are taken to be constant, independent of the reserve density. In general, we consider that the chemical composition of biomass (structure plus reserve) can change; this implies that the molar thermodynamic properties of biomass can also change. Whenever the reserve density is constant, the chemical composition of biomass and its molar thermodynamic properties are also constant.

4.3.1 Mass Balance Equation

Most of the dry mass of biological systems consists of proteins, lipids, carbohydrates and nucleic acids. The major chemical elements in the covalently bounded compounds are carbon, C, hydrogen, H, oxygen, O, nitrogen, N, phosphorus and sulphur. The first four elements stand for more than 96% of the total dry mass [27], so we focus on these elements only. A 'molecule' of structure is denoted by $CH_{n_{HV}}O_{n_{OV}}N_{n_{NV}}$, a 'molecule' of reserve by $CH_{n_{HE}}O_{n_{OE}}N_{n_{NE}}$, where the chemical index n_{ij} is the number of atoms of element *i* per atom of carbon in compound *j*. The mass balance equation for the organism is written on a molar basis for each element because there is no conservation of compounds due to the chemical transformations inside the organism. The mass balance equation is:

$$0 = \boldsymbol{n}_{\mathcal{M}} \dot{\boldsymbol{J}}_{\mathcal{M}} + \boldsymbol{n}_{\mathcal{O}} \dot{\boldsymbol{J}}_{\mathcal{O}}$$
(4.13)

where $\dot{J}_{\mathcal{M}}$ is the vector with the molar fluxes of the minerals $(\dot{J}_{CO_2}, \dot{J}_{H_2O}, \dot{J}_{O_2}, \dot{J}_{N_{waste}})$, $\dot{J}_{\mathcal{O}}$ is the vector with the molar fluxes of the organics $(\dot{J}_X, \dot{J}_P, -\dot{J}_V, -\dot{J}_E)$, $n_{\mathcal{M}}$ is the matrix with the chemical composition of minerals and $n_{\mathcal{O}}$ is the matrix with the chemical composition of organic compounds. Each entry in these matrices, n_{*1*2} , is the number of atoms of element *1 in compound *2.

The fluxes $\dot{J}_V \equiv \frac{d}{dt}M_V$ and $\dot{J}_E \equiv \frac{d}{dt}M_E$ are the change in C-mols of structure and reserve per unit time in the organism. The other fluxes, \dot{J}_j , mole numbers per unit time, are positive if they represent a net input into the thermodynamically defined organism and negative otherwise. In a heterotrophic organism CO_2 is usually but not always an output, O_2 is an input, H_2O is an output formed metabolically from other compounds,

nitrogenous waste is an output, food is an input and products are an output. Eq. (4.13) states that, for each element, the rate of mole numbers accumulation inside the organism (structure and reserve), equals the inputs minus the outputs (other organic and mineral flows).

4.3.2 Energy Balance Equation

The energy balance equation quantifies the organism's accumulation of energy as the result of inputs minus outputs of energy fluxes. The energy fluxes (Joule/s) are the net heat flux, the net work flux and energy fluxes associated with input and output molar fluxes.

In this analysis mechanical work will be considered negligible. This is supported by Garby et al. [17], who state that 'the energy transfer as heat (in animals) is relatively large and directed outwards, while the energy transfer as work is small'.

The temperature of structure, reserve and outgoing products, T, is assumed constant and equal to the temperature of the body. We think this is a reasonable first approximation because chemical reactions inside the organism occur for a limited temperature range mainly due to enzymatic action [22]. Anyway this will probably be a better approximation for endotherms or for ectotherms in an environment where they are able to keep their temperature constant by moving.

This set of assumptions, i.e., a constant temperature of the organism, negligible mechanical work and incoming fluxes with a temperature similar to the organism's imply that the net heat released by the organism equals the net heat produced in all chemical reactions inside the organism. Chemical reactions taking place are the degradation of food and reserve material in order to obtain energy (synthesize ATP from ADP) and release nutrients, and the building up of reserve and new structural material with the nutrients and energy obtained.

Here, we apply the energy balance equation to aerobic and to anaerobic organisms. The distinction between these types of organisms is useful because there are simplifications applicable only to aerobic organisms (sections 4.3.2 and 4.3.3).

Energy Balance Equation: General Organism

With the simplifications that were mentioned in section 4.3.2, the energy balance of the entire thermodynamic system is

$$0 = \overline{\boldsymbol{h}}_{\mathcal{M}}^{T} \dot{\boldsymbol{J}}_{\mathcal{M}} + \overline{\boldsymbol{h}}_{\mathcal{O}}^{T} \dot{\boldsymbol{J}}_{\mathcal{O}} + \dot{P}_{T+}$$
(4.14)

where \dot{P}_{T+} is the total released heat ³, $\overline{\boldsymbol{h}}_{\mathcal{M}}^{T} = \left(\overline{h}_{CO_2} \quad \overline{h}_{H_2O} \quad \overline{h}_{O_2} \quad \overline{h}_{N_{waste}} \right)^{T}$ and $\overline{\boldsymbol{h}}_{\mathcal{O}}^{T} = \left(\overline{h}_X \quad \overline{h}_P \quad \overline{h}_V \quad \overline{h}_E \right)^{T}$ and \overline{h}_i is the molar enthalpy of compound *i*. The internal energy and the flow work linked to the input and output molar fluxes are lumped in the enthalpy.

Eq. 4.14 is the energy balance for a non-steady state organism defined according to DEB theory. This supports the use of direct calorimetry (the direct measurement of released heat) to assess enthalpy changes in organisms as referred by [31]: if the organism was completely burned then the net heat release plus the enthalpy of the combustion products would be equal to the organism's total enthalpy.

The enthalpy can be substituted by $\overline{h} = \overline{u} + P\overline{v} = \overline{g} + T\overline{s} = \mu + T\overline{s}$, which is obtained using the definition of Gibbs energy $\overline{g} = \overline{u} - T\overline{s} + P\overline{v}$ and the equality between Gibbs energy and chemical potential $g = \mu$ for a single component:

$$0 = (\boldsymbol{\mu}_{\mathcal{M}} + T\overline{\boldsymbol{s}}_{\mathcal{M}})^T \dot{\boldsymbol{J}}_{\mathcal{M}} + (\boldsymbol{\mu}_{\mathcal{O}} + T\overline{\boldsymbol{s}}_{\mathcal{O}})^T \dot{\boldsymbol{J}}_{\mathcal{O}} + \dot{P}_{T+}$$
(4.15)

where \overline{s}_E is the molar entropy of the reserve, \overline{s}_V is the molar entropy of the structure, $\overline{\mu}_M$ and \overline{s}_M collect the values for the four minerals; $\overline{\mu}_O$ and \overline{s}_O do that for the organic compounds, as before.

Energy Balance Equation: Aerobic Organism

We assume that reactors have negligible mass and are at pseudo steady state. An additional assumption that can also be made is based on Garby and Larsen [17] who consider, based on empirical knowledge, that for most important reactions in biological systems $T\Delta s$ is very small compared to Δh and therefore the enthalpy of the reaction $\Delta h_{reaction}$ is approximated using its Gibbs energy $\Delta g_{reaction}$, since at constant temperature $\Delta g = \Delta h - T\Delta s \simeq \Delta h$. This assumption is valid only for aerobic reactions but it is less stringent than that of Kooijman [22], where entropy is set to zero.

³The heat follows the same sign convention as the mass flows: it is negative if the net input is negative, i.e., the heat released is negative

We apply the simplification that $T\Delta s$ is very small to the set of all reactors to obtain:

$$0 = \overline{\mathbf{s}}_{\mathcal{M}}^T \dot{\mathbf{J}}_{\mathcal{M}} + \overline{\mathbf{s}}_{\mathcal{O}}^T \dot{\mathbf{J}}_{\mathcal{O}}.$$
(4.16)

This equation can be disaggregated to

$$\overline{s}_V \dot{J}_V + \overline{s}_E \dot{J}_E = \overline{s}_{\mathcal{M}}^T \dot{J}_{\mathcal{M}} + \overline{s}_X \dot{J}_X + \overline{s}_P \dot{J}_P, \qquad (4.17)$$

to emphasize that the entropy variation of an aerobic organism (left hand side), equals the net import of chemical entropy given by the right hand side of Eq. 4.17⁴. With Eq. (4.17) we conclude that, for an aerobic organism, if weak homeostasis applies, i.e., reserve and structure are in constant proportions, then there is a positive net import of chemical entropy during an increase in biomass, a negative net import of chemical entropy during a decrease in biomass and a null net import at constant biomass (see the entries (1,1), (3,1) and (5,1)⁵ in Table 4.2). If weak homeostasis does not apply and the structure and reserve molar entropies are different then an organism that is increasing its biomass could either have a positive or negative net import of chemical entropy. This is in accordance with Stockar and Liu [31] who concluded that microbial growth might result either from a positive or negative net import of chemical entropy.

The total entropy of an organism is easily computed if the molar entropy values and the chemical composition of biomass are known. Otherwise, Eq. 4.16 has to be integrated to obtain the total entropy of an organism as a function of total inputs and outputs from birth t_b until time t.

We apply the energy balance equation to the set of all reactors assuming that the reactors are in steady-state but that the organism as a whole is not. We obtain 6 ,

$$0 = \dot{Q}_{reactions} + \overline{\mu}_{\mathcal{O}}^T \dot{J}_{\mathcal{O}} + \overline{\mu}_{\mathcal{M}}^T \dot{J}_{\mathcal{M}}$$
(4.18)

where $\dot{Q}_{reactions}$ is the rate of the total heat release by all chemical reactions, since the work is null and the accumulation of energy is also null. Eq. 4.18 is equivalent to the energy balance equation presented in [22, p153] but it is applicable only to an aerobic organism. It would only be applicable to other organisms if entropies were set equal to zero in Eq. 4.15. This is not a reasonable assumption because it is equivalent to assuming

⁴Chemical entropy is the entropy associated with mass.

⁵Entry is written as (line,column).

⁶We obtain this equation by summing the equations that describe the aggregate chemical reactions occurring in each reactor.

that the heat capacities of the various compounds including the minerals are null in the interval between the absolute temperatures 0 K and T. Eq. 4.18 can be disaggregrated to

$$\overline{\mu}_V \dot{J}_V + \overline{\mu}_E \dot{J}_E = \dot{Q}_{reactions} + \overline{\mu}_X \dot{J}_X + \overline{\mu}_P \dot{J}_P + \overline{\mu}_M^T \dot{J}_M.$$
(4.19)

4.3.3 Entropy Balance Equation

The entropy balance equation states that the change in entropy is equal to the entropy production inside the organism due to irreversible processes plus the net entropy flux associated with heat and mass fluxes. The entropy balance equation is different from the other balance equations because the accumulation of entropy inside the organism depends on the internal processes, which control the value of the entropy production.

Entropy Balance Equation: General Organism

The entropy balance for the organism is

$$0 = \dot{\sigma} + \frac{\dot{P}_{T+}}{T} + \overline{s}_{\mathcal{M}}^T \dot{J}_{\mathcal{M}} + \overline{s}_{\mathcal{O}}^T \dot{J}_{\mathcal{O}}, \qquad (4.20)$$

which can be written as

$$\overline{s}_V \dot{J}_V + \overline{s}_E \dot{J}_E = \dot{\sigma} + \frac{P_{T+}}{T} + \overline{s}_M^T \dot{J}_M + \overline{s}_X \dot{J}_X + \overline{s}_P \dot{J}_P, \qquad (4.21)$$

where \dot{P}_{T+}/T is the entropy exchange coupled with heat fluxes.

With Eq. 4.21 we conclude that for an anaerobic and exothermic organism ⁷ at constant biomass, if weak homeostasis applies then the organism can have a 1) positive or 2) negative net import of chemical entropy (see the entries (1,2) and (1,3) in Table 4.2). In case 1 (resp. case 2) $T\dot{\sigma} + \dot{P}_{T+} < 0$ (resp. $T\dot{\sigma} + \dot{P}_{T+} > 0$). In case 1 (resp. case 2), the organism will have a positive (resp. positive or negative) net import of chemical entropy when its biomass is increasing and a positive or negative (resp. negative) net import of chemical entropy when its biomass is decreasing (see the entries (3,2), (3,3) and (5,2), (5,3) in Table 4.2). For an aerobic and endothermic organism the net import of chemical entropy follows the same behaviour as in case 2 because $T\dot{\sigma} + \dot{P}_{T+} > 0$ (see the entries (1,4), (3,4) and (5,4) in Table 4.2).

Multiplying Eq. 4.20 by T and subtracting Eq. 4.15 we obtain

$$0 = \overline{\boldsymbol{g}}_{\mathcal{M}}^{T} \dot{\boldsymbol{J}}_{\mathcal{M}} + \overline{\boldsymbol{g}}_{\mathcal{O}}^{T} \dot{\boldsymbol{J}}_{\mathcal{O}} - \dot{\sigma}T, \qquad (4.22)$$

⁷Exothermic (endothermic) organisms are organisms whose overall metabolism results in a negative (positive) net import of heat from the exterior.

where \overline{g}_i is the molar Gibbs energy per mole of compound *i*. This equation can be disaggregated to

$$\overline{g}_V \dot{J}_V + \overline{g}_E \dot{J}_E + \dot{\sigma}T = \overline{g}_{\mathcal{M}}^T \dot{J}_{\mathcal{M}} + \overline{g}_X \dot{J}_X + \overline{g}_P \dot{J}_P.$$
(4.23)

to emphasize that the net import of Gibbs energy equals the rate of Gibbs energy change inside the organism plus an irreversibility production measure.

The net import of Gibbs energy by an anaerobic organism follows the same behaviour for exothermic and endothermic organisms (see Eq. 4.23). When the organism is either increasing its biomass or at constant biomass the net input of Gibbs energy is positive (see the entries (2,2), (2,3), (2,4), (4,2), (4,3) and (4,4)). When the organism is decreasing its biomass the net input of Gibbs energy is negative or positive (see the entries (6,2), (6,3) and (6,4)).

Entropy Balance Equation: Aerobic Organism

The results of the previous section can be further narrowed for aerobic organisms. Subtracting Eq. 4.16 from Eq. 4.20, we obtain

$$\dot{\sigma} = -\frac{\dot{P}_{T+}}{T},\tag{4.24}$$

i.e., the rate of heat released by the aerobic organism equals minus an irreversibility production measure.

Therefore, since the second law tells us that entropy production is always positive, the total heat obtained from the organism is negative (released), which means that the sum of the processes of assimilation, dissipation and growth must be exothermic for aerobic life. The result of Kooijman in [22], i.e., 'the second law of thermodynamics implies that [each of] the processes of assimilation, dissipation and growth is exothermic', is obtained only by considering that entropies are null.

The second law only forbids that processes as a whole are endothermic when the organism is aerobic and the heat released by the organism is equal to the heat released in all chemical reactions. For example, using direct calorimetry, Stockar et al. [32, 25] showed the existence of a chemotroph whose overall metabolic process (assimilation plus dissipation plus growth) is endothermic. This rare type of overall metabolic process is called 'enthalpy retarded growth'.

With Eq. 4.19 and Eq. 4.24 we conclude that for an aerobic organism, if weak homeostasis applies then there is a positive net import of Gibbs energy at constant and at increasing biomass (see the entries (2,1) and (4,1) in Table 4.2). When the organism's

		Aerobic and	Anaerobic and		Anaerobic and
		exothermic	exothermic		endothermic
Constant Biomass	Chemical Entropy	0	+	-	-
	Gibbs energy	+	+	+	+
Increasing biomass	Chemical Entropy	+	+	+/-	+/-
	Gibbs energy	+	+	+	+
Decreasing biomass	Chemical Entropy	-	+/-	-	-
	Gibbs energy	+/-	+/-	+/-	+/-

Table 4.2: Net import of chemical entropy and Gibbs free energy in the overall metabolism for 1) exothermic aerobic organisms, 2) exothermic anaerobic organisms, and 3) endothermic anaerobic organisms. All organisms are at constant chemical composition (weak homeostasis). Organisms are either at constant, increasing or decreasing biomass. There are two columns for the anaerobic and exothermic organisms because 1) they can either have a positive or negative net import of chemical entropy at constant biomass (1st line) and 2) this is related to their behavior when they are increasing or decreasing their biomass (3rd and 5th lines).

biomass is decreasing the net import of Gibbs energy is positive or negative (see the entry (6,1) in Table 4.2).

4.3.4 Constraints imposed by the second law and DEB Theory

In the previous sections we made predictions on the sign of the net input into the organism of Gibbs free energy and chemical entropy. These results are synthesized in Table 4.2. These predictions are made for aerobic exothermic organisms, anaerobic exothermic organisms and anaerobic endothermic organisms in a constant environment, for three situations: (1) steady-state (constant biomass), (2) non steady state with increasing biomass and (3) non steady state with decreasing biomass. The imposition of a constant environment is a sufficient condition for weak homeostasis, i.e., constant biomass molar entropy. Table 4.2 highlights the differences between aerobic and anaerobic organisms in a constant environment: (1) only anaerobic organisms can be endothermic, (2) only anaerobic organisms can have a net negative import of chemical entropy while increasing their biomass and (3) only anaerobic organisms can have a net positive import of chemical entropy while decreasing their biomass. Apparently, anaerobic organisms have a higher thermodynamic flexibility.

For steady state, the internal entropy production ($\dot{\sigma} > 0$) implies a relation between

the heat and the chemical entropy exchange with the environment (see Eq. 4.21). Based on this, Stockar and Liu [31] distinguish different overall metabolisms: 'entropy neutral growth' (1^{st} column in Table 4.2), 'entropy driven growth' (3^{rd} and 4^{th} columns in Table 4.2) and 'entropy retarded growth' (2^{nd} column in Table 4.2) respectively for a null, negative or positive net import of chemical entropy.

However, the classification of Stockar and Liu [31] developed for the steady-state should not be used for the non steady state because it is misleading. An example is provided by aerobic organisms increasing their biomass. In this case they would be classified as 'entropy retarded' when the mechanism used to get rid of entropy production is heat dissipation only (see Eq. 4.24).

In non steady-states, organisms can get rid of internal entropy production by using an additional mechanism: accumulation of chemical entropy in new biomass. This mechanism can be called 'dilution of entropy production by growth'. The importance of the mechanism of accumulation of chemical entropy in biomass is fully dependent on the distinction introduced by DEB theory between reserve and structure. For an organism that is not in steady state but has a constant chemical composition (weak homeostasis) the capacity of this mechanism remains constant per C-mol of biomass increase (decrease) because the additional C-mol has the same entropy. In constrast, when the reserve density changes, the chemical composition of biomass and its entropy also change. Thus, the additional C-mol of biomass increase (decrease) has a different capacity to accumulate chemical entropy.

4.3.5 Calorimetry

Indirect Calorimetry

Indirect calorimetry is an empirical method of estimating heat production based on the measurements of gaseous exchanges and the nitrogenous waste flux [8] using multiple linear regression; see [3] or [24] for a good overview.

A theoretical underpinning for indirect calorimetry was provided by Kooijman [22]. Here, we obtained the linearity between the dissipated heat flux and the fluxes in a simpler and more direct way by using only a subset of DEB theory: the existence of strongly homeostatic reserve and structure.

We obtain the linear dependence between the mineral fluxes and the dissipated heat

by combining Eq. 4.13 with Eq. 4.14,

$$\dot{P}_{T+} = (\overline{\boldsymbol{h}}_{\mathcal{O}}^T \boldsymbol{n}_{\mathcal{O}}^{-1} \boldsymbol{n}_{\mathcal{M}} - \overline{\boldsymbol{h}}_{\mathcal{M}}^T) \dot{\boldsymbol{J}}_{\mathcal{M}}.$$
(4.25)

The coefficients that are obtained by linear regression are given by the expression in parentheses and can be computed without knowing any biochemical details. Only the chemical composition and the enthalpies of the reserve, the structure and the input and output products are needed. This result is a theoretical basis for the linear dependence between the mineral fluxes and the dissipated heat because it equates the total dissipating heat to a weighted sum of consumed dioxygen, produced carbon dioxide, nitrogen waste and water.

Thornton's rule

In the literature, Thornton's rule [33] is used to estimate heat production in aerobic organisms. This rule establishes a constant proportionality between the heat released in the combustion of organic compounds and the consumed oxygen: 444 kJ per mol of O_2 consumed. Recently the usefulness of this rule has been questioned by an experimental study made by Walsberg and Hoffman [34] because significant variations were obtained experimentally in the amount of heat released per mol of O_2 in a Kangoroo rat and a dove.

We now use DEB theory to obtain the conditions that keep constant the proportionality coefficient, h_{OT} , in the amount of heat released,

$$\dot{P}_{T+} = h_{OT} \dot{J}_{O_2}.$$
(4.26)

If the heat and the oxygen flows in Eq. 4.26 are written as functions of the organic fluxes using Eq. 4.13 and Eq. 4.25, then the proportionality coefficient is given by:

$$h_{OT} = \frac{(\overline{\boldsymbol{h}}_{\mathcal{O}}^{T} - \overline{\boldsymbol{h}}_{\mathcal{M}}^{T} \boldsymbol{n}_{\mathcal{O}}^{-1} \boldsymbol{n}_{\mathcal{O}}) \dot{\boldsymbol{J}}_{\mathcal{O}}}{\boldsymbol{n}_{\mathcal{M}}^{-1}(3, :) \boldsymbol{n}_{\mathcal{O}} \dot{\boldsymbol{J}}_{\mathcal{O}}}$$
$$= \frac{\sum_{i=1}^{i=3} (\boldsymbol{h}_{\mathcal{O}}(i) - \overline{\boldsymbol{h}}_{\mathcal{M}}^{T} \boldsymbol{n}_{\mathcal{M}}^{-1} \boldsymbol{n}_{\mathcal{O}}(:, i)) \dot{\boldsymbol{J}}_{\mathcal{O}}(i)}{\sum_{i=1}^{i=3} \boldsymbol{n}_{\mathcal{M}}^{-1}(3, :) \boldsymbol{n}_{\mathcal{O}}(:, i) \dot{\boldsymbol{J}}_{\mathcal{O}}(i)}$$
(4.27)

where i = 1 is food, X, i = 2 is reserve, E, i = 3 is structure, V, $n_{\mathcal{O}}(:,i)$ is column i of matrix $n_{\mathcal{O}}$ and $n_{\mathcal{M}}^{-1}(3,:)$ is the 3^{rd} line of matrix $n_{\mathcal{M}}^{-1}$; it is the 3^{rd} line that appears because the 3^{rd} column of matrix $n_{\mathcal{M}}$ has the chemical composition of O_2 . Eq. 4.27 can be written as:

$$h_{OT} = \frac{\sum_{i=1}^{i=3} \dot{P}_{Ti+} \dot{\boldsymbol{J}}_{\mathcal{O}}(i)}{\sum_{i=1}^{i=3} \dot{J}_{O_2}(i) \dot{\boldsymbol{J}}_{\mathcal{O}}(i)} = \frac{\sum_{i=1}^{i=3} h_{OT_i} \dot{J}_{O_2}(i) \dot{\boldsymbol{J}}_{\mathcal{O}}(i)}{\sum_{i=1}^{i=3} \dot{J}_{O_2}(i) \dot{\boldsymbol{J}}_{\mathcal{O}}(i)},$$
(4.28)

where \dot{P}_{Ti+} is the heat released in the complete combustion of one C-mol of the organic compound *i* and $n_{\mathcal{M}}^{-1}(3,:)n_{\mathcal{O}}(:,i)$ is the number of O_2 moles consumed, $\dot{J}_{O_2}(i)$, in the complete combustion of a C-mol of the organic compound *i*. Because the heat released in the complete combustion of an organic compound is constant it can be written as the product of a constant h_{OT_i} and $\dot{J}_{O_2}(i)$.

With Eq. 4.28 the coefficient h_{OT} can be interpreted as a mean of the heats released per each mol of O_2 that would be spent in the complete combustion of each organic compound weighted by its net flow. For h_{OT} to be constant it has to be independent of the values of the organic flows $J_O(i)$. For this to occur, the coefficients h_{OT_i} must be equal, i.e., the heat released per mole of O_2 for each organic compound must be the same, which is usually not the case.

In the remainder of this section, we use DEB theory to obtain an expression that establishes the link between the coefficient of proportionality between the heat released and the oxygen flow and the internal energetics of the organism. In the literature this coefficient has already been used to assess the metabolic pathways in aerobic organisms. For example, Hansen et al. [21] obtained an expression that explains the difference between the mean accepted value for Thornton's coefficient and the observed proportionality coefficient by the existence of anaerobic reactions with an enthalpy change different from zero.

Eq. 4.4 and 4.9 establish a connection between the organic flows of food, j_X , and structure, j_V with the assimilation, \dot{p}_A , and growth powers, \dot{p}_G . The flow of reserve can be written as a function of the three powers using Eq. 4.2 and Eq. 4.1:

$$j_E = (\dot{p}_A - \dot{p}_M - \dot{p}_G)/\mu_E. \tag{4.29}$$

The flows j_X , j_V and j_E multiplied by M_V , can be assembled as:

$$\dot{\boldsymbol{J}}_{\mathcal{O}} = \boldsymbol{\eta}_{\mathcal{O}} \dot{\boldsymbol{p}} M_V, \tag{4.30}$$

where \dot{p} is the vector with the three powers of assimilation, growth and maintenance per C-mol of structure and $\eta_{\mathcal{O}}$ is the matrix with the coefficients that link each organic flow with \dot{p}_A , \dot{p}_M and \dot{p}_G . With Eq. 4.30 and Eq. 4.28 the coefficient of proportionality h_{OT} can be written as:

$$h_{OT} = \frac{\sum_{i=1}^{i=3} h_{OT_i} \dot{J}_{O_2}(i) \boldsymbol{\eta}_{\mathcal{O}}(i,:) \dot{\boldsymbol{p}}}{\sum_{i=1}^{i=3} \dot{J}_{O_2}(i) \boldsymbol{\eta}_{\mathcal{O}}(i,:) \dot{\boldsymbol{p}}}.$$
(4.31)

Eq. 4.31 establishes a link between the proportionality coefficient and the organism's internal energetic flows: assimilation, maintenance and growth. Again, if the coefficients h_{OT_i} are the same, h_{OT} is independent of the relative amounts of energy spent in each internal process; otherwise its change can be used to assess the internal allocation of energy in the organism between assimilation, growth and maintenance.

4.4 Empirical Application

To demonstrate the practical applicability of the well-tested DEB theory to quantify thermodynamic changes in organisms we use published data on *Klebsiella aerogenes* growing in a continuous culture at different steady-states obtained from Esener et al. [14, 15]. *Klebsiella aerogenes* is growing aerobically on glycerol as the sole carbon and energy source, using ammonia as the N source, at a temperature of 35 ° C and a pH of 6.8 [14]. The products are only biomass (structure plus reserve), carbon dioxide and water [13]. The synthetic medium used by Esener et al. [13] is described by Evans et al. in [16]. More details about the experimental setup are given by Esener et al. [14].

Measurements include 26 measures of the specific CO_2 production rate (mol of CO_2 per C-mol of biomass inside the chemostat), 26 measures of the specific O_2 consumption rate, 26 measures of the yield of biomass on substrate Y_{WX} (C-mol biomass produced per C-mol of substrate consumed) and 9 measurements of the biomass composition at different steady-states [14, 15]. Each steady-state is characterized by a different throughput or dilution rate, d. The dilution rate is equal to the growth rate, j_V , because otherwise the biomass concentration in the chemostat would increase or decrease. The maximum growth rate was also measured by [14, 15].

4.4.1 **DEB** parameters

We used the measurements to estimate the following essential DEB parameters: k_E , y_{XE} , y_{EV} , \dot{k}_M , g and the chemical compositions of a C-mol of structure $n_{CV} \equiv 1$, n_{HV} , n_{OV} , n_{NV} and a C-mol of reserve $n_{CE} \equiv 1$, n_{HE} , n_{OE} , n_{NE} .

The method is as follows. 1) We depart from a first guess of 13 DEB parameters. 2) For each steady-state we compute DEB variables like the functional response and the reserve density using the DEB parameters. 3) For each steady-state we calculate the predicted values of CO_2 , O_2 , Y_{WX} and biomass composition. 4) We compute the difference between the predicted values and the 105 measurements and make another estimation of the DEB parameters using the Newton-Raphson method. Steps 2-4 are repeated in order to minimize the sum of the squared errors. A detailed description of steps 1-3 is given below.

Estimation method

We depart from a first estimation of parameters k_E , y_{XE} , y_{EV} , k_M and g and the chemical compositions of structure n_{CV} , n_{HV} , n_{OV} , n_{NV} and reserve n_{CE} , n_{HE} , n_{OE} , n_{NE} .

Eq. 4.12 is then used to compute the functional response f for each steady-state. This equation is also used to compute the maximum growth rate, r_m , that occurs when f is equal to one.

The value for m_{Em} is computed from the definition of investment ratio g given by Eq. 4.11. For each steady-state Eq. 4.5 is used to compute the reserve density m_E because $dm_E/dt = 0$.

With the reserve density m_E and the parameters n_{CV} , n_{HV} , n_{OV} , n_{NV} , n_{CE} , n_{HE} , n_{OE} and n_{NE} , the chemical composition of a C-mol of biomass is computed using:

$$n_{HW} = (n_{HV} + m_E n_{HE}) / (1 + m_E), \tag{4.32}$$

$$n_{NW} = (n_{NV} + m_E n_{NE}) / (1 + m_E), \tag{4.33}$$

and

$$n_{OW} = (n_{OV} + m_E n_{OE}) / (1 + m_E).$$
(4.34)

The yield of biomass produced $j_V(1+m_E)$, on substrate consumed fj_{Xm} , is computed from:

$$y_{WX} = j_V (1 + m_E) / (f j_{Xm}).$$
(4.35)

The substrate consumption rate per mole of structure, j_X , the structure production rate per mole of structure, j_V , and the reserve production rate per mole of structure, j_E , are:

$$j_X = f j_{Xm}, \tag{4.36}$$

$$j_V = j_V, \tag{4.37}$$

$$j_E = j_V m_E. \tag{4.38}$$

The flows in Eq. 4.36, 4.37 and 4.38 are divided by $(1 + m_E)$ to be converted to flows per C-mol of biomass. Then, these flows are used together with the mass balance, Eq. 4.13, applied to each steady-state of the *Klebsiella aerogenes* culture to compute the CO_2 , H_2O , O_2 and NH_3 flows. In Eq. 4.13, n_M is the matrix with the chemical composition of minerals (CO_2 , H_2O , O_2 and NH_3) and n_O is the matrix with the chemical composition of organic compounds (X, E, V).

Parameter	Value	Units
\dot{k}_M	0.021	h^{-1}
\dot{k}_E	2.11	h^{-1}
y_{VE}	0.904	_
y_{XE}	1.345	_
g	1	—

Table 4.3: DEB parameters for *Klebsiella aerogenes*.



Figure 4.2: Measurements (points) and DEB model results (lines). Specific rate of consumption of O_2 (×), specific rate of production of CO_2 (+) and yield (*) vs. dilution rates. Units are: mol.C-mol⁻¹.h⁻¹ for O_2 and CO_2 and C-mol.C-mol⁻¹ for the yield.

Results

The elemental composition of structure and reserve are $CH_{1.64}O_{0.379}N_{0.198}$, and $CH_{1.66}O_{0.422}N_{0.312}$, respectively. The values obtained for the other parameters are listed in Table 4.3. The maximum growth rate measured is $1.052 h^{-1}$ and the value obtained with the DEB model is $1.044 h^{-1}$. The comparison between the other measurements and the DEB model results is presented in Fig. 4.2 and 4.3. The root mean square error for O_2 and CO_2 are 0.0088 and 0.0086 mol.C-mol⁻¹.h⁻¹ respectively and the root mean square error for Y_{WX} is $0.0249 \text{ C-mol.C-mol}^{-1}$. The root mean square error for n_{HW} , n_{OW} and n_{NW} are 0.009, 0.0191 and 0.0113 C-mol.C-mol⁻¹ respectively. Since the fits are very good, and the DEB model obeys mass balances, we have an automatic check on the empirical mass balances, i.e., the measurements obey the mass balance. The change in the chemical composition of biomass (see Fig. 4.3) is not very significant because the chemical compositions of a C-mol of structure and a C-mol of reserve are similar with the exception of the amount of nitrogen.

The values obtained for the reserve density for each dilution rate are in Fig. 4.4.



Figure 4.3: Measurements (points) and DEB model results (lines). The variable chemical composition of biomass, $n_{HW}(+)$, $n_{NW}(\times)$ and $n_{OW}(*)$ vs. dilution rates.



Figure 4.4: Reserve density vs. dilution rates.



Figure 4.5: The assimilation, growth and maintenance energy flows per C-mol of biomass vs. dilution rates.

We also obtained the flows per unit of biomass of assimilation, maintenance and growth (Eq. 4.4, Eq. 4.8 and Eq. 4.9) shown in Fig. 4.5. The ratio of the assimilation flow to the food flow,

$$\dot{p}_A/\dot{p}_X = (1/y_{XE})(\mu_E/\mu_X),$$
(4.39)

is obtained with $\dot{p}_X = j_X \mu_X$ and Eq. 4.4. This ratio is constant because Eq. 4.39 is a function of parameters only. The assimilation flow increases with the throughput rate because the flow of food, X, also increases. Although the maintenance flow per C-mol of structure is constant because it is a function of parameters only (Eq. 4.8), the maintenance flow per C-mol of biomass decreases with the dilution rate because the reserve density increases. This occurs because it is the structure that is costly in terms of maintenance and not the reserve. The growth flow per C-mol of biomass increases with the dilution rate because it is proportional to the specific growth rate, j_V , (see Eq. 4.9) and the specific growth rate (equal to the dilution rate) increases more than the reserve density (see Fig. 4.4).

The ratio of energy spent on maintenance to energy spent on growth, given by k_M/r , increases with a decreasing dilution rate, i.e., growth per C-mole of biomass becomes more expensive with decreasing dilution rate. The energetic explanation rooted in DEB theory for this behavior is: a decrease in the dilution rate translates into a lower catabolic power (defined in Fig. 5.1) and because maintenance has priority over growth a higher fraction of the catabolic power is spent on maintenance and a lower one on growth.

4.4.2 Structure and reserve specific enthalpies and entropies

The enthalpy balance

The enthalpy balance applied to the chemostat is given by Eq. 4.14. For each steady-state $\dot{J}_{\mathcal{M}}$ and $\dot{J}_{\mathcal{O}}$ are computed using the DEB model with the parameters estimated in the previous section. Formation enthalpies of CO_2 , O_2 , H_2O taken from [9] and formation enthalpy of glycerol taken from [5] were corrected for the temperature of the experiment using the specific heats at constant pressure taken from [9]. The formation enthalpy for NH_3 aq. is for 37 °C and was taken from [20]. Values used are in Table 4.4. The enthalpies of structure and reserve are unknown but constant for all steady-states (strong homeostasis assumption). The dissipating heat depends on the steady-state and is unknown.

Eq. 4.14 is applied to different steady-states and solved for the dissipating heats and the enthalpies of structure and the reserve. This system of equations involves 2 extra unknowns. The two missing constraints were taken from Roels [30] (i.e. dissipating heats for two steady-states).

Released heats are presented in Fig. 4.6. Specific released heat ranges from -20 kJ/Cmol. to -253 kJ/C-mol (see Fig. 4.6) for increasing throughput rates. The values obtained for the released heats are of the same magnitude as the released heats presented by [31] for aerobic growth of different microorganisms on various substrates. The heat flow for each dilution rate is much higher than any of the assimilation, growth and maintenance flows (Fig. 4.6). Its magnitude is four to five times higher than the magnitude of the assimilation flow. Therefore, the energy dissipated as heat, which in an aerobic organism is a measure of the production of entropy, is a very significant energy drain.

We obtained a molar enthalpy of formation of -33 kJ/C-mol for the reserve and a molar enthalpy of formation of -107 kJ/C-mol for the structure. Thus, the formation of 1 C-mol of structure and 1 C-mol of reserve from their components at a reference state are both exothermic reactions, with the former being more exothermic. The steady-state enthalpy of the biomass, $(\overline{h}_V + \overline{h}_E m_E)/(1 + m_E)$, decreases with the dilution rate from -76 kJ/C-mol to -105 kJ/C-mol. In the literature we found no formation enthalpy values for *Klebsiella aerogenes*. Some formation enthalpy values referred in the literature are: -95.68 kJ/C-mol for *Escherichia coli* growing aerobically on succinic acid [2], -97.8 kJ/Cmol for the same microorganism [1] and -133.09 kJ/C-mol for *Saccharomyces cerevisae*.



Figure 4.6: The heat production rate per C-mol of chemostat biomass per hour (+) and heat production per mol of O_2 consumed (\times) (Thornton's rule) vs. dilution rates. Units are: (\times) in kJ.mol O_2 and (+) in kJ.C-mol⁻¹.h⁻¹.

The entropy balance

The entropy balance applied to the chemostat is given by Eq. 4.16. For each steadystate $\dot{J}_{\mathcal{M}}$ and $\dot{J}_{\mathcal{O}}$ are computed using the DEB model with the parameters estimated in section 4.4.1. Absolute entropies were taken from Dean [9] and corrected for temperature (see Table 4.4). The entropies of structure and reserve are unknown but constant for all steady-states (strong homeostasis assumption).

Eq. 4.16 is applied to different steady-states and solved for the entropies of structure and the reserve with nonlinear regression. We obtained a molar entropy of 74.8 J/Cmol.K for the reserve and a molar entropy of of 52.0 J/C-mol.K for the structure. To test the reliability of these specific entropy values we computed the left-hand side of Eq. 4.16 for many steady-states. It is very close to zero, i.e., it is at maximum 0.04% of any other term in the equation. The first important remark is that these entropies are not null and are different from the entropies of the inputs and outputs. The steady-state entropy of the biomass, $(\bar{s}_V + \bar{s}_E m_E)/(1 + m_E)$, increases from 52.4 J/C-mol.K to 61.4 J/C-mol.K with increasing dilution rate (see Fig. 4.7). The molar biomass entropy increases with the increasing dilution rate because the reserve density increases. In the literature we found no absolute entropy values for *Klebsiella aerogenes*. Other absolute entropy values comprise: 94.4 J/C-mol.K for dried *Escherichia coli* growing on succinic acid [2] and 34.17 J/C-mol.K for *Saccharomyces cerevisae* [4].

We also compare the molar biomass entropy obtained with DEB with the entropy given by the empirical rule proposed by Battley [6] for organic substances (see Fig. 4.7). The entropy of the biomass computed by DEB theory increases more with the dilution rate and is significantly higher. However, the application of Battley's rule to dead bio-



Figure 4.7: Entropy of biomass computed with DEB model (+) and entropy of biomass given by Battley's rule (\times) vs. dilution rates.

Formula	State	Enthalpy (kJ/mol)	Entropy (J/mol.K)
CO_2	g	-393.14	214.70
H_2O	1	-285.83	72.331
O_2	g	0	205.80
NH_3	aq.	-132.5	112.34
C mol glycerol	aq. ⁸	-225.52	69.743

Table 4.4: Enthalpies and entropies at 35 °C.

mass of *Saccharomyces cerevisae* [6] gave a very similar result to the entropy obtained experimentally in [4]. Therefore: (1) the entropy of dead biomass is different from the entropy of living biomass and (2) Battley's rule should not be applied for living biomass. The entropies of structure and reserve should be computed for other organisms in order to evaluate the generality of these results.

In the literature [1] the entropy of biomass has been compared to the entropy of the substrate. In our case, the biomass entropy ranges from 3.03 J/g.K to 2.69 J/g.K while the entropy of glycerol is 2.03 J/g.K. The fact that 1 gram of biomass has a higher entropy then 1 gram of substrate is in accordance with results obtained by Battley [1] for *Escherichia coli* and succinic acid. Battley argues that this result points to the fact that specific entropy is not related with complexity, otherwise, how could the lower entropy value for the substrate be explained? We disagree because we think that 1 gram is an arbitrary quantity: why not compare 1 C-mol? In that case the entropy of a C-mol of structure is lower than the entropy of a C-mol of glycerol.

4.4.3 Calorimetry

The heats released in the complete combustion of compound *i* per mol of consumed O_2 are: -472 kJ for the food, -485 kJ for the reserve and -447 kJ for the structure of *Klebsiella aerogenes*. These values are more or less in agreement with the values obtained theoretically by Gnaiger and Kemp [18] for other organic compounds. Although, the heats of combustion of *X*, *E* and *V* are similar, the net flow of food is positive while the net flows of structure and reserve are negative, suggesting that the coefficient h_{OT} is not bounded by the values of the various h_{OT_i} . This is indeed the case for the dilution rates considered, where the ratio of the heat flow to the oxygen flow varies between -476 kJ and -507 kJ per mol of O_2 consumed (see Fig. 4.6).

4.5 Conclusions

The thermodynamic analysis made in this chapter is applicable to any organism because 1) it is based on a thermodynamic formalism applicable to any open thermodynamic system and 2) uses a general model to describe the internal structure of the organism - the Dynamic Energy Budget Model.

We obtain the thermodynamic constraints for organisms with constant food availability, i.e., organisms with a constant chemical composition (DEB's weak homeostasis assumption). These constraints are that only anaerobic organisms can (1) be endothermic, (2) have a net negative import of chemical entropy while increasing their biomass and (3) have a net positive import of chemical entropy while decreasing their biomass (see Table 4.2). Apparently, anaerobic organisms have a higher thermodynamic flexibility.

We obtain Thornton's coefficient as a function of either (1) the flows of organic compounds (see Eq. 4.28), or (2) assimilation, maintenance and growth (see Eq. 4.31) using DEB theory. These relationships are useful in providing new insights into the discrepancies obtained between Thornton's constant and experimental values.

We use experimental data on the aerobic growth of *Klebsiella aerogenes* to obtain molar enthalpies and entropies for the reserve and structure. The knowledge that these properties are constant (DEB's strong homeostasis assumption) is sufficient to compute changes in the enthalpy and in the entropy of living biomass that are known to accompany changes in the reserve density. The importance of being able to compute thermodynamic properties as a function of the amount of reserves has been acknowledged in the literature, e.g., Battley [1] computes the enthalpy of a C-mol of *E. coli* under conditions that impose

that 'no storage materials are produced'. Previously, the entropy of living organisms was obtained either by (1) experimental methods or by (2) Battley's empirical rule [4]. The use of DEB theory for these computations is better than the methods referred because (1) experimental methods are destructive and (2) Battley' rule does not give results similar to the results we obtained. This last point suggests that the entropy of living biomass is different from the entropy of dead biomass because Battley's rule has been validated with good results for dead biomass and organic compounds.

We introduce the mechanism of 'dilution of entropy production by growth' for organisms that are not in steady-state. The capacity of this mechanism to store entropy in new biomass changes with DEB's reserve density because the molar entropy of the reserve is different from the molar entropy of the structure. We proved this for *Klebsiella aerogenes* where the entropies obtained are different from zero and the structure's molar entropy is significantly lower than the reserve's. Additionally, this result suggest that the reserve density concept of DEB theory is essential in discussions concerning the relationship between organization and entropy because the entropy of the organism is a function of the reserve density but the entropy of the structure, which can be related with the organization of the organism, is not.

The development of the generic thermodynamic analysis carried out in this chapter can contribute to enlighten the discussions mentioned here and others including thermodynamic measures of biological organization, the explanation of evolutionary increase in size and evolutionary strategies of energy allocation.

4.6 Appendices

4.6.1 Appendix I

In this appendix we briefly explain the notation used throughout the chapter. A list of compounds is in Tab. 4.1.

Mass of compound *, M_* , is measured in moles for P (product) or X (food) and in C-moles for E (reserve), V (structure) and W (biomass). The ratio $m_E \equiv M_E/M_V$ is the reserve density of the organism and m_{Em} is the maximum reserve density.

Mass flows of compound *, j_* , are measured in moles (or C-moles) per C-mol of structure per time. There is one exception to this rule, j_{Xm} , which is the flow of X measured in moles per C-mol of structure per unit time at the maximum ingestion rate. If mass flows are measured in moles or C-moles per unit time they are represented as

 $J_* \equiv j_* M_V.$

Coefficients that relate two mass flows are y_{*1*2} . They represent the number of moles of *1 needed to produce one mol of *2. In the assimilation reactor food is converted into reserve, y_{XE} , and in the growth reactor reserve is converted into structure, y_{EV} .

Energy flows, \dot{p}_* , are measured in Gibbs energy per C-mol of structure per unit time. The * = X, A, C, M, G, stands for the process which the energy flow is associated with: X (feeding), A (assimilation), C (catabolism), M (maintenance) and G (growth). Chemical potentials convert mass flows to energy flows: μ_X converts the flow of food to \dot{p}_X ; μ_E converts the flow of reserve that exits the assimilation reactor into \dot{p}_A , the flow of reserve that exits the reserve compartment into $\dot{p}_C = \dot{p}_M + \dot{p}_G$.

The energy flow \dot{p}_X is associated with the adimensional functional response $f(X) \in [0, 1]$ that is equal to 1 at abundant food $(X \to \infty)$ and 0 at no food availability (X = 0). Other DEB parameters include: the reserve turnover rate \dot{k}_E and the maintenance rate coefficient \dot{k}_M , both parameters' dimensions are per time. The first is related with the velocity of use of the reserve and the second with the velocity of degradation of the structure. Also related with the structure there is the adimensional investment ratio g, a measure of the relative cost of building structure.

Thermodynamic properties have the usual notation: \overline{g}_* is the molar Gibbs energy, \overline{h}_* is the molar enthalpy, \overline{s}_* is the molar entropy and \overline{u}_* is the molar internal energy and μ_* is the chemical potential of compound *, T is the temperature, \dot{P}_{T+} is the rate of heat release by the organism and $\dot{Q}_{reactions}$ is the rate of heat release by all chemical reactions inside the organism.

Vectors and matrices are in bold. The transpose of a vector is indicated by T and the inverse of a matrix is indicated by $^{-1}$. An overbar means that it is a molar quantity.

The matrix with the chemical composition is n: $n_{\mathcal{O}}$ is the matrix with the chemical composition of the organic compounds (X, P, E, V) and $n_{\mathcal{M}}$ is the matrix with the chemical composition of the minerals $(CO_2, O_2, H_2O, N_{waste})$. Each entry in these matrices, n_{*1*2} , is the number of atoms of element *1 in compound *2.

4.6.2 Appendix II

The dynamics of the reserve density

$$\frac{dm_E}{dt} = fm_{Em}\dot{k}_E - \frac{\dot{p}_C}{\mu_E} - m_E \frac{1}{M_V} \frac{dM_V}{dt},$$
(4.40)

is obtained by combining Eq. 4.3 with Eq. 4.4, Eq. 4.6 and $j_X = f j_{Xm}$. One of the assumptions of DEB theory is that the mobilization of reserves, i.e., the catabolic power, cannot depend on food availability, which means that it can only depend on the state variables, the reserve density, m_E , and the amount of structure, M_V . Under this assumption, the last two terms in Eq. 4.40 are a function only of m_E and M_V . So, Eq. 4.40 can be written as:

$$\frac{dm_E}{dt} = fm_{Em}\dot{k}_E - \Phi(m_E, M_V).$$
(4.41)

At constant food, the weak homeostasis assumption implies that the reserve density is constant $m_E = m_E^*$. Thus,

$$fm_{Em}\dot{k}_E = \Phi(m_E^*, M_V).$$
 (4.42)

However, the weak homeostasis assumption also implies that m_E^* is dependent on food level but not on the amount of structure M_V because the organism can grow with a constant reserve density. Thus,

$$\Phi(m_E^*, M_V) = H(m_E^*), \tag{4.43}$$

because $fm_{Em}k_E$ does not depend on M_V . The function $\Phi(m_E, M_V)$ can be generalized out of steady-state as $\Phi(m_E, M_V) = H(m_E) + (m_E^* - m_E)G(m_E, M_V)$. With this specification for $\Phi(m_E, M_V)$ eq. 4.41 becomes

$$\frac{dm_E}{dt} = fm_{Em}\dot{k}_E - H(m_E) - (m_E^* - m_E)G(m_E, M_V).$$
(4.44)

With Eq. 4.40 and Eq. 4.44 the catabolic flux per C-mol of structure is:

$$\dot{p}_{C} = \mu_{E} H(m_{E}) + \mu_{E} (m_{E}^{*} - m_{E}) G(m_{E}, M_{V})$$

$$- \mu_{E} m_{E} \frac{1}{M_{V}} \frac{dM_{V}}{dt}.$$
(4.45)

Additionally, $G(m_E, M_V) = 0$ because m_E^* is a function of food availability and according to DEB theory the catabolic power cannot depend on food availability. Thus,

$$\dot{p}_C = \mu_E H(m_E) - \mu_E m_E \frac{1}{M_V} \frac{dM_V}{dt}.$$
 (4.46)

Eq. 4.46 can be written as a function of m_E , M_V and parameters using Eq. 4.10:

$$\dot{p}_{C} = \mu_{E} H(m_{E}) - \mu_{E} m_{E} \left(\frac{\dot{k}_{E} \frac{m_{E}}{m_{Em}} - \dot{k}_{M} g}{\frac{m_{E}}{m_{Em}} + g} \right).$$
(4.47)

To proceed with the derivation we need another of DEB's assumptions: the partitionability of reserves. If the organism's reserve is partitioned among different aggregates then the catabolic power that is mobilized from each aggregate must be proportional to the amount of energy embodied in it:

$$M_V \dot{p}_C(\lambda m_E, M_V, \lambda g) = \lambda M_V \dot{p}_C(m_E, M_V, g).$$
(4.48)

Also, the number of moles allocated to growth per mole of structure, from each aggregrate, must be proportional to the amount of energy embodied in it.

The imposition that the catabolic power given by eq. 4.47 must obey eq. 4.48 implies that $\lambda H(m_E) = H(\lambda m_E)$. Therefore $H(m_E) = \gamma m_E$ and Eq. 4.46 simplifies to:

$$\dot{p}_C = \mu_E m_E \gamma - \mu_E m_E \frac{1}{M_V} \frac{dM_V}{dt}.$$
(4.49)

With Eq. 4.49 the reserve density dynamics given by Eq. 4.40 simplifies to:

$$\frac{dm_E}{dt} = fm_{Em}\dot{k}_E - m_E\gamma. \tag{4.50}$$

At abundant food availability, (1) the ingestion rate $j_X = j_{Xm}$ implying that f = 1 and (2) the steady-state reserve density is $m_E^* = m_{Em}$. With conditions (1-2) $\gamma = \dot{k}_E$ and

$$\frac{dm_E}{dt} = \dot{k}_E \left(fm_{Em} - m_E \right). \tag{4.51}$$

References

- [1] E. Battley. On the enthalpy formation of *Escherichia coli* K-12 cells. *Biotechnology and Bioengineering*, 39:5–12, 1992.
- [2] E. Battley. Calculation of entropy change accompanying growth of *Escherichia coli* K-12 on succinic acid. *Biotechnology and Bioengineering*, 41:422–428, 1993.
- [3] E. Battley. The advantages and disadvantages of direct and indirect calorimetry. *Thermochimica Acta*, 250:337–352, 1995.
- [4] E. Battley. Heat capacity measurements from 10 to 300 K and derived thermodynamic functions of lyophilized cells of *Saccharomyces cerevisae* including the absolute entropy and the entropy formation at 298.15 K. *Thermochimica Acta*, 298:37– 46, 1997.
- [5] E. Battley. The development of direct and indirect methods for the study of the thermodynamics of microbial growth. *Thermochimica Acta*, 309:17–37, 1998.

- [6] E. Battley. An empirical method for estimating the entropy of formation and the absolute entropy of dried microbial biomass for use in studies on the thermodynamics of microbial growth. *Thermochimica Acta*, 326:7–15, 1999.
- [7] A. Bejan. Advanced Engineering Thermodynamics. Wiley, Durham, NC, 1988.
- [8] K. Blaxter. *Energy Metabolism in Animals and Man*. Cambridge University Press, Cambridge, 1989.
- [9] J. Dean. Lange's Handbook of Chemistry. McGraw-Hill, 12th Ed., New York, NY, 1979.
- [10] Y. Demirel and S. Sandler. Thermodynamics and bioenergetics. *Biophysical Chem-istry*, 97:87–111, 2002.
- [11] P. Duboc and U. Stockar. Energetic investigation of *Saccharomyces cerevisae* during transitions. Part 2. Energy balance and thermodynamic efficiency. *Thermochimica Acta*, 251:131–144, 1995.
- [12] A. Esener, J. Roels, and W. Kossen. Fed-batch culture: Modeling and applications in the study of microbial energetics. *Biotechnology and Bioengineering*, XXIII:1851– 1871, 1981.
- [13] A. Esener, J. Roels, and W. Kossen. The influence of temperature on the maximum specific growth rate of *Klebsiella pneumoniae*. *Biotechnology and Bioengineering*, XXIII:1401–1405, 1981.
- [14] A. Esener, J. Roels, and W. Kossen. Dependence of the element composition of *K*. *pneumoniae* on the steady-state specific growth rate. *Biotechnology and Bioengineering*, XXIV:1445–1449, 1982.
- [15] A. Esener, J. Roels, and W. Kossen. Theory and applications of unstructured growth models: kinetic and energetic aspects. *Biotechnology and Bioengineering*, XXV:2803–2841, 1983.
- [16] C. Evans, D. Herbert, and D. Tempest. *The continuous culture of microorganisms*. *Construction of a Chemostat.* Academic Press, London, 1970.
- [17] L. Garby and Larsen P. *Bioenergetics its thermodynamic foundations*. Cambridge University Press, 1995.

- [18] E. Gnaiger and R. B. Kemp. Anaerobic metabolism in aerobic mammalian cells: information from the ratio of calorimetric heat flux and respirometric oxygen flux. *Biochimica et Biophysica Acta*, 1016:328–332, 1990.
- [19] S. R. de Groot and P. Mazur. *Non-Equilibrium Thermodynamics*. North Holland, Amsterdam, 1962.
- [20] Y. Guan and R. B. Kemp. Detection of the changing substrate requirements of cultured animal cells by stoichiometry growth equations validated by enthalpy balances. *Journal of Biotechnology*, 69:95–114, 1999.
- [21] L. D. Hansen, C. Macfarlane, N. McKinnon, B. N. Smith, and R. S. Criddle. Use of calorespirometric ratios, heat per CO₂ and heat per O₂, to quantify metabolic paths and energetics of growing cells. *Thermochimica Acta*, 422:55–61, 2004.
- [22] S. A. L. M. Kooijman. Dynamic Energy and Mass Budgets in Biological Systems. Cambridge University Press, Cambridge, 2000.
- [23] S. A. L. M. Kooijman. Quantitative aspects of metabolic organization: a discussion of concepts. *Philosophical Transactions of the Royal Society of London B*, 356:331– 349, 2001.
- [24] I. Lamprecht. Calorimetry and thermodyamics of living systems. *Thermochimica Acta*, 405:1–13, 2003.
- [25] J. Liu, I. Marison, and U. Stockar. Microbial growth by a net heat up-take: A calorimetric and thermodynamic study on acetrophic methanogenesis by *Methanosarcina barkeri*. *Biotechnology and Bioengineering*, 75:170–180, 2001.
- [26] M. Moran and H. Shapiro. Fundamentals of Engineering Thermodynamics. Wiley, New York, NY, 2000.
- [27] H. Morowitz. Energy Flow in Biology. Academic Press, New York, NY, 1968.
- [28] R. Nisbet, E. Muller, K. Lika, and S. A. L. M. Kooijman. From molecules to ecosystems through dynamic energy budget models. *Journal of Animal Ecology*, 69:913– 926, 2000.
- [29] H. Qian and D. A. Beard. Thermodynamics of stoichiometric biochemical networks in living systems far from equilibrium. *Journal of Biophysical Chemistry*, 114:213– 220, 2005.
- [30] J. Roels. Energetics and Kinetics in Biotechnology. Elsevier, Amsterdam, 1983.
- [31] U. Stockar and J. Liu. Does microbial life always feed on negative entropy? Thermodynamic analysis of microbial growth. *Biochimica et Biophysica Acta - Bioenergetics*, 1412:191–211, 1999.
- [32] U. Stockar, I. Marison, and J. Liu. Endothermic microbial growth. a calorimetric investigation of an extreme case of entropy driven microbial growth. *Pure and Applied Chemistry*, 72:1831–1838, 2000.
- [33] W. Thornton. The relation of oxygen to the heat of combustion of organic compounds. *Philosophical Magazine and Journal of Science*, 33:196–203, 1917.
- [34] G. E. Walsberg and T. C. Hoffman. Direct Calorimetry reveals large errors in respirometric estimates of energy expenditure. *The Journal of Experimental Biology*, 208:1035–1043, 2005.

Chapter 4

98

Chapter 5

From empirical patterns to theory: A formal metabolic theory of life

Abstract

The diversity of life on Earth raises the question of whether it is possible to have a unique theoretical description of the quantitative aspects of the organisation of metabolism. However, similarities between organisms, like von Bertalanffy's growth curves and Kleiber's law on metabolic rate, suggest that mechanisms that control the uptake and use of metabolites are common to all organisms. These and other empirical patterns widespread in biology should be the ultimate test to any metabolic theory that hopes for generality. The present study (1) collects empirical evidence on growth, stoichiometry, feeding, respiration and indirect calorimetry and presents it in a simplified manner as stylized empirical biological facts, (2) formalizes into assumptions and propositions a metabolic theory and (3) proves that these assumptions and propositions are coherent with the stylized empirical facts. This formalized metabolic theory is fully consistent with the Dynamic Energy Budget (DEB) Theory.

Keywords: Dynamic energy budget (DEB) theory, Kleiber's law, von Bertalanffy growth

5.1 Introduction

In the literature, two main approaches are followed to get insights into biological phenomena: (1) the study of the complex set of biochemical reactions ocurring at different rates and (2) the study of the organization of metabolism described by the mass and energy flows inside the organisms. We believe that the modeling of the biochemical networks of reactions that are taking place in the organism is useful but will not by itself lead to an understanding of life because the set of biochemical reactions occuring in the organism can be species-specific and too complex, especially for multicellular organisms. Also, the standard modeling of biochemical networks neglects the spatial structure and the complex transport and allocation processes in the organism.

In contrast, this chapter builds on the premise that the mechanisms that are responsible for the organization of metabolism are not species-specific [41]. This hope for generality is supported by (1) the universality of physics and evolution and (2) the existence of widespread biological empirical patterns among organisms.

The roadmap of this chapter is as follows. In section 5.2, the empirical patterns that characterize metabolism are synthesized and presented in a stylized manner. They are of the utmost importance because any biological non-species specific metabolic theory should predict a qualitative behaviour that is compatible with these facts. We believe that such a theory has already been developed - the Dynamic Energy Budget (DEB) Theory. This theory aims to capture the quantitative aspects of the organisation of metabolism at the organism level with implications for the sub- and supra-organismic levels [41, 47, 60]. In section 5.3, DEB theory is formalized for its standard model, which considers an isomorphic organism, with 1-reserve and 1-structure. This model is assumed to be appropriate for most heterotrophic unicellulars and animals. This theory is formalized in such a way that (1) the assumptions are highlighted and separated from the propositions and (2) the reasoning behind the assumptions and propositions is supported by the stylized empirical patterns or by universal laws. In DEB theory, the difference between species reduces to differences in the set of parameter values. In section 5.4, DEB theory for the relationship between parameters among different species is formalized. Section 5.5 synthesizes the links between empirical patterns, assumptions and propositions and concludes.

5.2 Empirical Patterns

In this section, we summarize the stylized empirical patterns in biology that are essential for a theoretical description of metabolic organization (see Tables 5.1, 5.2). These patterns are related with (1) the metabolic processes that are known to be widespread in organisms including feeding, growth, reproduction, maturation and maintenance; (2) the life-stages, i.e., embryo, juvenile and adult, and (3) the stoichiometry of organisms.

A theory that describes the metabolism of organisms should also be compatible with

		Stylized Facts	Empirical Evidence
Feeding	F1	During starvation, organisms are able to reproduce	animals [29, 36, 37]
	F2	During starvation, organisms are able to grow	animals
			[13, 18, 66, 70, 77, 88]
	F3	During starvation, organisms are able to survive	animals [55, 76]
		for some time	bacteria [53]
Growth	G1	The growth of isomorphic organisms at abundant	animals
		food is well described by the von Bertalanffy	[8, 15, 17, 69, 72, 78]
		growth curve [3, 64]	
	G2	Many species do not stop growing after reproduction has	animals [35, 74]
		started, i.e., they exhibit indeterminate growth [28, 48]	
	G3	Foetuses increase in weight proportional to cubed	animals [30, 89]
		time [30]	
	G4	The logarithm of the von Bertalanffy growth rate of	bacteria [41, p.276-282]
		different species corrected for a common body	yeasts [41, p.276-282]
		temperature decreases almost linearly with the	animals [41, p.276-282]
		logarithm of the species maximum size	
	G5	The logarithm of the von Bertalanffy growth rate for	animals [19], [41, p.96]
		organisms of the same species at different food	
		availabilities decreases linearly with ultimate length	
Respiration	R1	Freshly laid eggs do not use dioxygen in significant	animals [6, 62, 67, 87]
		amounts	
	R2	The use of dioxygen increases with decreasing mass in	animals
		embryos and increases with mass in juveniles and adults	[6, 11, 62, 65, 67, 71, 87]
	R3	The use of dioxygen scales with body weight raised	animals [11, 65, 71]
		to a power close to 0.75 [38]	
	R4	Organisms show a transient increase in metabolic rate	animals [7, 27, 34, 59, 68]
		after ingesting food independent of their body mass -	
		the heat increment of feeding	

Table 5.1: Stylized facts and empirical evidence on feeding, growth and respiration.

		Stylized Facts	Empirical Evidence
Stoichiometry	S 1	Well-fed organisms have a different body chemical	animals [9, 14, 29, 56]
		composition than poorly-fed organisms	yeasts [20]
	S2	Organisms growing with constant food density	animals
		converge to a constant chemical composition	[10, 16, 31, 49, 75]
Indirect	I1	Dissipating heat is a weighted sum of three mass flows:	
Calorimetry		carbon dioxide, dioxygen and nitrogeneous waste	animals [73]
Cells	C1	Cells in a tissue are metabolically very similar	
		independently of the size of the organisms [57]	

Table 5.2: Stylized facts and empirical evidence on stoichiometry, indirect calorimetry and cells.

physics and evolutionary theory. The following laws or physical principles should have a crucial role: (P1) mass and energy are conserved entities; (P2) any energy conversion process leads to dissipation, i.e., entropy production is always positive; (P3) mass transport beween different environments occurs across surfaces; (P4) mass and energy flows into the cell depend only on intensive properties; (P5) the morphology of an organism, i.e., the dependence of the surface area on the volume as the organism grows, controls the relative importance of processes that are proportional to surface area and processes that are proportional to volume. The following evolutionary principle should have an important role: (P6) organisms increased their control over their metabolism during evolution, which allows for some adaptation to environmental changes in a short period. Also, any theory that captures all differences between individuals in differences in parameter values should have (P7) parameter values that are individual-specific and inheritable in a sloppy way, which allows for some adaptation to environmental changes across generations.

5.3 Theory on Metabolic Organization

The standard DEB model considers an isomorphic organism, i.e., an organism whose surface area is proportional to volume to the power 2/3, with 1-reserve and 1-structure. Figure 5.1 summarizes the standard DEB model while Tables 5.3 and 5.4 summarize the notation used throughout the paper.

Assumption 5.3.1 (Strong homeostasis) The biomass of the organism can be partitioned into two categories of generalised compounds, i.e. mixtures of a large number of compounds, which do not change in chemical composition: the structure, V, and the reserve, E. The organism feeds on a resource, X, and produces products, P, also of fixed chemical compositions.

The rationale behind strong homeostasis is (P6). Organisms control their own metabolism through the production of enzymes that catalyse chemical transformations in a constant chemical environment. However, empirical evidence on the variable chemical composition of the organisms, (S1), justifies the need for two aggregate chemical compounds, i.e., structure, V, and reserve, E.

Assumption 5.3.2 (Metabolic Processes) Metabolism can be characterized by the following processes: (1) Feeding, \dot{p}_X , i.e., the uptake of food by the organism; (2) assimilation, \dot{p}_A , i.e., the set of reactions that transform food into reserve and (3) catabolism, \dot{p}_C ,



Figure 5.1: Metabolism in a DEB organism. Circles are processes; rectangles are state variables; arrows are flows of reserve (E), structure(V), minerals, food (X), products(P) or offspring (E_R).

Variable	Dimension	Interpretation	Variable	Dimension	Interpretation
t	Т	time	\dot{p}_X	$e T^{-1}$	Feeding power
X	$\#L^{-3}$	Substrate density	\dot{p}_A	$e T^{-1}$	Assimilation power
E	e	Energy in reserve	\dot{p}_C	$e T^{-1}$	Catabolic power
e	-	Scaled reserve density	\dot{p}_M	$e T^{-1}$	Volume related
					maintenance power
V	L^3	Structural volume	\dot{p}_T	$e T^{-1}$	Surface related
					maintenance power
L	L	Volumetric length	\dot{p}_J	$e T^{-1}$	Maturity maintenance
					power
E_H	e	Energy allocated to	\dot{p}_G	$e T^{-1}$	Growth power
		maturation			
f	_	Scaled functional response	\dot{p}_R	eT^{-1}	Reproduction power

Table 5.3: List of symbols. Dimensions: – no dimension; L length; M mass; T time; # moles or C-moles; e energy. Symbols with (1) $\{\cdot\}$ are per unit surface area, (2) $[\cdot]$ are per unit of structural volume and (3) \cdot are per unit time.

i.e., the mobilization of reserve. So, reserve dynamics is given by:

$$\frac{dE}{dt} = \dot{p}_A - \dot{p}_C. \tag{5.1}$$

The mobilized reserve is allocated to (4) growth, \dot{p}_G , i.e., the increase of structure; (5) somatic maintenance, $\dot{p}_M + \dot{p}_T$, i.e., the maintenance of the structure where \dot{p}_M and \dot{p}_T are associated with volume and surface maintenance costs respectively; (6) maturity maintenance, \dot{p}_J , i.e., the maintenance of the complexity of the structure and (7) maturation, \dot{p}_R , i.e., the increase in the complexity of the structure or (8) reproduction, \dot{p}_R , i.e., the flow of energy to offspring, i.e.,

$$\dot{p}_C = \dot{p}_M + \dot{p}_T + \dot{p}_G + \dot{p}_J + \dot{p}_R.$$
(5.2)

The fraction of catabolic power allocated to somatic maintenance and growth is a general function $0 \le \kappa(V, E) \le 1$, i.e.,

$$\dot{p}_M + \dot{p}_T + \dot{p}_G = \kappa \dot{p}_C. \tag{5.3}$$

and

$$\dot{p}_J + \dot{p}_R = (1 - \kappa)\dot{p}_C.$$
 (5.4)

All metabolic processes are a function of V, E and parameters. The feeding and assimilation processes are also a function of X.

Parameter	Dimension	Interpretation
$\{\dot{p}_{Am}\}$	$e L^{-2} T^{-1}$	Surface-specific assimilation power
$[E_m]$	$e L^{-3}$	Maximum reserve density
$[\dot{p}_M]$	$e L^{-3} T^{-1}$	Volume-specific maintenance power
$\{\dot{p}_T\}$	$e L^{-2} T^{-1}$	Surface-specific maintenance power
$[E_G]$	$e L^{-3}$	Volume-specific growth costs
\dot{v}	LT^{-1}	Energy conductance
κ	-	Fraction of catabolic power spent on maintenance plus growth
κ_R	_	Fraction of reproduction power fixed in eggs
g	_	Investment ratio
\dot{k}_M	T^{-1}	Maintenance rate coefficient
\dot{k}_J	T^{-1}	Maturity rate coefficient
L_m	L	Maximum length
L_h	L	Heating length
E_H^b	e	Threshold of energy investment at birth
E_H^p	e	Threshold of energy investment at puberty
E_0	e	Energy cost of one egg
μ_E	$e M^{-1}$	Chemical potential
Compound specifier	Dimension	Interpretation
X	-	Substrate (food)
E	_	Reserve
V	_	Structure
P	-	Products
M_i	-	Mineral compound <i>i</i>
Process specifier	Dimension	Interpretation
Α	-	Assimilation
C	-	Catabolism
M	-	Maintenance (volume related)
T	-	Maintenance (surface related)
G	-	Growth
R	_	Reproduction or Maturation

Table 5.4: List of symbols. Dimensions: – no dimension; L length; M mass; T time; # moles or C-moles; e energy. Chemical compound and processes specifiers appear as subscripts to other variables. Symbols with (1) {·} are per unit surface area, (2) [·] are per unit of structural volume and (3) · are per unit time.

Empirical evidence (R4) shows that there are processes in the organism associated with food processing only, which suggests that food goes through a set of chemical reactions that transform it into reserves. Organisms spend energy on growth, maintenance and reproduction in the absence of food (F1, F2, F3) suggesting that the energy mobilized for growth, maintenance and reproduction is obtained from the reserve and not from the environment. The energy mobilized for maturation is also obtained from the reserve and not from the environment because eggs do not feed but they allocate energy to maturation. The allocation of energy to growth, somatic maintenance, maturation and reproduction covers the well-known metabolic processes common to all organisms. Maturity maintenance includes maintaining regulating mechanisms and concentration gradients and defense systems. The need to allocate energy to maturity maintenance is intimately related with the second law of thermodynamics (P2): the level of maturity would decrease in the absence of energy spent in its maintenance because the higher the level of maturity the further away is the system from equilibrium.

Empirical evidence (R1) shows that freshly laid eggs do not require maintenance because aerobic organisms use dioxygen in chemical reactions that yield energy. Thus, 1) freshly laid eggs do not have structure and 2) reserve does not require maintenance. Reserves can have active metabolic functions but do not require maintenance because they are continuously used passively and replenished.

The use of dioxygen increases with decreasing mass in an embryo (R2). This means that the structure that is built using the reserve requires maintenance. This is also coherent with the fact that respiration increases with mass in juveniles and adults (R2). Structural materials require maintenance because they are continuously degraded actively and reconstructed.

Proposition 5.3.1 (Partitionability of reserve dynamics) If the organism's reserve, E, is partitioned in the organism among categories of chemical compounds, $E_i \equiv \lambda_i E$, then 1) their constant relative abundances λ_i are constant and 2) the catabolic power mobilized from each aggregate must be proportional to the amount of energy embodied in it, i.e.,

$$\dot{p}_C(\lambda E, V) = \lambda \dot{p}_C(E, V), \tag{5.5}$$

with $0 \le \lambda \le 1$ and 3) the catabolic power mobilized to maintenance and growth, from each aggregate, must be proportional to the amount of energy embodied in it, i.e.,

$$\kappa(\lambda E, V)\dot{p}_C(\lambda E, V) = \kappa(E, V)\lambda\dot{p}_C(E, V).$$
(5.6)

Proof: The mobilization of the different categories of chemical compounds, E_i , must be coordinated, such that their constant relative abundances λ_i are constant, otherwise the chemical composition of reserves as a whole would change, violating Assumption 5.3.1.

Eq. 5.1 and the definition of $E_i \equiv \lambda_i E$ imply that $dE_i/dt = \dot{p}_A(E_i, V) - \dot{p}_C(E_i, V)$ and 2) $dE_i/dt = \lambda_i dE/dt = \lambda_i (\dot{p}_A(E, V) - \dot{p}_C(E, V))$. Eq. 5.5 follows from these two equations.

Also, the allocation to growth and maintenance of the different categories of chemical compounds, E_i , must be coordinated, such that growth and maintenance do not depend on the partitioning of the reserves (see Assumption 5.3.2), i.e,

$$\sum_{i=1}^{n} \kappa(E_i, V) \dot{p}_C(E_i, V) = \kappa(E, V) \dot{p}_C(E, V).$$
(5.7)

With Eq. 5.5, we can simplify Eq. 5.7:

$$\sum_{i=1}^{n} \lambda_i \kappa(\lambda_i E, V) = \kappa(E, V).$$
(5.8)

Eq. 5.8 implies that $\kappa(\lambda_i E, V) = \kappa(E, V)$ because $\sum_{i=1}^n \lambda_i = 1$. Eq. 5.6 follows. **Proposition 5.3.2 (Metabolic Stoichiometry)** The stoichiometry of each metabolic process defined in Assumption 5.3.2 is constant. In particular, the stoichiometries of somatic maintenance, maturity maintenance, maturation and the overhead of reproduction, $(1 - \kappa_R)\dot{p}_R$, are the same. For this reason, the sum of these powers is identified as the dissipation power

$$\dot{p}_D = \dot{p}_M + \dot{p}_J + \dot{p}_T + (1 - \kappa_R)\dot{p}_R, \tag{5.9}$$

where $\kappa_R = 0$ for the embryo and juvenile stages.

Also, the following conversion factors that characterize each metabolic process are constant. (1) $\eta_{*1*2} \equiv \mu_{*2*1}^{-1}$, between any mass flow *1 and any energy flow *2 and (2) y_{*1*2} , between any mass flows *1 and *2.

Proof: The stoichiometry of the assimilation process is (see Fig 5.1):

$$a_1M_1 + \ldots + a_xM_x + X \longrightarrow b_1P + b_2E, \tag{5.10}$$

where M_1 to M_x are the mineral compounds, b_1 and b_2 are the number of C-moles of product and reserve produced per each C-mol of food processed and a_1 to a_x are the net number of moles consumed of compounds M_1 to M_x , respectively. To obtain the x + 2stoichiometric coefficients a_1 to a_x , b_1 and b_2 we have to compute the mass balance for at least x + 2 chemical elements. These stoichiometric coefficients are constant because the chemical composition of X, E and P are also constant (see Assumption 5.3.1). Therefore, 1) the factor $y_{XE} = 1/b_2$ that converts food to reserve moles is constant, and 2) the factor $\eta_{XE} = 1/(b_2\mu_E)$ that converts food moles to reserve's energy is also constant because μ_E is constant (see Assumption 5.3.1). A similar reasoning can be applied to the other metabolic processes.

Also, the processes of maturity maintenace, somatic maintenance and the overhead of reproduction that consist in transforming reserve plus minerals into minerals have the same stoichiometry.

The overhead of reproduction is associated with the conversion from the reserve energy of the mother into the reserve energy of the embryo. The existence of an overhead is consistent with the dissipation principle (P2).

Proposition 5.3.3 (Organism Stoichiometry) The overall stoichiometry of the organism has 3 degrees of freedom. More specifically, the flow of any compound produced or consumed in the organism is a weighted average of the flows of any three other compounds.

Proof: The net flows (input-output) of minerals, products, heat and other compounds, at the boundary of the organism, are given by:

$$\dot{J}_{*1} = \dot{J}_{*1A} + \dot{J}_{*1D} + \dot{J}_{*1G}$$
(5.11)

where *1 stands for CO_2 , O_2 , heat, N_{waste} , H_2O and other compounds, \dot{J}_{*1A} , \dot{J}_{*1G} and \dot{J}_{*1D} are the net flows of *1 in the assimilation, growth and dissipation processes, respectively. Using the definition for η_{*1*2} (see Assumption 5.3.2):

$$J_{*1} = \eta_{*1A}\dot{p}_A + \eta_{*1D}\dot{p}_D + \eta_{*1G}\dot{p}_G \tag{5.12}$$

where $\dot{J}_{*1A} \equiv \eta_{*1A}\dot{p}_A$, $\dot{J}_{*1G} \equiv \eta_{*1G}\dot{p}_G$ and $\dot{J}_{*1D} \equiv \eta_{*1D}\dot{p}_D$.

To make this proof simpler we specify three compounds: CO_2 , O_2 and N_{waste} . We write Eq. 5.12 for each compound:

$$J_{CO_2} = \eta_{CO_2A}\dot{p}_A + \eta_{CO_2D}\dot{p}_D + \eta_{CO_2G}\dot{p}_G$$
(5.13)

$$\dot{J}_{O_2} = \eta_{O_2A} \dot{p}_A + \eta_{O_2D} \dot{p}_D + \eta_{O_2G} \dot{p}_G \tag{5.14}$$

$$\dot{J}_{N_{waste}} = \eta_{N_{waste}A}\dot{p}_A + \eta_{N_{waste}D}\dot{p}_D + \eta_{N_{waste}G}\dot{p}_G \tag{5.15}$$

By manipulating equations 5.13, 5.14 and 5.15, we obtain the following set of equations for the powers \dot{p}_A , \dot{p}_D and \dot{p}_G

$$\dot{p}_A = \alpha_{CO_2} \dot{J}_{CO_2} + \alpha_{O_2} \dot{J}_{O_2} + \alpha_{N_{waste}} \dot{J}_{N_{waste}},$$
(5.16)

$$\dot{p}_D = \beta_{CO_2} \dot{J}_{CO_2} + \beta_{O_2} \dot{J}_{O_2} + \beta_{N_{waste}} \dot{J}_{N_{waste}},$$
(5.17)

$$\dot{p}_G = \gamma_{CO_2} \dot{J}_{CO_2} + \gamma_{O_2} \dot{J}_{O_2} + \gamma_{N_{waste}} \dot{J}_{N_{waste}},$$
(5.18)

where α_i , β_i and γ_i are constants. Each power is a weighted average of the flows of 3 compounds. Thus, the flow of compound *1 (Eq. 5.12) can be written as a weighted average of 3 compounds if \dot{p}_A , \dot{p}_D and \dot{p}_G are replaced by equations 5.16, 5.17 and 5.18, respectively.

The method of indirect calorimetry (I1) is a particular case of Proposition 5.3.3. **Proposition 5.3.4 (Structure dynamics)** *Structure dynamics is controlled by growth,*

$$\frac{dV}{dt} = \frac{\dot{p}_G}{[E_G]},\tag{5.19}$$

where \dot{p}_G is the amount of energy allocated to growth and $[E_G]$ is the specific cost of growth.

Proof: The dynamics of structure is linked to growth directly by Assumption 5.3.2. The specific growth cost of structure is constant because $[E_G]$, a conversion factor between energy and mass, is constant (see Proposition 5.3.2).

Assumption 5.3.3 (Life-history events) Life stage events occur when E_H exceeds certain thresholds. The initiation of feeding occurs when $E_H = E_H^b$ and the initiation of allocation to reproduction coupled to the ceasing of maturation occurs when $E_H = E_H^p$. The dynamics of the amount of energy invested into maturation is:

$$\frac{dE_H}{dt} = \dot{p}_R, \qquad E_H < E_H^p. \tag{5.20}$$

Other life history events, such as cell division, metamorphosis or other stage transitions (e.g. to the pupal stage) occur also at threshold values for E_H .

The life history of organisms that reproduce by fission is well described by one single life-stage, the juvenile, while multicellular organisms typically start as embryo that does not feed and reproduce as adults. Whenever $E_H^p > E_H^b$ the three life stages are: the embryonic or foetal, the juvenile and the adult.

The logical reasoning behind this assumption is the following. An organism that develops and produces offspring increases its complexity (or maturity) from the embryo to the adult stage. Among organisms of the same species the level of maturity at the onset of feeding behaviour is the same. Also, initiation of allocation to reproduction, among organisms of the same species, occurs when they reach the same level (usually higher) of maturity. Therefore, it is reasonable to assume that the amount of energy invested to achieve the degrees of maturity that organisms need to start feeding or allocating to reproduction are intra-species constants.

Assumption 5.3.4 (Dependence on the Environment - Feeding) The ingestion at abundant food is proportional to surface area, $\dot{J}_{Xm} = {\{\dot{J}_{Xm}\}}V^{2/3}$, where ${\{\dot{J}_{Xm}\}}$ is the maximum surface-specific feeding rate. So, the feeding power is

$$\dot{p}_X = \{\dot{J}_{Xm}\} V^{2/3} f(X) \mu_X = \dot{J}_X \mu_X, \tag{5.21}$$

where $0 \le f(X) \le 1$, the nondimensional functional response, is

$$f(X) = \frac{\dot{J}_X}{\dot{J}_{Xm}},\tag{5.22}$$

 J_X is the rate of ingestion at food density X and μ_X is the chemical potential of food.

Feeding is proportional to surface area within the same species because acquisition processes and digestion and other food processing activities depend on mass transport processes that occur through surfaces (P3).

Proposition 5.3.5 (Dependence on the Environment - Assimilation) *The assimilation power is proportional to surface area,*

$$\dot{p}_A = \{\dot{p}_{Am}\}V^{2/3}f(X),$$
(5.23)

where $\{\dot{p}_{Am}\} \equiv \frac{\{\dot{J}_{Xm}\}}{\eta_{XA}}$ is the maximum surface-specific assimilation rate. **Proof:** By writing \dot{J}_X in Eq. 5.21 as a function of

$$\eta_{XA} \equiv \frac{p_A}{\dot{J}_X},\tag{5.24}$$

it follows directly that

$$\dot{p}_A = \frac{\{\dot{J}_{Xm}\}}{\eta_{XA}} V^{2/3} f(X).$$
(5.25)

Proposition 5.3.2 implies that η_{XA} is constant and therefore $\{\dot{p}_{Am}\} \equiv \frac{\{\dot{j}_{Xm}\}}{\eta_{XA}}$ is also constant.

Metabolic processes can be classified as supply-driven or demand-driven. Demanddriven processes are controlled by the state of the organism, i.e., by V and E, while supply-driven processes are controlled by inputs, e.g., by X. Feeding and assimilation are simultaneously supply and demand driven processes because they depend on the concentration of food in the environment, X and on the physiological state of the organism through the surface area (body size).

Assumption 5.3.5 (The κ allocation rule) *The* κ *function is a constant.*

Note that Assumption 5.3.5 only imposes that κ is also independent of V because in Proposition 5.3.1 we proved that κ is independent of E.

Good support for this assumption is the empirical evidence that many organisms do not stop growing after reproduction has started (G2) because it suggests that growth and reproduction are parallel processes that do not compete directly. Many species of holometabolic insects are an exception to this empirical evidence because they only start reproduction if they are (about) fully grown. In this case, reproduction cannot compete with growth.

Also, ovary and somatic cells have information on blood energy content (an intensive variable) but not on each other's activities (P4). Therefore, they take energy from the blood proportionally to their carriers membrane concentration, i.e., a fixed fraction of the energy content.

Assumption 5.3.6 (Weak homeostasis) For any constant food level, $X = X^* > 0$, there is a reserve density, $[E]^*(X^*)$, which remains constant along the growth process. Furthermore $\lim_{X^*\to\infty} [E]^* \equiv [E_m] < \infty$, where $[E_m]$ is the maximum reserve density.

The weak homeostasis assumption is supported by the empirical evidence that growing biomass converges to a constant chemical composition as long as food density remains constant (S2). Also, empirical evidence supports the existence of a maximum size (G4,G5) which implies that the maximum reserve density, $[E_m]$, is limited.

Proposition 5.3.6 (Catabolic power) Catabolic power is given by:

$$[\dot{p}_C] = [E] \left(\dot{v} V^{-1/3} - \frac{1}{V} \frac{dV}{dt} \right).$$
(5.26)

where $\dot{v} \equiv {\dot{p}_{Am}}/{[E_m]}$ is the energy conductance.

Proof:

By the definition of reserve density, i.e., $[E] \equiv E/V$:

$$\frac{d[E]}{dt} = \frac{1}{V}\frac{dE}{dt} - \frac{[E]}{V}\frac{dV}{dt}.$$
(5.27)

Replacing $\frac{dE}{dt}$ by Eq. 5.1 and \dot{p}_A by Eq. 5.23 Eq. 5.27 simplifies to:

$$\frac{d[E]}{dt} = \{\dot{p}_{Am}\}V^{-1/3}f(X) - [\dot{p}_C] - \frac{[E]}{V}\frac{dV}{dt}.$$
(5.28)

Eq. 5.28 can be written as,

$$\frac{d[E]}{dt} = \{\dot{p}_{Am}\}V^{-1/3}f(X) - \Phi(V, [E]),$$
(5.29)

because 1) \dot{p}_C is a function of E and V (see Assumption 5.3.2), 2) dV/dt is proportional to \dot{p}_G (see Eq. 5.19) and 3) \dot{p}_G is also a function of E and V (see Assumption 5.3.2).

If the organism is kept at constant food level for some time, weak homeostasis (Assumption 5.3.6) applies and Eq. 5.29 simplifies to

$$\{\dot{p}_{Am}\}V^{-1/3}f(X) = \Phi(V, [E]^*), \tag{5.30}$$

proving that $[E^*]$ is a function of resource density. Also, Assumption 5.3.6 says that $[E^*]$ is independ of volume because it remains constant along the growth process, implying that $\Phi(V, [E]^*) = V^{-1/3}H([E]^*)$. The function $\Phi(V, [E])$ can be generalized out of steady-state as $\Phi(V, [E]) = V^{-1/3}H([E]) + ([E^*] - [E])G(V, [E])$.

Using Eq. 5.28, Eq. 5.29 and $\Phi(V, [E]) = V^{-1/3}H([E]) + ([E^*] - [E])G(V, [E])$, the catabolic flux can be written as:

$$[\dot{p}_C] = V^{-1/3}H([E]) + ([E^*] - [E])G(V, [E]) - \frac{[E]}{V}\frac{dV}{dt}.$$
(5.31)

The catabolic power is a demand driven process, i.e., it cannot depend on X (see Assumption 5.3.2). However, Eq. 5.31 depends on $[E^*]$ and $[E^*]$ depends directly on X (see Eq. 5.30). This implies that G(V, [E]) = 0. Thus,

$$[\dot{p}_C] = V^{-1/3} H([E]) - \frac{[E]}{V} \frac{dV}{dt},$$
(5.32)

or

$$\dot{p}_C = V^{2/3} H([E]) - [E] \frac{dV}{dt}.$$
(5.33)

The partitionability Assumption 5.3.1 is used to specify H([E]). However, in the case of growing organisms, Eq. 5.5 is not enough, to specify H([E]), because dV/dt is an unknown function of (V, E). In the case of a full grown adult (dV/dt = 0), Eq. 5.5 implies that

$$\lambda H([E]) = H(\lambda[E]), \tag{5.34}$$

or $\lambda H([E]) = H(\lambda[E])$. Therefore $H([E]) = \dot{v}[E]$ and Eq. 5.32 simplifies to:

$$[\dot{p}_C] = \dot{v}[E]V^{-1/3} - [E]\frac{1}{V}\frac{dV}{dt}.$$
(5.35)

With Eq. 5.35 the reserve density dynamics is:

$$\frac{d[E]}{dt} = V^{-1/3} \left(\{ \dot{p}_{Am} \} f(X) - \dot{v}[E] \right).$$
(5.36)

Assumptions 5.3.4 and 5.3.6 impose that f(X) = 1 when $[E] = [E_m]$ implying that

$$\dot{v} = \frac{\{\dot{p}_{Am}\}}{[E_m]}.$$
 (5.37)

The mobilization of reserves (catabolic power) is a demand-driven process because it is independent of the environment (food) and it is dependent on V and E. This is reasonable because 1) it occurs inside the organism at a molecular level, and at that level no information concerning the external environment is available (P4), (2) this independence provides the organism an increased protection against environmental fluctuations and an increased control over its own metabolism (P6) and (3) it is uncoupled from the metabolic functions of feeding and assimilation. If metabolic functions were dependent on each other then it would be much more difficult to change a particular node in the metabolic network, while avoiding complex consequences for the whole organism. Also, changes in metabolic nodes typically occur "randomly" (blind watchmaker), and if several of them are required at the same time to improve the whole organism, such improvements would be rare. The result would be that evolutionary progress would stop, while the environment continues to change.

The catabolic power decreases with the relative structural growth because the energy density decreases (dilution by growth). The catabolic power per unit of structural volume of an organism that is not growing is

$$[\dot{p}_C] = \dot{v} \frac{[E]}{V^{1/3}}.$$
(5.38)

This is similar to a diffusion law where the flux of reserves per unit of structural volume is proportional to a reserve density gradient and to the energy conductance \dot{v} . Also, $[\dot{p}_C]$, is constant for a full grown organism at constant food level because dV/dt = 0 and $[E] = [E]^*$ (see Assumption 5.3.6). This means that, at constant food level, an organism that behaves accordingly to Assumption 5.3.6 and Proposition 5.3.6 has a higher degree of control over its metabolism (P6).

Proposition 5.3.7 (Reserve density at weak homeostasis) *The reserve density under weak homeostasis is a function of resource density only:*

$$[E^*] = f(X)[E_m].$$
(5.39)

Proof: At weak homeostasis $\frac{d[E]}{dt} = 0$ and Eq. 5.28 simplifies to:

$$[\dot{p}_C^*] = [E^*] \left(\dot{v} V^{-1/3} \frac{f(X^*)}{[E^*]/[E_m]} - \frac{1}{V} \frac{dV}{dt} \right).$$
(5.40)

Eq. 5.39 is now obtained inserting the expression for catabolic power given by Eq. 5.26.

Assumption 5.3.7 (Maintenance Powers) Maintenance costs \dot{p}_M and \dot{p}_T are proportional to structural volume and to surface area:

$$\dot{p}_M = [\dot{p}_M]V,\tag{5.41}$$

$$\dot{p}_T = \{\dot{p}_T\} V^{2/3},\tag{5.42}$$

where $[\dot{p}_M]$ and $\{\dot{p}_T\}$ are constant specific costs, i.e., costs per time and per unit of structural volume or area, respectively.

Maturity maintenance costs \dot{p}_J are proportional to the cumulative amount of energy invested into maturation,

$$\dot{p}_J = k_J E_H,\tag{5.43}$$

where $E_H \leq E_H^p$ and \dot{k}_J is a maturity rate coefficient.

Maintenance includes all metabolic processes that the organism must perform to 'stay alive'. Maintenance costs associated with size include costs that are proportional to the structural volume \dot{p}_M , e.g., protein turnover, and processes that are proportional to the surface area \dot{p}_T , e.g., heating in endotherms.

The chemical composition of the structure is constant (see Assumption 5.3.1) which means that each unit of structure needs the same type and amount of materials and amount of energy per unit of time, i.e., specific costs are constant.

The energy already invested in maturation, E_H , contributed to increase the amount of regulating mechanisms and gradients in the organism. Henceforth, the maturity maintenance costs, \dot{p}_J , increase proportional to E_H . In an adult the maturity maintenance costs are constant because maturity does not increase after the onset of reproduction.

Proposition 5.3.8 (Maturation and Reproductive Power) The amount of energy allocated to maturation in a juvenile is

$$\dot{p}_R = (1 - \kappa)\dot{p}_C - \dot{k}_J E_H,$$
(5.44)

while the amount of energy allocated to reproduction in an adult is

$$\dot{p}_R = (1 - \kappa)\dot{p}_C - \dot{k}_J E_H^p.$$
(5.45)

Proof: It follows from Eq. 5.4 and Eq. 5.43. ■

The amount of energy that an adult invests in reproduction is invested by a juvenile into maturation. If an organism is kept at a low food density such that the accumulated amount of energy invested into maturation never reaches the threshold E_H^p then the organism will never reproduce. The energy drain to maturity maintenance, \dot{p}_J , 1) decreases \dot{p}_R (see Eq. 5.44 and Eq. 5.45) 2) and increases the duration of the juvenile stage (see Eq. 5.20).

Proposition 5.3.9 (Allocation Priorities) *Maintenance has priority over growth and maturity maintenance has priority over maturation or reproduction.*

Proof: Maintenance powers, \dot{p}_M , \dot{p}_T and \dot{p}_J given by Eq. 5.41, Eq. 5.42 and Eq. 5.43 are set by the state of the organism V, by κ and other parameters that are constant. The energy that is not needed for maintenance purposes is then allocated to growth by Eq. 5.19 and to maturation or reproduction by Eq. 5.44 or Eq. 5.45.

Maintenance and maturity maintenance have priority because they are demand-driven processes, i.e., they are completely controlled by the state of the organism. Growth and maturation or reproduction are supply-driven processes because they do not depend on the state of the organism, i.e., they depend on the amount of energy not needed for maintenance and maturity maintenance, respectively.

Proposition 5.3.10 (Maximum Size) Organisms of the same species have a maximum size, V_m , i.e., organisms do not grow indefinitely.

Proof: Somatic maintenance competes directly and has priority over growth (see Prop. 5.3.9). This is crucial because somatic maintenance increases proportional to size (see Assumption 5.3.7), which imposes a maximum size on the organism. ■

In the literature, the existence of a maximum size (including reserve and structure) is generally accepted (G4,G5) which implies that the structure also has a maximum size, V_m .

Proposition 5.3.11 (Dynamic energy budget) *The dynamic energy budget of an organism is:*

$$\frac{de}{dt} = \dot{v}L^{-1}(f(X) - e)$$
(5.46)

$$\frac{dL}{dt} = \frac{\dot{v}\,e - L_h/L_m - L/L_m}{3 \, g + e}$$
(5.47)

where $e \equiv [E]/[E_m]$ is the scaled reserve density, $L \equiv V^{1/3}$ is the volumetric length,

$$g \equiv \frac{[E_G]}{\kappa[E_m]} \tag{5.48}$$

is the investment ratio, i.e., the ratio between the costs of growth and the maximum amount of energy allocated to growth and maintenance, $L_h \equiv {\dot{p}_T}/{\dot{p}_M}$ is the heating length and

$$L_m \equiv \frac{\kappa \{ \dot{p}_{Am} \}}{[\dot{p}_M]} = \frac{\dot{v}[E_G]}{g[\dot{p}_M]}$$
(5.49)

is the maximum length.

Proof: Eq. 5.46, is obtained by multiplying Eq. 5.30 by $[E_m]$ and using the definition of scaled reserve density.

Eq. 5.47 is derived by 1) replacing \dot{p}_G in Eq. 5.19 by Eq. 5.3, 2) replacing \dot{p}_C with Eq. 5.26 multiplied by V, 3) replacing \dot{p}_M and \dot{p}_T with Eq. 5.41 and Eq. 5.42, respectively, 4) rearranging to obtain

$$\frac{dV}{dt} = \frac{[E]\kappa \dot{v} V^{2/3} - [\dot{p}_M]V - \{\dot{p}_T\}V^{2/3}}{[E_G] + \kappa[E]},$$
(5.50)

and 5) replacing V with L^3 , [E] with $e[E_m]$, $\{\dot{p}_T\}$ with $L_h[\dot{p}_M]$, $\kappa[E_m]$ with $[E_G]/g$ and $[E_G]\dot{v}/g[\dot{p}_M]$ with L_m in Eq. 5.50.

The heating length is the reduction in length due to the energy allocated to surface maintenance costs. For endotherms, these surface maintenance costs are associated mainly with heating. If an organism has no surface maintenance costs, i.e., $\dot{p}_T = 0$ and $L_h = 0$, then its maximum volumetric length, L_m , achieved at e = 1, is given by Eq. 5.49 (see also Eq. 5.47).

The energy budget of the organism (Eq. 5.46 and 5.47) is determined by powers and not the reverse, which means that in special circumstances (extreme starvation), the energy budget can be different.

Proposition 5.3.12 (von Bertalanffy law) The growth curve of an isomorphic individual at constant food availability X^* or at abundant food ($f \approx 1$) is:

$$\frac{dL}{dt} = \dot{r}_B (L_\infty - L). \tag{5.51}$$

The ultimate length L_{∞} and the von Bertalanffy growth rate \dot{r}_B are given by

$$L_{\infty} = \left(fL_m - L_h\right),\tag{5.52}$$

$$\dot{r}_B = \frac{\dot{v}}{3L_m} \frac{1}{g+f} = \frac{\dot{k}_M g}{3(g+f)} = \left(\frac{3}{\dot{k}_M} + \frac{3L_h}{\dot{v}} + \frac{3L_\infty}{\dot{v}}\right)^{-1},$$
(5.53)

where $k_M \equiv [\dot{p}_M]/[E_G]$ is the maintenance rate coefficient, i.e., the ratio between the costs of maintenance and growth of structure.

Proof: If resource density is constant, X^* , reserve density is $e^* \equiv [E^*]/[E_m] = f(X^*)$ through most of the individual's life (see Eq. 5.39). Hence, the growth curve (see Eq. 5.47) is:

$$\frac{dL}{dt} = \frac{\dot{v}}{3} \frac{f - L_h/L_m - L/L_m}{g + f}.$$
(5.54)

Eq. 5.51 is obtained by combining Eq. 5.54 with Eq. 5.52 and Eq. 5.53. ■

Von Bertalanffy's law (Eq. 5.51) is one of the most universal biological patterns (G1). Also, organisms of the same species at different food availabilities exhibit von Bertalanffy growth rates that are inversely proportional to ultimate length (G5) (Eq. 5.53).

Proposition 5.3.13 (Foetal Development) If the reserves of the mother, continuously supplied to the foetus via the placenta, are considered very large, i.e., $e = \infty$ then foetal growth is given by

$$\frac{dL}{dt} = \frac{\dot{v}}{3}.\tag{5.55}$$

Proof: Eq. 5.55 is obtained dividing the numerator and the denominator of the right hand side of Eq. 5.47 by e and then replacing it with ∞ .

According with Eq. 5.55 the structural volume of the foetus is proportional to cubed time:

$$V(t) = \left(\frac{\dot{v}t}{3}\right)^3.$$
(5.56)

The structural volume can be converted to weight using the following auxiliary proposition.

Proposition 5.3.14 (Volume to weight) *The relationship between weight, w, and structural volume, V, is*

$$w = \left(\frac{[E]}{\mu_E} + d_V\right) V = \left(\frac{[E_m]}{\mu_E}e + d_V\right) V,$$
(5.57)

where d_V is the density of the structure and μ_E is the chemical potential of reserve.

Proof: The volume of the organism can be written as,

$$\frac{[E]V}{d_E\mu_E} + V,\tag{5.58}$$

where the first term is the volume of the reserve, i.e., the ratio between the reserve's energy, [E]V, and the reserve's energy per unit volume of reserve, $d_E\mu_E$, and d_E is the density of the reserve. The weight of the organism (Eq. 5.57) is obtained by multiplying the volume of the reserve by d_E and the volume of the structure by d_V .

The structural volume of the foetus can be converted to weight (Eq. 5.57) using a constant because he has unlimited access to food, i.e., the reserves of the mother. In this case, Eq. 5.56 is validated by the empirical data that suggests that foetal weight is proportional to cubed time (G3).

Proposition 5.3.15 (Intraspecific Kleiber's Law) The metabolic rate measured by the dioxygen consumption of animals of the same species, \dot{J}_{O_2} kept under fasting conditions,

for a short time, is proportional to w^{α} with $\alpha \in [0.66, 1]$. If animals have the same reserve density *e* then the proportionality constant is the same.

Proof: For organisms that are kept under fasting conditions, dioxygen is consumed in growth, reproduction or maturation and maintenance, i.e.,

$$\dot{J}_{O_2} = \eta_{OD}\dot{p}_M + \eta_{OD}\dot{p}_J + \eta_{OD}\dot{p}_T + (1 - \kappa_R)\eta_{OD}\dot{p}_R + \eta_{OG}\dot{p}_G,$$
(5.59)

where η_{OD} and η_{OG} are constant because the stoichiometry of each process is constant (see Proposition 5.3.2).

Eq. 5.59 can be simplified to:

$$\dot{J}_{O_2} = \eta_{OD} \frac{1 - \kappa_R + \kappa_R \kappa}{\kappa} (\dot{p}_M + \dot{p}_T + \dot{p}_G) + \eta_{OD} \kappa_R \dot{p}_J$$
(5.60)

by first inserting the expression for \dot{p}_R given by Eq. 5.4 and then replacing \dot{p}_C with Eq. 5.3.

The somatic and maturity maintenance powers are given by Eq. 5.41, 5.42 and 5.43 and the growth power,

$$\dot{p}_G = [E_G] \dot{v} \frac{(e - L_h/L_m)V^{2/3} - V/L_m}{g + e},$$
(5.61)

is obtained by combining Eq. 5.19 with Eq. 5.47. The powers \dot{p}_M and \dot{p}_J are proportional to V while \dot{p}_T is proportional to $V^{2/3}$ and \dot{p}_G is a linear combination of $V^{2/3}$ and V. The dioxygen consumption must be approximately proportional to V^{α} with $\alpha \in [2/3, 1]$ because it is a linear combination of $V^{2/3}$ and V. If the animals of the same species have a similar reserve density then the proportionality constant between \dot{J}_{O_2} and w^{α} is the same (see Eq. 5.57).

Empirical evidence on Kleiber's law is amply available in the literature (R3). However, if organisms of the same species do not have similar reserve densities then the available data in the literature is harder to evaluate because respiration is typically linked to body weight, which has contributions from both reserve and structure. DEB Proposition 5.3.15 provides an explanation for the differences in the values of α in Kleiber's law within the same species: differences in the allocation of energy to growth and surface related maintenance.

5.4 Theory on Parameter Values

In DEB theory, the set of parameter values is individual-specific. Individuals differ in parameter values and selection leads to evolution characterized by a change in the (mean)

value of these parameters (P7). The differences between species is just an evolutionary amplification of the difference between individuals, i.e., it reduces to differences in the mean value of DEB parameters. In this section, the theory for the covariation of (mean) parameter values among species is presented.

Assumption 5.4.1 (Primary molecular based parameters) Molecular based parameters are related to molecular processes and, for related species, are similar and independent of the ultimate size of the organism (see Tab. 5.5). These parameters include: $[E_G], [\dot{p}_M], {\dot{p}_T}, \dot{k}_J, \kappa, \kappa_R, \dot{v}, [E_H^b] = \frac{E_H^b}{V_m} and [E_H^p] = \frac{E_H^p}{V_m}.$

Parameters that characterize molecular based processes (see Tab. 5.5) are considered to be identical for related species because cells are very similar independently of the size of the organism (C1). Therefore cells of about equal size have similar growth, maintenance and maturation costs, i.e., $[E_G]$, $[\dot{p}_M]$, $\{\dot{p}_T\}$, \dot{k}_J , κ_R , $[E_H^b]$ and $[E_H^p]$ are equal for related species. The partition of energy mobilized from reserves is done at the level of the somatic and reproductive cells and therefore κ is also a molecular based process. Kooijman and Troost [46] present a possible molecular mechanism that makes clear that \dot{v} is a molecular based parameter. A simpler but less precise argument to justify this is presented next. Two full grown organisms with the same V and the same [E], that belong to different but related species with different maximum lengths, have similar metabolic needs. Therefore they must have a similar rate of mobilization of reserves. This occurs if \dot{v} is the same because the mobilization of reserves is proportional to \dot{v} (see Eq. 5.26).

Proposition 5.4.1 (Secondary molecular based parameters) *The heating length* L_h *is similar between related species and independent of the ultimate size of the organism.*

Proof: The heating length L_h is independent of the size of the organism because both $\{\dot{p}_T\}$ and $[\dot{p}_M]$ are molecular based parameters and $L_h = \frac{\{\dot{p}_T\}}{[\dot{p}_M]}$.

Proposition 5.4.2 (Physical design parameters) *Physical design parameters are proportional to the maximum length of the organism,* L_m (see Tab. 5.5). These parameters in*clude:* $\{\dot{p}_{Am}\}, [E_m], E_H^p, E_H^b$ and g.

Proof: Suppose that a reference species and species A are related and that $z = \frac{L_m^A}{L_m}$ is the ratio of maximum lengths.

The surface specific assimilation rate is proportional to maximum length (see Eq. 5.49) because the molecular based parameters are equal for both species, $\kappa = \kappa^A$ and $[\dot{p}_M] = [\dot{p}_M^A]$:

$$z = \frac{L_m^A}{L_m} = \frac{\{\dot{p}_{Am}^A\}}{\{\dot{p}_{Am}\}}.$$
(5.62)

Molecular Based Parameters	Physical Design Parameters
$\kappa^A = \kappa$	$\{\dot{p}^A_{Am}\} = \{\dot{p}_{Am}\}z$
$\dot{v}^A = \dot{v}$	$g = zg^A$
$\kappa_R^A = \kappa_R$	
$\dot{k}_J^A = \dot{k}_J$	
$[E_G^A] = [E_G]$	$[E_m^A] = [E_m]z$
$[\dot{p}_M^A] = [\dot{p}_M]$	
$\{\dot{p}_T^A\} = \{\dot{p}_T\}$	
$[E_b^{H,A}] = [E_b^H]$	$E_b^{H,A} = E_b^H$
$[E_p^{H,A}] = [E_p^H]$	$E_p^{H,A} = E_p^H$
$L_h^A = L_h$	

Table 5.5: Body Scaling Relations between different species with different maximum body sizes of L_m and L_m^A where $z = \frac{L_m^A}{L_m}$.

The maximum reserve density $[E_m]$ is proportional to maximum length, i.e., $[E_m^A] = z[E_m]$ because $\dot{v} = \dot{v}^A$ and $\{\dot{p}_{Am}^A\} = z\{\dot{p}_{Am}\}$ (see Eq. 5.37).

The energy investment threshold at puberty E_p^H is proportional to maximum length, i.e., $E_p^{H,A} = z^3 E_p^H$ because $[E_p^{H,A}] = [E_p^H]$. Also, the energy investment threshold at birth E_b^H is proportional to maximum length, i.e., $E_b^{H,A} = z^3 E_b^H$ because $[E_b^{H,A}] = [E_b^H]$.

The investment ratio g is proportional to maximum length, i.e., $g = zg^A$ because $[E_G] = [E_G]_A$, $\kappa = \kappa^A$ and $[E_m^A] = z[E_m]$ (see Eq. 5.48).

Proposition 5.4.3 (von Bertalanffy growth rate) *The von Bertalanffy growth rate for species A at abundant food is:*

$$\log \dot{r}_B^A = \log \frac{\dot{v}}{3} - \log(gL_m + L_m^A).$$
(5.63)

where all parameters with the exception of L_m^A are for a reference species.

Proof: At abundant food, i.e., f = 1, the von Bertalanffy growth rate (Eq. 5.53) for species A is:

$$\dot{r}_B^A = \frac{\dot{v}^A}{3L_m^A} \frac{1}{g^A + 1}.$$
(5.64)

Eq. 5.63 is obtained by (1) rewriting the DEB parameters of species A as a function of the DEB parameters of the reference species using Tab. 5.5 and then (2) applying logarithms.

This proposition on the inter-species comparison of Von Bertalanffy growth rate corrected for a common body temperature is supported by empirical data (G5)(for a comparison between empirical data and DEB model predictions see [41, Fig. 8.3]).

Proposition 5.4.4 (Interspecific Kleiber's Law) The metabolic rate measured by the dioxygen consumption of full grown adult animals kept under fasting conditions, of species of different maximum body size, scales to the α power of its mass, i.e., \dot{J}_{O_2} is proportional to w^{α} with $\alpha \in [0.5, 1]$.

Proof: For full grown organisms that are kept under fasting conditions, the dioxygen is consumed in reproduction and somatic and maturity maintenance, i.e.,

$$J_{O_2} = \eta_{OD} \dot{p}_M + \eta_{OD} \dot{p}_J + \eta_{OD} \dot{p}_T + (1 - \kappa_R) \eta_{OD} \dot{p}_R$$
(5.65)

where η_{OD} is constant because the stoichiometry of each process is constant (see Proposition 5.3.2).

Eq. 5.65 can be simplified to Eq. 5.60 where $\dot{p}_G = 0$ by first inserting the expression for \dot{p}_R given by Eq. 5.4 and then replacing \dot{p}_C with Eq. 5.3.

The somatic and maturity maintenance powers can be written for full grown adults $(l = 1 \text{ and } \min(E_H, E_H^p) = E_H^p)$:

$$\dot{p}_M = [\dot{p}_M] V_m^A, \tag{5.66}$$

$$\dot{p}_T = \{\dot{p}_T\} V_m^{A\,2/3},$$
(5.67)

$$\dot{p}_J = \dot{k}_J [E_H^p] V_m^A,$$
 (5.68)

where all parameters are for the reference species with the exception of the maximum body size of the species being considered V_m^A . Eq. 5.66 to 5.68 were obtained from Eq.5.41, 5.42 and 5.43 respectively, by first replacing body size V with V_m because l = 1 and then rewriting DEB parameters of the species being considered, species A, as a function of the DEB parameters of a reference species using Tab. 5.5 with the exception of the maximum body size, V_m^A .

The powers \dot{p}_M and \dot{p}_J are proportional to maximum body size volume V_m^A while \dot{p}_T is proportional to $V_m^{A\,2/3}$. The dioxygen consumption must be approximately proportional to $V_m^{A\,\alpha}$ with $\alpha \in [2/3, 1]$ because it is a linear combination of $V_m^{A\,2/3}$ and V_m^A .

The relationship between the weigth and the volume of the species considered,

$$w = \frac{[E_m]e}{\mu_E V_m^{1/3}} V_m^{A\,4/3} + d_V V_m^A,\tag{5.69}$$

is obtained from Eq. 5.57 by replacing body size V with V_m , energy density [E] with $[E_m]e$ and rewriting the parameters of species A as a function of the parameters of a reference species with the exception of V_m^A . Thus, (1) the mass is approximately proportional to V_m^β with $\beta \in [1, 4/3]$ and (2) the dioxygen consumption is approximately proportional to $w^{\alpha/\beta}$ with $\alpha/\beta \in [0.5, 1]$.

The proposition obtained that the power in Kleiber's law can be in the interval [0.5, 1] is more adequate that a unique value of 3/4. This is supported by Dodds et al. [12] reanalyses of datasets. These authors tested whether the power is 3/4 or 2/3 finding little evidence for rejecting the power 2/3. Also, Vidal and Whitledge [86] found powers of 0.72 and 0.85 for crustaceans and Phillipson [63] found values of 0.66 for unicellulars and 0.88 for ectotherms.

According to Proposition 5.4.4, the proportionality between dioxygen production and weight, in interspecies comparisons of full grown adults, has a variable power because of a variable allocation to surface related maintenance costs, e.g., heating, \dot{p}_T .

5.5 Conclusions

Syntheses of DEB theory and discussions of the underlying concepts have already been presented in the literature [60, 47, 81, 82]. However, in this chapter, we formalize DEB in a set of assumptions and propositions proving 1) that DEB is a theory for the metabolic organization of organisms fully supported by the empirical biological patterns and the universal laws of physics and evolution and 2) a theory for metabolic organization can be as formal as physics.

This theory considers that body mass is partitioned into structure and reserves which is supported by empirical evidence that organisms can have a variable stoichiometry (S1). Reserve does not require maintenance because it is passively continuously used and replenished while structure requires maintenance because it is actively continuously degraded and reconstructed. These two metabolic facts are supported by the fact that freshly laid eggs do not use dioxygen in significant amounts and that the use of dioxygen increases with decreasing mass in the embryo and with increasing mass in the juvenile and adult (R1, R2).

Feeding is considered to be proportional to surface area within a species because transport occurs across surfaces (P3). In the organism (1) food is transformed into reserve and (2) reserve is mobilized to fuel growth, maturation, maintenance and reproduction. This internal organization is suggested by the empirical evidence on the heat increment of feeding (R4) and by the fact that starving organisms survive, grow and reproduce (F1-F3).

Additionally to the processes of growth, maturation, maintenance and reproductions, organisms also allocate energy to maturity maintenance, which is imposed by the need to spend energy to keep the organism far away from equilibrium (P2). The allocations of energy to repoduction and growth do not compete with each other; this is suggested by empirical evidence on growth (G2) and the laws of mass and energy transfer (P4). Also, energy allocation to growth competes directly with the energy allocation to somatic maintenance, which imposes a maximum size within a species (G4, G5).

Organisms tend to a constant chemical composition in an environment with constant food availability; this is supported by empirical evidence on a constant stoichiometry under certain conditions (S2) and motivated by evolutionary theory (P6).

The theory for the covariation of parameter values among species is based on the empirical evidence that cells are similar across species (C1).

The propositions obtained explain the following empirical findings: (1) the method of indirect calorimetry (I1), (2) body size growth (von Bertalanffy curves), (3) the variation of von Bertalanffy's growth rates within (G5) and across species (G4), (4) Kleiber's law on metabolic rate (R3) and (5) the pattern of foetal growth (G3).

These assumptions and propositions are at the core of DEB theory. This theory has already been tested for many organisms including bacteria [4, 5, 43, 79], crustaceans [40], collemboles [33], appendicularians [41], rotifiers [2], yeasts [20], molluscs [21, 24, 25, 26, 83], unicellular alga [41], lichens [41], nematodes [1, 22, 23, 32], fishes [84, 85] and birds and mammals [41, 89].

In this chapter, we focus on the standard DEB model for isomorphs with one reserve and one structure. They are ideal to explain the concepts, and demonstrate the importance surface area - volume interactions, which is an important organising principle, in combination with mass and energy conservation. However, from an evolutionary perspective they represent an advanced state that evolved from systems with more reserves and, therefore, less homeostatic control. The evolution of metabolism as a dynamic system is discussed in [46]. Extensions to the standard DEB model that were not discussed in this chapter include: (1) shape corrections for the surface area of organisms that do not behave as isomorphs but deviate from this in predictable ways [41, p.26-29]; (2) the dependence of physiological rates on body temperature [41]; (3) the inclusion of more reserves (for organisms feeding on simple substrates) and more structures (plants) [41, p.168]; (4) an ageing model that explains the phenomenological Weibull [41, p.141] and the Gompertz laws [54]; (5) shrinking whenever the catabolic power mobilized from reserves is not enough to pay maintenance [80] and (6) implications for cellular levels [45], trophic chains and population dynamics [39, 50, 51, 52, 58] and ecosystem dynamics [42, 44, 61].

This chapter contributes to a research agenda on a non-species specific metabolic theory by (1) proposing a set of stylized empirical patterns that are the ultimate test to any metabolic theory and (2) using these facts to establish a set of assumptions and obtaining the propositions that follow. The validity of each assumption and empirical fact considered can be independently discussed leading to a wider consensus in the metabolic field.

References

- A. O. Alvarez, T. Jager, S. A. L. M. Kooijman, and J Kammenga. Responses to stress of *Caenorhabditis elegans* populations with different reproductive strategies. *Functional Ecology*, 19:656–664, 2005.
- [2] M. O. Alver, J. A. Alfredsen, and Y. Olsen. An individual-based population model for rotifier (*Brachionus plicatilis*) cultures. *Hydrobiologia*, 560:93–108, 2006.
- [3] L. von Bertalanffy. A quantitative theory of organic growth (Inquiries on growth laws. II.). *Human Biology*, 10:181–213, 1938.
- [4] B. W. Brandt, F. D. L. Kelpin, I. M. M. van Leeuwen, and S. A. L. M. Kooijman. Modelling microbial adaptation to changing availability of substrates. *Water Research*, 38:1003–1013, 2004.
- [5] B. W. Brandt, I. M. M. van Leeuwen, and S. A. L. M. Kooijman. A general model for multiple substrate biodegradation. Application to co-metabolism of non structurally analogous compounds. *Water Research*, 37:4843–4854, 2003.
- [6] T. L. Bucher. Parrot eggs, embryos, and nestlings patterns and energetics of growth and development. *Physiological Zoology*, 56:465–483, 1983.
- [7] M. A. Chappell, G. C. Bachman, and K. A Hammond. The heat increment of feeding in house wren chicks: Magnitude, duration, and substitution for thermostatic costs. *Journal of Comparative Physiology B-Biochemical Systemic and Environmental Physiology*, 167:313–318, 1997.

- [8] Y. Chen, D. A. Jackson, and H. H. Harvey. A comparison of von Bertalanffy and polynomial functions in modeling fish growth data. *Canadian Journal of Fisheries* and Aquatic Sciences, 49:1228–1235, 1992.
- [9] Y. Chen, C. H. Ke, S. Q. Zhou, and F. X. Li. Effects of food availability on feeding and growth of cultivated juvenile *Babylonia formosae habei* (Altena and Gittenberger 1981). *Aquaculture Research*, 36:94–99, 2005.
- [10] Y. Chilliard, C. Delavaud, and M. Bonnet. Leptin expression in ruminants: Nutritional and physiological regulations in relation with energy metabolism. *Domestic Animal Endocrinology*, 29(1):3–22, 2005.
- [11] A. Clarke and N. M. Johnston. Scaling of metabolic rate with body mass and temperature in teleost fish. *Journal of Animal Ecology*, 68:893–905, 1999.
- [12] P. S. Dodds, D. H. Rothman, and J. S. Weitz. Re-examination of the '3/4-law' of metabolism. *Journal of Theoretical Biology*, 209:9–27, 2001.
- [13] S. Dou, R. Masuda, M. Tanaka, and K. Tsukamoto. Feeding resumption, morphological changes and mortality during starvation in japanese flounder larvae. *Journal* of Fish Biology, 60:1363–1380, 2002.
- [14] S. B. Du and K. S. Mai. Effects of starvation on energy reserves in young juveniles of abalone *Haliotis discus hannai Ino. Journal of Shellfish Research*, 23:1037–1039, 2004.
- [15] B. P. Ferreira and G. R. Russ. Age validation and estimation of growth-rate of the coral trout, *Plectropomus-leopardus*, (Lacepede 1802) from Lizard Island, Northern Great Barrier Reef. *Fishery Bulletin*, 92:46–57, 1994.
- [16] P. Fink, L. Peters, and E. Von Elert. Stoichiometric mismatch between litoral invertebrates and their periphyton food. *Archiv Fur Hydrobiologie*, 165(2):145–165, 2006.
- [17] N. B. Frazer, J. W. Gibbons, and J. L. Greener. Exploring fabens growth interval model with data on a long-lived vertebrate, *Trachemys-scripta* (reptilia, testudinata). *Copeia*, 1:112–118, 1990.

- [18] C. S. Gallardo, C. Manque, and M. Filun. Comparative resistance to starvation among early juveniles of some marine muricoidean snails. *Nautilus*, pages 121– 126, 2004.
- [19] V.F. Galluci and T.J. Quinn. Reparameterizing, fitting, and testing a simple growthmodel. *Transactions of the American Fisheries Society*, 108:14–25, 1979.
- [20] P. P. F. Hanegraaf, A. H. Stouthamer, and S. A. L. M. Kooijman. A mathematical model for yeast respiro-fermentative physiology. *Yeast*, 16:423–437, 2000.
- [21] R. J. F. van Haren. Application of Dynamic Energy Budgets to xenobiotic kinetics in Mytilus edulis and population dynamics of Globodera pallida. PhD thesis, Vrije Universiteit, June 1995.
- [22] R. J. F. van Haren and E. M. L. Hendrikx. Population dynamic models of potato cyst nematodes based on dynamic energy budgets. *Nematologica*, 38:414, 1992.
- [23] R. J. F. van Haren and E. M. L. Hendrikx. Feeding, growth and reproduction of potato cyst nematodes. *Nematologica*, 41:351, 1995.
- [24] R. J. F. van Haren, E. M. L. Hendrix, and H. J. Atkinson. Growth curve analysis of sedentary plant parasitic nematodes in relation to plant resistance and tolerance. In *Predictability and Nonlinear Modelling in Natural Sciences and Economics*, (eds Grasman, J. and Straten, G. van), pages 172–183. Kluwer Academic Publishers, Dordrecht, 1994.
- [25] R. J. F. van Haren and S. A. L. M. Kooijman. Application of the dynamic energy budget model to *Mytilus edulis* (L). *Netherlands Journal of Sea Research*, 31:119– 133, 1993.
- [26] R. J. F. van Haren, H. E. Schepers, and S. A. L. M. Kooijman. Dynamic energy budgets affect kinetics of xenobiotics in the marine mussel *Mytilus edulis*. *Chemosphere*, 29:163–189, 1994.
- [27] P.A.J. Hawkins, P.J. Butler, A.J. Woakes, and G.W. Gabrielsen. Heat increment of feeding in Brunnich's guillemot *Uria lomvia*. *Journal of Experimental Biology*, 200:1757–1763, 1997.

- [28] M. Heino and V. Kaitala. Evolution of resource allocation between growth and reproduction in animals with indeterminate growth. *Journal of Evolutionary Biology*, 12:423–429, 1999.
- [29] H.J. Hirche and G. Kattner. Egg production and lipid content of *Calanus glacialis* in spring: indication of a food-dependent and food-independent reproductive mode. *Marine Biology*, 117:615–622, 1993.
- [30] A. S. Huggett and W. F. Widdas. The relationship between mammalian foetal weight and conception age. *The Journal of Physiology*, 114:306–317, 1951.
- [31] Y. Ingenbleek. The nutritional relationship linking sulfur to nitrogen in living organisms. *The Journal of Nutrition*, 136:1641–1651, 2006.
- [32] T. Jager, O. A. Alvarez, J. E. Kammenga, and S. A. L. M Kooijman. Modelling nematode life cycles using dynamic energy budgets. *Functional Ecology*, 19:136– 144, 2005.
- [33] T. Jager, T. Crommentuijn, C. A. M. van Gestel, and S. A. L. M. Kooijman. Simultaneous modelling of multiple endpoints in life-cycle toxicity tests. *Environmental Science and Technology*, 38:2894–2900, 2004.
- [34] D. N. Janes and M. A. Chappell. The effect of ration size and body-size on specific dynamic action in adelie penguin chicks, *Pygoscelis-adeliae*. *Physiological Zoology*, 68:1029–1044, 1995.
- [35] C. Jorgensen and O. Fiksen. State-dependent energy allocation in cod (*Gadus morhua*). *Canadian Journal of Fisheries and Aquatic Sciences*, 63:186–199, 2006.
- [36] K. L. Kirk. Life-history responses to variable environments: Starvation and reproduction in planktonic rotifers. *Ecology*, 78:434–441, 1997.
- [37] O. S. Kjesbu, J. Klungsoyr, H. Kryvi, P. R. Witthames, and M. G. Walker. Fecundity, atresia, and egg size of captive atlantic cod (*gadus-morhua*) in relation to proximate body-composition. *Canadian Journal of Fisheries and Aquatic Sciences*, 48:2333– 2343, 1991.
- [38] M. Kleiber. Body size and metabolism. *Hilgardia*, 6:315–353, 1932.

- [39] B. W. Kooi, L. D. J. Kuijper, and S. A. L. M. Kooijman. Consequences of symbiosis for food web dynamics. *Journal of Mathematical Biology*, 3:227–271, 2004.
- [40] S. A. L. M. Kooijman. Population dynamics on the basis of budgets. In J. A. J. Metz and O. Diekmann, editors, *The dynamics of physiologically structured populations*, Springer Lecture Notes in Biomathematics, pages 266–297. Springer-Verlag, Berlin, 1986.
- [41] S. A. L. M. Kooijman. Dynamic Energy and Mass Budgets in Biological Systems. Cambridge University Press, Cambridge, 2000.
- [42] S. A. L. M. Kooijman, H. A. Dijkstra, and B. W. Kooi. Light-induced mass turnover in a mono-species community of mixotrophs. *Journal of Theoretical Biology*, 214:233–254, 2002.
- [43] S. A. L. M. Kooijman, E. B. Muller, and A. H. Stouthamer. Microbial-growth dynamics on the basis of individual budgets. *Antonie Van Leeuwenhoek International Journal of General and Molecular Microbiology*, 60:159–174, 1991.
- [44] S. A. L. M. Kooijman and R. M. Nisbet. How light and nutrients affect life in a closed bottle. In S. E. Jørgensen, editor, *Thermodynamics and ecological modelling*, pages 19–60. CRC Publ., Boca Raton, FL, USA, 2000.
- [45] S. A. L. M. Kooijman and L. A. Segel. How growth affects the fate of cellular metabolites. *Bulletin of Mathematical Biology*, 67:57–77, 2005.
- [46] S. A. L. M. Kooijman and T. A. Troost. Quantitative steps in the evolution of metabolic organisation as specified by the dynamic energy budget theory. *Biological Reviews (in press)*, 2006.
- [47] S.A.L.M. Kooijman. Quantitative aspects of metabolic organization: a discussion of concepts. *Philosophical Transactions of the Royal Society of London B*, 356:331– 349, 2001.
- [48] J. Kozlowski. Optimal allocation of resources explains interspecific life-history patterns in animals with indeterminate growth. *Proceedings of the Royal Society of London Series B-Biological Sciences*, 263:559–566, 1996.
- [49] E. Krol, P. Redman, P. J. Thomson, R. Williams, C. Mayer, J. G. Mercer, and J. R. Speakman. Effect of photoperiod on body mass, food intake and body composition

in the field vole, *Microtus agrestis*. *Journal of Experimental Biology*, 208:571–584, 2005.

- [50] L. D. Kuijper, B. W. Kooi, T. R. Anderson, and S. A. L. M. Kooijman. Stoichiometry and food chain dynamics. *Theoretical Population Biology*, 66:323–339, 2004.
- [51] L. D. J. Kuijper, T. R. Anderson, and S. A. L. M. Kooijman. C and N gross efficiencies of copepod egg production studies using a Dynamic Energy Budget model. *Journal of Plankton Research*, 26:213–226, 2004.
- [52] L. D. J. Kuijper, B. W. Kooi, C. Zonneveld, and Kooijman S. A. L. M. Omnivory and food web dynamics. *Ecological Modelling*, 163:19–32, 2003.
- [53] E. R. S. Kunji, T. Ubbink, A. Matin, B. Poolman, and W. N. Konings. Physiologicalresponses of lactococcus-lactis ml³ to alternating conditions of growth and starvation. *Archives of Microbiology*, 159:372–379, 1993.
- [54] I. M. M. Leeuwen, F. D. L. Kelpin, and S. A. L. M. Kooijman. A mathematical model that accounts for the effects of caloric restriction on body weight and longevity. *Biogerontology*, 3:373–381, 2002.
- [55] B. H. Letcher, J.A. Rice, L. B. Crowder, and F. P. Binkowski. Size-dependent effects of continuous and intermittent feeding on starvation time and mass loss in starving yellow perch larvae and juveniles. *Transactions of the American Fisheries Society*, (125):14–26, 1996.
- [56] T. Molnar, A. Szabo, G. Szabo, C. Szabo, and C. Hancz. Effect of different dietary fat content and fat type on the growth and body composition of intensively reared pikeperch *Sander lucioperca* (1.). *Aquaculture Nutrition*, 12:173–182, 2006.
- [57] H. J. Morowitz. Energy Flow in Biology. Academic Press, New York, 1968.
- [58] E. B. Muller, R. M. Nisbet, S. A. L. M. Kooijman, J. J. Elser, and E. McCauley. Stoichiometric food quality and herbivore dynamics. *Ecol. Letters*, 4:519–529, 2001.
- [59] R. F. Nespolo, L. E. Castaneda, and D. A. Roff. The effect of fasting on activity and resting metabolism in the sand cricket, *Gryllus firmus*: a multivariate approach. *Journal of Insect Physiology*, 51:61–66, 2005.

- [60] R. Nisbet, E. Muller, K. Lika, and S.A.L.M. Kooijman. From molecules to ecosystems through dynamic energy budget models. *Journal of Animal Ecology*, 69:913– 926, 2000.
- [61] A. W. Omta, J. Bruggeman, S. A. L. M. Kooijman, and H. A. Dijkstra. The biological carbon pump revisited: feedback mechanisms between climate and the Redfield ratio. *Geophysical Research Letters*, 2006.
- [62] T. N. Pettit. Embryonic oxygen-consumption and growth of Laysan and blackfooted albatross. *American Journal of Physiology*, 242:121–128, 1982.
- [63] J. Phillipson. Bioenergetic options and phylogeny. In *Physiological Ecology: An Evolutionary Approach to Resource Use*, (eds. C. R. Townsend and P. Calow). Blackwell Science Publisher, Oxford, 1981.
- [64] A. Putter. Studies on the physiological similarity. VI. Similarities in growth. *Pflugers* archiv fur die gesamte physiologie des menschen und der tiere, 180:280, 1920.
- [65] S. Richman. The transformation of energy by *Daphnia pulex*. *Ecological Monographs*, 28:273, 1958.
- [66] R. D. Roberts, C. Lapworth, and R. J. Barker. Effect of starvation on the growth and survival of post-larval abalone (*Haliotis iris*). *Aquaculture*, 200:323–338, 2001.
- [67] C. Romijn and W. Lokhorst. Foetal respiration in the hen the respiratory metabolism of the embryo. *Physiologia Comparata et Oecologia*, 2:187–197, 1951.
- [68] D. A. S. Rosen and A. W. Trites. Heat increment of feeding in Steller sea lions, *Eumetopias jubatus*. *Comparative Biochemistry and Physiology A-Physiology*, 118:877–881, 1997.
- [69] J. L. Ross, T. M. Stevens, and D. S. Vaughan. Age, growth, mortality, and reproductive-biology of red drums in North Carolina waters. *Transactions of the American Fisheries Society*, 124:37–54, 1995.
- [70] N. R. Russell and R. J. Wootton. Appetite and growth compensation in the european minnow, *Phoxinus-phoxinus* (cyprinidae), following short periods of food restriction. *Environmental Biology of Fishes*, 34:277–285, 1992.

- [71] V. M. Savage, J. F. Gillooly, W. H. Woodruff, G. B. West, A. P. Allen, B. J. Enquist, and J. H. Brown. The predominance of quarter-power scaling in biology. *Functional Ecology*, 18:257–282, 2004.
- [72] C. C. Schwartz and K. J. Hundertmark. Reproductive characteristics of Alaskan moose. *Journal of Wildlife Management*, 57:454–468, 1993.
- [73] J. L. Seale, W. V. Rumpler, and Moe P. W. Description of a direct-indirect roomsized calorimeter. *American Journal of Physiology*, 260(2):E306–E320, 1991.
- [74] R. Shine and J. B. Iverson. Patterns of survival, growth and maturation in turtles. *Oikos*, 72:343–348, 1995.
- [75] R. Steenbergen, T. S. Nanowski, R. Nelson, S. G. Young, and J. E. Vance. Phospholipid homeostasis in phosphatidylserine synthase-2-deficient mice. *Biochimica et Biophysica Acta - Lipids and Lipid Metabolism*, 1761(3):313–323, 2006.
- [76] B. A. Stockhoff. Starvation resistance of gipsy moth, *Lymantria dispar* (l.) (lepidoptera: Lymantriidae): tradeoffs among growth, body size and survival. *Oecologia*, 88:422–429, 1991.
- [77] T. Stromgren and C. Cary. Growth in length of *Mytilus edulis-l* fed on different algal diets. *Journal of Experimental Marine Biology and Ecology*, 76:23–34, 1984.
- [78] S. C. Strum. Weight and age in wild olive baboons. American Journal of Primatology, 25:219–237, 1991.
- [79] C. Tolla. *Modelling of Microbial Population Dynamics in variable Environments*. PhD thesis, University of Marseille & Vrije Universiteit, Amsterdam, 2006.
- [80] C. Tolla, S. A. L. M. Kooijman, and J. C. Poggiale. A kinetic inhibition mechanism for the maintenance process. *Journal of Theoretical Biology (submitted)*, 2006.
- [81] J. van der Meer. An introduction to Dynamic Energy Budget (DEB) models with special emphasis on parameter estimation. *Journal of Sea Research*, 56:85–102, 2006.
- [82] J. van der Meer. Metabolic theories in ecology. *Trends in Ecology and Evolution*, 21:136–140, 2006.

- [83] H. W. van der Veer and M. Alunno-Bruscia. Journal of Sea Research Dynamic Energy Budgets in Bivalves, volume 56. Elsevier, 2006.
- [84] H. W. van der Veer, S. A. L. M. Kooijman, and J. van der Meer. Body size scaling relationships in flatfish as predicted by Dynamic Energy Budgets (DEB theory): implications for recruitment. *Journal of Sea Research*, 50:255–270, 2003.
- [85] H. W. van der Veer, S. A. L. M. Kooijman, and J. van der Meer. Intra- and interspecific comparison of energy flow in North Atlantic flatfish species by means of Dynamic Energy Budgets. *Journal of Sea Research*, 45:303–320, 2001.
- [86] J. Vidal and T. E. Whitledge. Rates of metabolism of planktonic crustaceans as related to body weight and temperature of habitat. *Journal of Plankton Research*, 4:77–84, 1982.
- [87] P. J. Whitehead. Respiration of *Crocodylus johnstoni* embryos. In *Wildlife Management: Crocodiles and alligators*, Webb, G.J.W. and Manolis, S.C. and Whitehead, P.J. (Eds), pages 173–497. Sydney: Beatty, 1987.
- [88] H. P. Zheng, C. H. Ke, S. Q. Zhou, and F. X. Li. Effects of starvation on larval growth, survival and metamorphosis of ivory shell *Babylonia formosae habei* Altena et al., 1981 (Neogastropoda : Buccinidae). *Aquaculture*, 243:357–366, 2005.
- [89] C. Zonneveld and S. A. L. M. Kooijman. Comparative kinetics of embryo development. *Bulletin of Mathematical Biology*, 55:609–635, 1993.